



Trade Science Inc.

# Organic CHEMISTRY

*An Indian Journal*
*Full Paper*

OCAIJ, 7(6), 2011 [376-379]

## A new method of deprotection for the hydroxide radicals: Boron trifluoride etherate/sodium iodide

Yu-Qing Cao\*, Xiao-Jun Yang, Xiang-Tao Xu, Ding-Xiang Du, Li-Ya Xu, Fang-Rui Song  
College of Pharmacy, Hebei University, Baoding - 071 002, (P.R.CHINA)

E-mail: coffeeyang2017@163.com

Received: 13<sup>th</sup> March, 2011 ; Accepted: 23<sup>rd</sup> March, 2011

### ABSTRACT

A new efficient method for dealkylation of ethers is reported. Ethers could transform into corresponding alcohols with boron trifluoride etherate and sodium iodide in acetonitrile after hydrolysis. This reaction can proceed at room temperature, and the yield is excellent.

© 2011 Trade Science Inc. - INDIA

### KEYWORDS

Boron trifluoride etherate;  
Sodium iodide;  
Deprotection;  
Ethers.

### INTRODUCTION

Hydroxide radical is very common in organic compounds, especially in nature extract, it could be transformed into ester, halogenated hydrocarbon, aldehydes, ketone, carboxylic acid and unsaturated hydrocarbon easily. It is necessary to protect the hydroxide radical from transforming when we do not want the hydroxide radical change. Alkylation of a hydroxyl moiety is regarded as one of the most effective protection methodologies. At last we need to process a deprotection step to return the hydroxide radical. It's useful in both commercial and academic pursuits, such as natural product, fine chemical and pharmaceutical syntheses.


Many researchers bring forward their means to accomplish this dealkylation reaction since mid-20th century. Lewis acids are usually used in the dealkylation reaction. The first reagent for dealkylation is  $\text{BBr}_3$ ,<sup>[1]</sup> but the usage of boron tribromide is limited for its intensive toxicity and explosibility when heated. Later people found some other dealkylation reagents,  $\text{Me}_3\text{SiI}$ <sup>[2]</sup> was used in this reaction more common, such as  $\text{Me}_3\text{SiI}/\text{CDCl}_3$ ,<sup>[3]</sup>

$\text{Me}_3\text{SiI}/\text{Pyridine}$ <sup>[4]</sup>,  $\text{Me}_3\text{SiCl}/\text{NaI}$ .<sup>[5]</sup> Olivero<sup>[6]</sup> found the electrochemistry means could transform some ethers into alcohols too. Except these reagents,  $\text{OsO}_4/\text{N}$ -methylmorpholine N-oxide,<sup>[7]</sup>  $\text{NaBH}_4/\text{THF}$ ,<sup>[8]</sup>  $\text{SmI}_2/\text{THF}$ ,<sup>[9]</sup>  $\text{Pd}(\text{PPh}_3)_4/\text{MeOH}$ <sup>[10]</sup> were test and verify to be feasible for some specially ethers. Shanthaveerappa<sup>[11]</sup> used the high nucleophilicity of bromide ion in the form of ionic liquid, 1- n -butyl-3-methylimidazolium bromide, for the nucleophilic displacement of an alkyl group to regenerate a phenol from the corresponding aryl alkyl ether. This is a novel method and the yield is excellent, but the ionic liquid make the process too complicated.

Some ethers with special substitutional groups may need special reagent combination.  $\text{Ce}(\text{OTf})_4$  could react with triphenylmethyl ethers and changed them into corresponding alcohols in mild conditions.<sup>[12]</sup> Silyl ethers could changed into corresponding alcohols with sodium periodate in excellent yield.<sup>[13]</sup> t-Butyldimethylsilyl ethers were efficiently and selectively deprotected using silica supported sodium hydrogen sulfate ( $\text{NaHSO}_4 \cdot \text{SiO}_2$ ) as a heterogeneous catalyst at room temperature to regenerate the parent alcohols in high yields.<sup>[14]</sup> Shino

found that the propargyl ethers could be processed reductively deprotection reaction with  $\text{SmI}_2$ -amine-water system.<sup>[15]</sup> Farhad developed a method for direct oxidation of trimethylsilyl, tetrahydropyranyl and methoxymethyl ethers to their corresponding carbonyl compounds using trinitratocerium bromated supported on  $\text{NaHSO}_4 \cdot \text{H}_2\text{O}$  under solvent-free conditions.<sup>[16]</sup> 1-Decanethiol has been found to be an excellent reagent for the deprotection of aryl methyl ethers. A clear advantage of 1-decanethiol over the more commonly used thiols is the easy extraction of both the deprotecting reagent and the reaction byproducts into the aqueous phase, it could make the process more simple.<sup>[17]</sup>

We now report that a combination of boron trifluoride etherate/sodium iodide in acetonitrile, it could react with several kinds of ethers, then hydrolysis into alcohols. It is a mild, low-cost reagents system, the yield is satisfying.



Scheme 1

















## EXPERIMENTAL

The reaction is monitored with TLC (thin layer chromatography) and GC (gas chromatography). Melting points were determined on a microscopy apparatus (SGW X-4) and uncorrected. All the liquid parent materials were freshly distilled. The products were characterised by comparison of their melting points and boiling points with the literature values.








## General procedure

To a stirred solution of ether (10mmol) and sodium iodide (20mmol, 3.0g) in dry acetonitrile (25ml) is added through a constant pressure funnel boron trifluoride etherate (20mmol, 5.42ml) smoothly and the mixture is stirred at room temperature. The mixture is then treated with saturated sodium hydrogen carbonate solution (25ml) and extracted with ether (3×10ml). The organic layer is washed with aqueous sodium thiosulfate solution (20ml) followed by water (50ml), dried with anhydrous sodium sulfate, and evaporated. The residue is then taken in ether (25ml) and triturated. The ether layer is decanted from the precipitated salt; and distilled to yield the pure alcohol.

TABLE 1 : Dealkylation of ethers

Entry	Substrate	Product	Time (h)	Yield (%)	m.p.(°C)	
					Found	Reported <sup>C</sup>
1			8	82	39-40 <sup>a</sup>	40-42
2			18	47	193-195 <sup>a</sup>	195
3			8	85	30-32 <sup>a</sup>	32-34
4 <sup>b</sup>			8	87	170-172 <sup>a</sup>	172-175
5			18	38	39-40 <sup>a</sup>	40-42
6			8	83	121-123 <sup>a</sup>	122-123
7			6	89	157-159	160-161
8			6	94	158-159	160-161

## Full Paper

Entry	Substrate	Product	Time (h)	Yield (%)	m.p.(°C)	
					Found	Reported <sup>c</sup>
9			6	95	137-139	139-140
10			6	95	38-40 <sup>a</sup>	38-40
11			6	94	103-104	104
12			6	91	195-196	196
13			6	92	176-179	177-180

<sup>a</sup>Determination of melting point; <sup>b</sup>The proportion of the reagent is ether: sodium iodide: boron trifluoride etherate=1:4:4; <sup>c</sup>The Merck Index, 2006, 13th Edn.

## RESULTS AND DISCUSSION

We attempt to add the reagents at different scale and in different solvents, at last we found the best proportion is Ether: NaI: BF<sub>3</sub>·Et<sub>2</sub>O=1: 2 : 2, and the most suitable solvent is acetonitrile.


As the results show in TABLE 1, we could see most of the ethers could convert to the corresponding alcohols or phenols in excellent yield, the aliphatic ethers need shorter time and the yield is higher than the aromatic ethers some ethers with special groups behave sluggishly comparatively.

The electron pair on the oxygen atom that attach with benzene ring direct could conjugate with the benzene ring, which would reduce the activity of this oxygen atom, make it hard to combine with Lewis acid BF<sub>3</sub> (entry 1-6, TABLE 1). An electron-withdrawing group on the benzene ring can reduce the activity of oxygen atom, make the reactions proceed more difficult (entry 2, TABLE 1). The electron donating group on the benzene ring can make the reactions proceed easier (entry 3-4, TABLE 1).

For the asymmetry ethers, the C-O bond rupture has selectivity between the two side of oxygen atom. Generally speaking, the rupture order of C-O bond from easy to hard is tertiary carbon, secondary carbon, primary carbon and aryl-carbon. Because the electron pair on the oxygen atom conjugate with the benzene ring, the C-O bond between benzene and oxygen atom has some characteristic of double bond, so this bond rupture more difficult than the other one. This is why the

product of entry 5 in TABLE 1 is phenol, but not isoamylol.

The mechanism of this reaction we conjecture: firstly, BF<sub>3</sub> combine with oxygen atom of ethers to generate a complex compound, then the iodine ion attack this complex compound, which led to the bond breaking between carbon and oxygen atom and generate a new compound A, A transform into alcohols or phenols through hydrolysis (Scheme 2).



Scheme 2

## CONCLUSION

In summary, a new method for the conversion of ethers into alcohols or phenols using boron trifluoride etherate-sodium iodide-acetonitrile system has been developed. This method is inexpensive, practical and with less pollution and easy work-up.

## REFERENCES

- [1] M.Felix; J.Org.Chem., **39**, 1427 (1974).
- [2] F.L.Ho, G.A.Olah; Angew.Chem.Int.Ed., **15**, 774 (1976).

- [3] M.E.Jung, M.A.Lyster; *J.Org.Chem.*, **42**, 3761 (1977).
- [4] M.E.Jung, M.A.Lyster; *Organic Syntheses*, **59**, 35 (1979).
- [5] G.A.Olah, S.C.Narang, B.G.Gupta; *J.Org.Chem.*, **44**, 1247 (1979).
- [6] S.Olivero, E.Dunach; *Chem.Commun.*, **24**, 2497 (1995).
- [7] H.B.Mereyala, S.R.Gurralla, S.K.Mohan; *Tetrahedron.*, **55**, 11331 (1999).
- [8] K.P.Chary, G.H.Mohan, D.S.Iyengar; *Chem.Lett.*, **28**, 1223 (1999).
- [9] A.Dahlen, A.Sundgren, M.Lahmann, S.Oscarson, G.Hilmersson; *Org.Lett.*, **5**, 4085 (2003).
- [10] D.R.Vutukuri, P.Bharathi, K.Rajasekaran, M.Tran, S.Thayumanavan; *J.Org.Chem.*, **68**, 1146 (2003).
- [11] S.K.Boovanahalli, D.W.Kim, D.Y.Chi; *J.Org.Chem.*, **10**, 3341 (2004).
- [12] K.N.Ali, F.A.Reza; *Tetrahedron.*, **57**, 6805 (2001).
- [13] W.Mijuan, L.Chun, Y.Dali, L.Xiaotian; *Tetrahedron Letters*, **43**, 8727 (2002).
- [14] M.Shino, U.Akiharu, I.Yukishige; *Tetrahedron Letters*, **49**, 5159 (2008).
- [15] S.Farhad, S.Saeideh; *Chinese Chemical Letters*, **19**, 676 (2008).
- [16] D.Biswanath, R.K.Ravinder, P.Thirupathi; *Tetrahedron Letters*, **47**, 5855 (2006).
- [17] K.Bhima, S.Ananta, S.Bapurao, G.Samir; *Tetrahedron Letters*, **51**, 3075 (2010).