



ANALYSIS OF OIL EXTRACTED FROM *ANETHUM SOWA*

NAMRATA VATS* and RAGINI PANDEY^a

R.S.K.D. P.G. College, JAUNPUR (U.P.) INDIA

^aIndian Institute of Medical & Paramedical Sciences, JAUNPUR (U.P.) INDIA

ABSTRACT

The dried powder of *Anethum sowa* extracted for cold percolation followed by hydro distillation. The oil obtained by the hydro-distillation method from the fresh leaves and stems of *Anethum sowa* herb was analyzed by GC-MS. Twenty compounds were isolated and identified. The major constituents were apiole (26.15%), o-cymene (13.76%), α -thujene (16.53%), β -phellandrene (7.17%), 6,6-dimethyl-2-(3-oxobutyl) bicyclo (3.1.1) heptan-3-one (7.03%), exo-2-hydroxycineol (4.98%), limonene (4.16%), 3-isopropyl-4-methyl-1-pentyn-3-ol (2.89%), myristicine (2.16%) and dithydroumbellulone (2.19%).

Key words: *Anethum sowa* L., Essential oil, GC-MS, Apiole.

INTRODUCTION

Anethum sowa L. (Hindi-Sowa) belongs to the family *Apiaceae* (*Umbelliferae*) has a primitive herb use as vegetable. It is an annual, glabrous aromatic cold weather condiment crop in northern plains of India¹. *Anethum sowa* has also been some-times referred to as a variety of *Anethum graveolens*. Its fruits are longer but less fragrant than *Anethum graveolens* L. (*Umbelliferae*). The Dill herb grows up to 76 cm in height, which has small feathery leaves and stands on sheathing foot-stalks with linear and pointed leaflets. Stem is erect, branched, cylindrical, striated, smooth and pale green^{2,3}. Both seed and freshly cut entire herb including stem, leaf and fruit are widely used for flavoring of food, beverages, fragrance in cosmetics and in various medicinal preparations. The oil and its emulsion in water are used in gastrointestinal disorders and an important ingredient of gripe water^{2,3}. Biological and pharmacological studies of this plant revealed antimicrobial, anti-oxidative and antispasmodic activities^{4,5}.

Essential oils of Dill herb and seed have been reported by many researchers⁶⁻⁹. The purpose of this study was to determine the chemical constituents of Dill herb essential oil,

* Author for correspondence; E-mail: dr.girija59@yahoo.com

the content of aroma causing compounds as well as to determine the feasibility of large scale herb production in eastern U.P. India. The literature search shows that, there is no previous report on the chemical composition of the essential oil of Dill herb grown in North India.

EXPERIMENTAL

Materials and methods

The fresh Dill herb was collected from the district of Jaunpur (U.P.) India in January 2012. The plant was identified by Deptt. of Taxonomy, Kuteer P.G. College Chakke, Jaunpur and authenticated with preserved sample under supervisor of Dr. R. A. Singh.

The essentials oil of leaves and stems were obtained by hydro-distillation method from finely chopped fresh herb after separation of its root in a Clevenger type apparatus for 2 hours. The oil was collected, dried over anhydrous sodium sulphate and refrigerated at 6°C. The yield of the oil was 0.35% (w/w) on fresh weight basis.

GC-MS analysis

The analysis of the oils was carried out by GC-MS electron impact ionization method on GC-17a gas chromatograph (Shimadzu) with FID detector coupled to a GC-MS QP-5050A mass spectrometer (Shimadzu); fused silica capillary column (30 m × 0.25 mm film thickness) coated with DB-1 (J & W). Column temperature 40°C (2 min.) was raised to 250°C at the rate of 5°C/min. Injection port temperature was 350°C and injection volume 0.1 µL. Helium gas was used as carrier gas at constant pressure of 100 Kpa, flow rate 20 mL/min. Acquisition parameters full scan : scan range 40-450 amu. Identification of the compounds were confirmed by computer matching of their mass spectral fragmentation pattern with those of compounds in NIST (National Institute of Standard and Technologies) 107 built in Lis Libraries Shimadzu Corporation, Japan. Sample was dissolved in chloroform. Mass spectra were recorded at 70 ev. The GC-MS data of the oil are shown in Table 1.

Table 1: Chemical composition of the Dill herb (leaf and stem) essential oil

Compounds	Relative %
α -pinene ^c	0.98
α -thujene ^c	16.53
o-cymene ^c	13.76
β -phellandrene ^c	7.17

Cont...

Compounds	Relative %
Limonene ^c	4.16
Terpinolene	1.53
Undecane ^f	1.56
Dill ether ^b	1.64
1,3-cyclohexadiene,2-methyl-5-(1-ethylethyl)- monoepoxide ^b	1.82
6,6-dimethyl-2-(3-oxobutyl) bicycle (3.1.1) heptan-3-one	7.03
Camphenol-6 ^b	1.34
Lilac alcohol B ^b	1.17
exo-2-hydroxycineole ^b	4.98
3-isopropyl-4-methyl-1-pentyn-3-ol ^g	2.89
Dihydroumbellulone ^b	2.19
Myristicin ^a	2.16
Apiol ^a	26.15
α -selinene ^d	1.36
Phytol ^e	1.58

Notes : Compounds arranged in the order of the peak numbers. Superscripts a, b, c, d, e, f and g indicates for phenyl-propanoid, oxygenate monoterpenes, monoterpene hydrocarbons, sesquiterpene, diterpene, alkanek, alcohol constituents respectively.

RESULTS AND DISCUSSION

In the study Dill yielded 0.32% (w/w) light yellow coloured oil with a typical herb odor on fresh weight basis. It is reported in the literature that the oil percentages ranges from 0.1% to 0.46% on fresh weight basis^{10,11}.

Nineteen constituents have been identified in the oil by GC and GC-MS (Table 1). The constituents are listed in the order of their peak. The oil is characterized by fairly high concentration of monoterpenes (56.87%) comprising of monoterpene hydrocarbons (44.01%) and oxygenated monoterpenes (12.86%). Ondly one sesquiterpene a-selinene (1.08%) consisted of the sesquiterpene hydrocarbon, a diterpene compound phytol (1.58%) consisted

of the oxygenated diterpene, phenylpropanoid consists apiole (26.15%) and myristicine (2.16%) were identified in the oil. Undecane (1.56%) and 3-isopropyl-4-methyl-1-pentyn-3-ol (2.89%) were also found in the investigation as alkane and alcohol type compounds, respectively.

Other components were present in an amount of 8.42%. Beside these, the predominant components were; o-cymene (13.76%), α -thujene (16.53%), β -phellandrene (7.17%), 6,6-dimethyl-2-(3-oxobutyl)bicycloheptan-3-one (3.11%), heptan-3-one (7.03%), exo-2-hydroxycineole (4.98%), limonene (4.16%), and dihydroumbellulone (2.19%). Apiole an undesirable toxic constituent was found in the highest amount of the oil. This ingredient brings down its quality. This constituent could be separated or removed from the oil by fractional distillation¹². Though it is toxic in nature it can be used as synergist with insecticides. It has also shown a stronger oviposition deterrent against *Callosobruchus maculatus* than^{13,14}.

The literature survey revealed that the compounds like α -thujene (0.19-0.41%), α -pinene (0.98-2.16%), β -phellandrene (11.87-19.08%), dill ether (16.62-30.18%) and myristicine (0.01-0.19%) were found in the herb oil¹⁰. In another publication¹⁵, the compounds α -pinene (1.2-1.4), β -phellandrene (6.4-8.5%), dill ether (21.6-37.7%), limonene (2.1-2.8%) and myristicine (0.3-2.1%) were reported. In the present study, almost the similar compounds as the earlier reports^{10,15} were found. In addition to that, another 5 compounds like (1) phytol, (2) α -selenene, (3) 6,6-Dimethyl-2-(3-oxobutyl) bicycloheptan-3-one, (4) 3-Isopropyl-4-methyl-1-pentyn-3-ol and (5) cyclohexanone, 2-isopropyl-2,5-dimethyl were also found. It appears that these five compounds were first identified in this study.

CONCLUSION

The results of this study are of interest in respect to the processing of dill herb. The distillation and the drying process are to be performed in such a way that minimum loss of volatile components occurs. These essential oils have the potential to be used in food as flavoring and natural preservatives, pharmaceuticals, cosmetics and may also be considered as active ingredients in medical preparations.

REFERENCES

1. A. Husain, O. P. Virmani, A. Sharma and L. N. Misra, Major Essential Oil-Bearing Plants of India, CIMAP, Lucknow (1988) pp. 87-95.
2. Anonymous, The Wealth of India, Raw Materials, Council of Scientific and Industrial Research, New Delhi, India, **Vol. 1-A** (1985) pp. 272-275.

3. R. N. Chopra, I. C. Chopra, K. L. Honda and L. D. Kapur, Chopra's Indigenous Drugs of India, 2nd Ed., Academic Publishers, New Delhi, India (1982) pp. 692-693.
4. B. K. Baslas and R. K. Baslas, Chemical Studies into the Essential Oils from the Plants of *Anethum Graveolens* & *Anetham Sowa* (Dill Oils), Indian Perfumer, **15**, 27-29 (1971).
5. A. Woolf, Essential Oil Poisoning, J. Clinical Toxicol., **37**, 721-727 (1999).
6. E. Guenther, The Essential Oils, **Vol. IV**, 1st Edn., D. Van Nostrand Co. Inc. New York (1953) p. 633.
7. B. Lawrence, New Trends in Essential Oils, Perfume Flavour, **5**, 6-16 (1980).
8. H. Husain, Status Report on Aromatic and Essential Oil-Bearing Plants in NAM Countries Publication and Information Directorate (CSIR), **1**, 71-72 (1994).
9. A. Ahmed, L. N. Misra and M. C. Nigam, A Dihydrobenzofuran from India Dill Seed Oil, J. Phytochemistry, **29(6)**, 2035-2037 (1990).
10. J. C. Denys, E. S. James and W. Markp, Characterization of Essential Oil of Dill (*Anethum graveolens* L) J. Essent Oil Res., **7**, 11-20 (1995).
11. S. E. El-Gengaihi and L. Hornok, The Effect of Plant Age on Content and Composition of Dill Essential Oil *Anethum Graveolens* L. Acta Horticult., **73**, 213-218 (1978).
12. N. B. Shankaracharya, R. L. Mohan, J. Puranaik, J. and S. Nagalakshmi, Studies on Chemical and Technological Aspects of Indian Dill Seed (*Anethum Sowa*, Roxb.), J. Food Sec. Technol., **37(4)**, 368-372 (2000).
13. S. S. Tomar, M. L. Maheshwari and S. K. Mukerjee, Synthesis and Synergistic Activity of Dillapiole Based Pyrethrum Synergists, J. Agric. Food Chem., **27(3)**, 547 (1979).
14. A. K. Tripathi, V. Prajapati, K. K. Aggarwal and S. Kumar, Insecticidal and Ovicidal Activity of the Essential Oil of *Anethum Sowa* Kurz Against *Callosobruchus Maculatus* F. (Coleoptera : Bruchidae) Insect Sci. Applic., **21(1)**, 61-66 (2001).
15. S. Halva, R. Huopalahti and M. Franzch, Herb Yield and Essential Oil of Dill (*Anethum Graveolens* L.) at Different Locations, J. Agric. Sci., Finland, **60**, 93-100 (1988).