

Design Synthesis and Preliminary Antibacterial Evaluation of Schiff Base Metal Complexes of Ampicillin and Cephalexin

Abbas H Abdulsada*, Noor H Nasser and Ahmed K Hussein

Department of Pharmaceutical Chemistry, University of Kufa, Najaf, Iraq

*Corresponding author: Abbas H Abdulsada, Department of Pharmaceutical Chemistry, University of Kufa, Najaf, Iraq, E-Mail: abhart80@gmail.com

Received: January 19, 2017; Accepted: February 19, 2018; Published: March 13, 2018

Abstract

In the present study, new Schiff bases derived from two beta lactam antibiotics, ampicillin and cephalexin were synthesized from condensation of the aforementioned drugs with Terephthalaldehyde and their metal complexes counterparts with copper (II) have been synthesized. In both drugs the free amino group of the acylamino side chain allowed to react with Terephthalaldehyde to obtain Schiff bases which were, subsequently, allowed to react with copper chloride separately to form Schiff base copper complexes. All the derivatives and metal complexes are analyzed by FT-IR, elemental analysis and ¹H NMR analysis. The antibacterial activity of the prepared compounds was screened against several bacterial strains namely S. aureus, E. coli and P. aeruginosa. The antibacterial screening tests elicit that both copper complexes besides Schiff bases exhibited promising antibacterial activity against these microorganisms so that they are of improved antibacterial activity than the parent drugs.

Keywords: Schiff base; Ampicillin; Cephalexin; Metal complexes; Terephthalaldehyde.

Introduction

Schiff bases are compounds that are synthesized by condensation reaction of primary amines with carbonyl groups [1,2]. In the past many Schiff base derivatives have been prepared and employed for applications like catalysis and enzymatic reactions, luminescent material, magnetism and molecular architectures [3]. However recently Schiff bases metal complexes gain massive attention in the domain of biological chemistry and coordination chemistry [4].

The incidence of infection has increased significantly as a result of multidrug resistance which gain a great clinical significant that motivate additional urgency for finding out a novel antibacterial agent which preferably act by a different mechanism of action or by broadening the antibacterial spectrum of the current antibiotics for example by forming a mutual prodrug [5,6].

The terrifying development of outbreaks by multidrug-resistant organisms constitute a great challenge when treating these infections, as these isolates often respond to a limited number of antibiotics and are in most cases life threatening [1,7,8].

Schiff bases are among the most efficient strategies tried to enhance the antimicrobial activity of many compounds including antibiotics and/or extend the spectrum of that individual antibiotic to include a resistant bacterial strain or a common bacterial pathogen [9-14].

Schiff's bases represent an important class of pharmacologically active molecules so that have engaged a great concern by medicinal chemist as they possess a variety of pharmacological properties. *A number of Schiff's base derivatives have been reported to exert notably antibacterial, antifungal, antitubercular, antitumor, antileishmanial, DNA-binding activities, etc*

[6,15-18]. It is believed that different biological activity of Schiff bases like antitumor, antibacterial, antifungal and herbicidal activities are attributed to the azomethine linkage [19,20].

Many Schiff base derivatives of β -lactam antibiotics have been synthesized and screened for their antimicrobial activity in order to enhance their potency and/ or extend their spectrum of antibacterial activity; Schiff base ligands and their metal complexes derived from different aldehydes with cefadroxil [21], cephadrine [2,22], ceftriaxone [2,23] and cephalothin [24], amoxicillin [25-29] have been synthesized and generally found that there is an enhancement in the activity of the prepared ligands and their metal complexes counterparts over the parent drug.

Ampicillin and cephalixin have been synthesized as metal complexes and as Schiff base derivatives by combining the drug molecule with different aldehydes, metal complexes for the prepared Schiff bases also synthesized. The prepared compounds were tested for their prospective antibacterial properties [30-50]. Pharmacological research interest in the design of metal–drug complexes since it was found that metal – drug combination help boost its activity and lower the incidence of toxic effects [19].

Experimental

Materials

All chemicals and solvents used were of analytical grade (Merk, Fluka and Sigma- Aldrich) and were used without further purification.

Measurement

Melting points were measured by open capillary method and are reported uncorrected (**TABLE 1**). IR spectra were obtained on Perkin Elmer IR spectrophotometer (KBr disc) in faculty of pharmacy/ university of kufa (**TABLE 2**). Elemental microanalysis was accomplished at the Jordanian University using CHN Elemental Analyzer (Euro-vector EA3000A, Italy) (**TABLE 1**). Nuclear magnetic resonance spectrum ^1H NMR for Schiff base and copper complexes was recorded in DMSO- d_6 using a Bruker 500 MHz instrument using a tetra methyl silane (TMS) as an internal standard in Terbia Modares University.

Preparation of schiff base ligands

Two Schiff base ligands were produced by reacting the drug substances with Terephthalaldehyde separately. The method of preparation was as follows. The solution of Terephthalaldehyde in absolute methanol was appended drop wise into 25 mL of either cephalixin or ampicillin in the same solvent with persistent stirring in 1:2 molar ratios. To this, KOH (0.1% in methanol) was appended to adjust the PH of the medium between pH 7 and 8 and the mixture was kept under reflux for 5 hours. After that the crude mixture has condensed to about one-third of its volume by vacuum evaporation. A bright yellow colored Schiff base was extracted by the addition of diethyl ether which was then filtered, followed by the addition of methanol and then drying under vacuum. Recrystallization was done in methanol.

TABLE 1. Physical properties and micro analytical data of ligand and its metal complexes.

| Compound | Color | Melting point | Percent of yield | Elemental analysis found (calculated) | | | |
|---------------------------------------|-------------|---------------|------------------|---------------------------------------|----------------|------------------|----------------|
| | | | | C | H | N | M |
| terephthalaldehydeampicillin | yellow | 229 | 79 | 60.83 (60.29) | 4.97 (5.06) | 10.19 (10.55) | - |
| terephthalaldehydecephalexin | yellow | 195 | 74 | 60.15 (60.59) | 4.22 (4.58) | 10.78 (10.60) | - |
| Terephthalaldehydeampicillin- Cu (II) | green | 318 | 68 | 51.94 (51.58) | 4.54 (4.33) | 9.37 (9.02) | 6.38 (6.82) |
| Terephthalaldehydecephalexin- Cu(II) | Light green | 274 | 66 | 52.07 (51.81) | 3.59 (3.91) | 9.33 (9.06) | 6.48 (6.85) |

FIG.1. Synthesis of terephthalaldehydecephalexin.





FIG.2. Synthesis of terephthalaldehydeneampicillin.



Preparation of Schiff base copper complexes

2 mmol of the previously prepared Schiff base ligand dissolved in methanol (25 ml) then added to 25 ml of methanol containing 2 mmol of copper chloride Cu (II). The mixture was then kept under reflux for 4 h. Then reducing the mixture volume under vacuum, filtering it off, washing with methanol and finally drying under vacuum.

Antimicrobial activity

The prepared Schiff bases along their metal complexes were screened for their potential activity against some selected microorganisms, especially against *Escherichia coli*, *staphylococcus aureus*, and *Pseudomonas aeruginosa* by measuring the zone of inhibition in culture medium in mm. The antibacterial activity was performed according to filter paper disc plate method at concentration 100 $\mu\text{g/mL}$ and the results reported in **TABLE 3**. The culture medium was Muller Hinton agar & Sabouroud Dextrose agar while the solvent control was DMSO.

Results and Discussion

The Schiff base metal complexes were obtained by reacting with copper chloride with Schiff base ligands in methanol under reflux separately. The solubility of the intermediates and final products was assayed in water and certain organic solvents. The ligands were soluble in water, methanol, DMF (Dimethylformamide) and DMSO (Dimethyl sulfoxide), while metal complexes were soluble in DMF and DMSO but not dissolve in more nonpolar organic solvents. The Schiff bases and the copper complexes were sufficiently hygroscopic.

It is possible to explore the bonding pattern of ligands to the metal elements from the comparison of FT-IR spectral data of the Schiff bases to those of copper complexes counterparts. These data indicate the formation of Schiff base ligands through the $-C=N-$ bond and ligands coordinate with copper ion through nitrogen and oxygen.

TABLE 2. IR spectral data of the synthesized compounds.

| Compound | C=O | C=N | M-N | M-O |
|--------------------------------------|------|------|-----|-----|
| Terephthalaldehydeampicillin | 1672 | 1638 | - | - |
| Terephthalaldehydecephalexin | 1668 | 1632 | - | - |
| Terephthalaldehydeampicillin-Cu (II) | 1655 | 1618 | 483 | 436 |
| Terephthalaldehydecephalexin-Cu(II) | 1643 | 1615 | 507 | 439 |

TABLE 3. The antibacterial activity of ligands and complexes.

| Compound | <i>E. coli</i> | <i>S. aureus</i> | <i>P. aeruginosa</i> |
|--------------------------------------|----------------|------------------|----------------------|
| Terephthalaldehydeampicillin | 25 | 29 | 18 |
| Terephthalaldehydecephalexin | 27 | 27 | 16 |
| Terephthalaldehydeampicillin-Cu (II) | 28 | 32 | 20 |
| Terephthalaldehydecephalexin-Cu(II) | 29 | 30 | 15 |

The absorption bands for azomethine group of the Schiff base ligands appears in the range of 1620-1650 while those for metal complexes are slightly revolved to slightly higher values denoting the coordination of Schiff bases with metal ion through $-N=C$ which contrary to bands related to the carboxylic acid group which remain in its range as this group does not participate with the metal ion. A new absorption bands exist in the range of 480- 510 for (M-N) and 415- 450 for (M-O) indicating the coordination of the ligands with the metal ion through N and O atoms.

The 1H NMR spectrum of terephthalaldehydeampicillin show a signal that is related to the chemical shift of the two methyl protons on the Thiazolidine ring existed at δ 1.2 and 1.39 ppm. The protons of N-CH and CO-CH on the beta-lactam ring and NH proton of amide have chemical shifts of δ 4.75, 4.55 and 8.23 ppm respectively. The azomethine proton appeared at δ 8.45 ppm while aromatic protons found at 6.85- 7.97 (m). No signal in the chart attributed to the chemical shift of the free amino group of ampicillin as it takes part in azomethine group formation. The 1H NMR spectrum of terephthalaldehydeampicillin Cu (II) complex in DMSO exhibited signals belong to chemical shifts of the protons of aromatic moiety, protons of CO-CH and N-CH on the beta-lactam ring observed at δ (6.68-7.76), 4.71, 4.95 ppm respectively. While signals occurred at δ 8.72 and 8.71 are related to the NH proton of the amide and azomethine proton respectively [32,33,51-53].

The ¹H NMR spectrum of terephthalaldehydecephalexin exhibited signals that are related to amide proton, azomethine proton and aromatic protons occurred at δ 9.07, 8.65 and (6.71-7.85) respectively. The chemical shifts occurred at δ 2.25, 4.52 and 5.03 were attributed to protons of methyl group, protons of CO-CH and N-CH on the beta-lactam ring respectively. There is no signal to appear for the free amino group of cephalixin as it is utilized in formation of Schiff base. The ¹H NMR spectrum of terephthalaldehydecephalexin Cu (II) complex in DMSO showed signals occurred at δ 2.33, 5.27 and 4.85 and were attributed to chemical shifts of methyl protons, protons of N-CH and CO-CH on the beta-lactam ring respectively. The chemical shift values at δ 9.38, 8.97 and (6.65-7.72) are attributed to the protons of amide, azomethine and aromatic group respectively [32,33,51-53].

Conclusion

Two new Schiff base derivatives of ampicillin or cephalixin with Terephthalaldehyde were synthesized at first then these Schiff bases are complexed with copper to form metal complex, the structures of the metal complexes as well as Schiff bases were proved by elemental analysis, NMR and FTIR spectra. The elemental analysis suggest that the 1:1 (M₂L₂) composition of the metal complexes where "M" depicts metal ion and "L" depicts the free ligand, indicating that the four azomethine groups of the two Schiff base ligands are coordinated with two central metal ions via "N" and "O" atoms, which is advocated by the FT-IR spectra.

REFERENCES

1. Al-Amiery AA, Al-Majedy KY, Ibrahim HH, et al. Antioxidant, antimicrobial, and theoretical studies of the thiosemicarbazone derivative Schiff base 2-(2-imino-1-methylimidazolidin-4-ylidene) hydrazinecarbothioamide (IMHC). *Org Med Chem Lett*, 2012;2.
2. Abdulsada HA, Synthesis Characterization and Antibacterial Evaluation of Cephadrine and Ceftriaxone Schiff Base Copper Complexes. *J Curr Chem Pharm Sci*, 2017;7:102.
3. Zhang X, Han JJ, Huang M, et al. Synthesis, Crystal Structures and Antibacterial Activities of Schiff Base Ligand and Its Cobalt(II) Complex. *Russ J Coordi Chem*, 2012;38:560-566.
4. Dai HC, Mao FL. Synthesis, Structures, and Antibacterial Activity of Schiff_Base Cobalt (III) Complexes, *Russian J Coordi Chem*, 2014; 40:120-124.
5. Shi L, Fang QR, Xue YJ, et al. Synthesis, Characterization, and Antibacterial and Cytotoxic Study of Metal Complexes with Schiff Base Ligands. *Aust J Chem*, 2008;61:288-296.
6. Makawana AJ, Sangani BC, Teraiya BS, et al. Schiff's base derivatives bearing 2-thiophenoxyquinoline nucleus as new antibacterial agents, *Med Chem Res*, 2014;23:471-479.
7. Raman N, Jeyamurugan R, Rajkapoor B, et al. Metal-based antitumor, cytotoxic and antimicrobial activity: pharmacological evaluation of Knoevenagel condensate β -diketone Schiff base thiosemicarbazone Cu (II) and Zn (II) complexes. *Appl Organometal Chem*, 2009;23:283-290.

8. Aziz AA, Naby MAS, Sayed AM, et al. Synthesis, structural characterization, thermal studies, catalytic efficiency and antimicrobial activity of some M(II) complexes with ONO tridentate Schiff base N-salicylidene-o-aminophenol (saphH₂). *J Mole Struc*, 2012;1010:130-138.
9. Sadeek AS, El-Attar MS, El-Hamid AMS. Synthesis, characterization and antibacterial activity of some new transition metal complexes with ciprofloxacin-imine, *Bull Chem Soc Ethiop*, 2015;29:259-274.
10. Kothari R, Sharma B, Sahawal S, et al. Synthesis, characterization and antibacterial evaluation of copper (II) complex with ciprofloxacin antibiotic, *World J Pharm Pharmaceutic Sci*, 2015;4:696-707.
11. Nazir F, Bukhari HI, Aziz MAM, et al. Antibacterial studies and Schiff base metal complexes with some novel antibiotics, *Int J Curr Pharm Res*, 2013;5:40-47.
12. Arshad R, Bukhari HI, Anum F, et al. Synthesis, Spectral and Biological studies of transition metal complexes of schiff base derived from Ofloxacin, *Int. J. Adv. Res. Biol. Sci.* 2016;3:99-108.
13. Krátký M, Dzurková M, Janoušek J, et al. Sulfadiazine Salicylaldehyde-Based Schiff Bases: Synthesis, Antimicrobial Activity and Cytotoxicity. *Molecules*, 2017;22:1573.
14. Chaudhary KN, Mishra P. *In vitro* antimicrobial screening of metal complexes of Schiff base derived from streptomycin and amoxicillin: Synthesis, characterization and molecular modeling. *Am J Appl Chem* 2014;2:19-26.
15. Tantarua G, Popescub CM, Bilda V, et al. Spectroscopic, thermal and antimicrobial properties of the copper(II) complex of Schiff base derived from 2-(salicylidene) aminopyridine. *Appl Organometal Chem*, 2012;26:356-361.
16. Kumar G, Kumar D, Devi S, et al. Synthesis, spectral characterization and antimicrobial evaluation of Schiff base Cu (II), Ni (II) and Co (II) complexes, *European Journal of Medicinal Chemistry*, 2010;45:3056-3062.
17. Nejo AA, Kolawole AG, Nejo OA, Synthesis, characterization, antibacterial, and thermal studies of unsymmetrical Schiff-base complexes of cobalt (II). *J Coord Chem*, 2010;63:4398-4410.
18. Shebl M, Khalil MES, Ahmed AS, et al. Synthesis, spectroscopic characterization and antimicrobial activity of mono-, bi- and tri-nuclear metal complexes of a new Schiff base ligand. *J Mole Struc*, 2010;980:39-50.
19. Raman N, Selvan A, Sudharsan S, Metallation of ethylenediamine based Schiff base with biologically active Cu(II), Ni(II) and Zn(II) ions: Synthesis, spectroscopic characterization, electrochemical behaviour, DNA binding, photonuclease activity and in vitro antimicrobial efficacy, *Spec Chim Acta Part A*, 2011;79:873-883.
20. Etaiw HES, El-Aziz AMD, El-Zaher AHE, et al. Synthesis, spectral, antimicrobial and antitumor assessment of Schiff base derived from 2-aminobenzothiazole and its transition metal complexes. *Spectrochim Acta Part A*, 2011;79:1331-1337.
21. Bukhari HI, Arif M, Nazir F, et al. Synthesis, characterization and antimicrobial studies of Schiff base transition metal complexes of Cr (II), Mn (II), Co (II), Ni (II), Zn (II) and Cd (II) derived from cefadroxil. *Int J Pharmaceu Chem*, 2013;03:17-23.
22. Bukhari IH, Arif M, Akbar J, et al. Preparation, characterization and biological evaluation of schiff base transition metal complexes with Cephradine. *Pak J Biol Sci.* 2005;8:614-617.

23. Fareed G, Kalhoro AM, Versiani AM, et al. 7-[[2-(4-Hydroxyphenyl)methylidene]amino]-1,3-thiazol-4-yl)-2-(methoxyimino)acetyl]amino]-3-[[2-methyl-5,6-dioxo-1,2,5,6-tetrahydro-1,2,4-triazin-3-yl)sulfanyl]methyl]-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic Acid. *Molbank*, 2012.
24. Anacona JR, Noriega N, Camus J. Synthesis, characterization and antibacterial activity of a tridentate Schiff base derived from cephalothin and sulfadiazine, and its transition metal complexes. *Spectrochim Acta A Mol Biomol Spectrosc*, 2015;137:16-22.
25. Chaudhary KN, Mishra P. Metal Complexes of a Novel Schiff Base Based on Penicillin: Characterization, Molecular Modeling, and Antibacterial Activity Study. *Bioinorgan Chem Appl*, 2017.
26. Al-masoudi WA, Mohammad TH, Hama AA. Synthesis characterization and pharmacologic study of new Schiff base derived from amoxicillin drug, *Int Res J Pharm*, 2015;6:386-389.
27. Al-Noor TH, Aziz RM, AL- Jeboori TA. Synthesis, characterization and antimicrobial activities of [Fe(II), Co(II), Ni(II), Cu(II) and Zn(II)] mixed ligand complexes Schiff base derived from amoxicillin drug and 4-(dimethyl amino)benzaldehyde with nicotinamide. *J Chem Pharmaceu Res*, 2014;6:1225-1231.
28. Joshi S, Pawar V, Uma V. Synthesis characterization and biological studies of Schiff base metal complexes Co (II), Zn (II) Ni (II) and Mn (II) derived from amoxicillin trihydrate and various aldehydes, *International Journal of Pharma and Bio Sciences*, 2011;2(1):240-250.
29. Anacona RJ, Marquez V. Synthesis of Some Salicylidene- β -Lactam Antibiotics and Their Antibacterial Activity, *Latin Am J Pharm*, 2013;32:887-91.
30. Kaushal R, Sheetal, Kaushal R, et al. Synthesis characterization and biological screening of titanium (IV) complexes derived from antibiotic drugs. *Int J Pharm Pharmaceu Sci*, 2014;6:374-378.
31. Naz N, Khatoon S, Ajaz H, et al. Synthesis, Spectral Characterization and Biological Evaluation of Schiff Base Transition Metal Complexes Derived from Ampicillin with D-Glucose. *Asian J Chem*, 2013;25 (4):2239-2242.
32. AL-Mudhafar JMM, synthesis characterization and preliminary antimicrobial evaluation of new Schiff bases of ampicillin and amoxicillin derived from isatin derivatives, *Int J Pharm Sci*, 2016;8:113-116.
33. Al-Noor HT, Aziz RM, AL- Jeboori TA. Synthesis characterization and antimicrobial activities of {Fe(II), Co(II), Ni(II), Cu(II), and Zn(II) mixed ligand complexes Schiff base derived from ampicillin drug and 4(dimethylamino)benzaldehyde with nicotinamide. *Int J Technical Research and Applications*, 2014;2(4):187-192.
34. Kaushal R, Thakur S. Syntheses and Biological screening of Schiff base complexes of Titanium (IV). *Chem Engi Trans*, 2013;32:1801-1806.
35. Arun TN, Gowramma B. Synthesis and biological evaluation of some novel Schiff bases of cephalixin. *IJPSR*, 2014;5:1008-1014.
36. Al-Noor HT , AL- Jeboori AT, Aziz RM. Preparation, Characterization and Antimicrobial activities of {Fe(II),Co(II),Ni(II),Cu(II),and Zn(II)}Mixed Ligand Complexes Schiff base derived from Cephalixin drug and 4(dimethylamino)benzaldehyde with Nicotinamide. *Adv Phy Theo Appl*, 2013;18:1-8.

37. Elgellal AA, Preparation Characterization and Biological Evaluation of Schiff-Base of Some Drug Substances, *J Med Bioeng*, 2014;3:187-189.
38. Iqbal SM, Bukhari HI, Arif M. Preparation, characterization and biological evaluation of copper(II) and zinc(II) complexes with Schiff bases derived from amoxicillin and cephalexin, *Appl Organometal Chem* 2005;19:864-869.
39. Srivastava KP, Singh A, Singh KS. Microwave Assisted Synthesis, Characterisation and Antibacterial Study of Drug based Schiff Bases and their Zn (II) Complexes. *Am Int J Res Sci Technol Eng Math*, 2014;6:286-292.
40. Narjis N, Zubi S, Zafar MI, synthesis spectral and biological studies of novel Schiff base metal complexes derived from cephalexin and sugars. *Asian J Chem*, 2013;25:297-300.
41. Al-Noor HT, Jarad JA, Hussien OA. Synthesis physico- chemical and antimicrobial properties of some metal (II) Mixed ligand complexes of tridentate Schiff base derives from B- lactam antibiotic {(cephalexin mono hydrate) -4-chlorobenzophenone} and saccharin. *Int J Chem Pro Engi Res*, 2014;1:109-120.
42. Anacona J, Pineda Y, Alina Bravo A, et al. Synthesis, Characterization and Antibacterial Activity of a Tridentate Schiff Base Derived from Cephalexin and 1,6-Hexanediamine and its Transition Metal Complexes. *Med Chem*, 2016;6:467-473.
43. Joshi S, Pawar V , Uma V. Antibacterial and Antioxidant Properties of Mn (II), Co (II), Ni (II) and Zn (II) Complex of Schiff base derived from Cephalexin. *RJPBCS*, 2011;2:61- 70.
44. Srivastava KP, Singh A, Singh KS, et al. Green and Efficient Synthesis, Characterization and Anti- Bacterial Activity of Copper (II) Complexes with Unsymmetrical Bidentate Schiff Base Ligands. *J Appl Chem*, 2014;7:16-23.
45. Chohan HZ, Pervez H, Khan MK, et al. Antifungal Cobalt(II), Copper(II), Nickel(II) and Zinc(II) Complexes of Furanyl-,Thiophenyl-, Pyrrolyl-, Salicylyl- and Pyridyl-derived Cephalexins, *J Enzy Inhibition Med Chem*, 2004;19:85–90.
46. Iqbal SM, Abid R, Sabir MA, et al. Preparation, Characterization and Biological Evaluation of Copper(II) and Zinc(II) Complexes with Cephalexin, *J Pharm Pharmacol* 1999;51:371-375.
47. Srivastava KP, Singh A, Singh KS. Eco-friendly and Efficient Synthesis, Characterisation and Antibacterial Studies of Unsymmetrical Bidentate Schiff Bases and their Zn (II) Complexes. *Orient J Chem*, 2014;30:1233-1241.
48. Deepak A, Chauhan S, Verma KK, et al. Preparation, Characterization and Biological Studies on Pyridoxylidene-Cephalexin Schiff Base Complexes of Aryltellurium (IV). In *J Chem Sci*, 2017;15:1-12.
49. Anacona JR, Rodriguez JL, Camus J, Synthesis, characterization and antibacterial activity of a Schiff base derived from cephalexin and sulphathiazole and its transition metal complexes, *Spectrochimica Acta*, 2014;129:96-102.
50. Anacona JR, Rangel V, Lorono M, et al. Tetradentate metal complexes derived from cephalexin and 2,6-diacetylpyridine bis(hydrazone):Synthesis, characterization and antibacterial activity, *Spectrochimica Acta Part A:Molecular and Biomolecular Spectroscopy*, 2015;149:23-29.
51. Al-Bayati RI, Al-Akidi SM, Al-Azzawi IK. Synthesis of new cephalosporin derivatives Al- Mustansiriya *J Sci*. 2009;20:8-18.

52. Alwan SM. Synthesis and preliminary antimicrobial activities of new arylideneamino-1,3,4-thiadiazole-(thio/dithio)-acetamido cephalosporanic acids. *Molecules* 2012;1025-1038.
53. Abdulghani AJ, Mohuee SK. Synthesis and Characterization of New Schiff Bases Derived from Ceftriaxone Sodium with 1H- Indole-2, 3- Dione (Isatin) and 1-Acetyl Indoline-2, 3-Dione (N-Acetylisatin) and their Platinum (IV) Complexes. *J Chem Bio Phy Sci.* 2016;6:579-595.