

## Development and validation of a RP-HPLC method for simultaneous determination of ritonavir and lopinavir in combined dosage form

Raja Abhilash Punagoti<sup>1\*</sup>, Venkateshwar Rao Jupally<sup>2</sup>

<sup>1</sup>S.V.S.Group of Institutions, School of Pharmacy, Bhemaram, hanamkonda. Warangal,A.P (INDIA)

<sup>2</sup>TallaPadmavathi College of Pharmacy, Urus, Warangal, A.P (INDIA)

<sup>3</sup>Department of pharmacy, Acharyanagarjuna university, Guntur, A.P (INDIA)

E-mail : abhilashmpharm@gmail.com

### ABSTRACT

A reverse phase HPLC method was developed and validated for the simultaneous determination of two antiviral drugs viz. lopinavir and ritonavir. Chromatography was carried out on C18 Column, Inertsil (150 x 4.6 mm, 5  $\mu$ ) with mobile phase of Buffer: Methanol: Acetonitrile (30:60:10v/v/v) pH 6.5, at a flow rate of 0.6 ml/min and detection was made at 226 nm. The different analytical performance parameters such as linearity, precision, accuracy, specificity, limit of detection (LOD), limit of quantification (LOQ) and robustness were determined according to ICH guidelines. The linearity for each analyte in the desired concentration range was good ( $r^2 > 0.9$ ). The recovery of the method was 99.86% and 100.16% for ritonavir and lopinavir respectively. Hence the proposed method was sensitive, precise and accurate and it can be successfully apply to estimate the amount of ritonavir and lopinavir in the formulations by easily available low cost materials.

© 2014 Trade Science Inc. - INDIA

### KEYWORDS

RP-HPLC;  
Ritonavir;  
Lopinavir;  
Method development;  
validation.

### INTRODUCTION

In recent years, the treatment of human immunodeficiency virus (HIV)-1 infection and AIDS has been advanced by the development of highly active antiretroviral therapy.

Lopinavir inhibits the HIV viral protease enzyme. This prevents cleavage of the gagpol poly protein and, therefore, improper viral assembly results. This subsequently results in non-infectious, immature viral particles. Combination therapy with the HIV protease inhibitors lopinavir and ritonavir has been shown to be effective against drug-resistant HIV-13. These agents are me-

tabolized by cytochrome P-450 (CYP) 3A in the liver. When lopinavir is administered with ritonavir, ritonavir inhibits the CYP 3A- mediated metabolism of lopinavir, thereby providing increased plasma levels of lopinavir<sup>[1]</sup>.

Lopinavir is chemically designated as [1S-[1R\*, (R\*), 3R\*, 4R\*]]-N-[4-[[[(2,6-dimethylphenoxy) acetyl] amino]-3-hydroxy-5-phenyl-1-(phenylmethyl) pentyl] tetrahydro- $\alpha$ -(1-methylethyl)-2-oxo-1(2H)-pyrimidineacetamide. Its molecular formula is  $C_{37}H_{48}N_4O_5$ , and its molecular weight is 628.801<sup>[1]</sup>.

Ritonavir is chemically designated as 10-Hydroxy-2-methyl-5-(1-methylethyl)-4-thiazolyl]-3,6-dioxo-8,11-bis(phenylmethyl)-2,4,7,12-tetraazatridecan-13-

oic acid, 5-thiazolylmethyl ester, [5S-(5R\*,8R\*,10R\*,11R\*)]. Its molecular formula is  $C_{37}H_{48}N_6O_5S_2$  and its molecular weight<sup>[1,2]</sup> is 720.95. The structures of Lopinavir and Ritonavir are depicted in figure 1.




Figure 1 : Structures of Lopinavir and Ritonavir

To analyze these drugs, several determination methods have been reported. These included high performance liquid chromatography- mass spectrometry (LC/MS)<sup>[3-5]</sup>, matrix-assisted laser desorption/ionization, time-of-flight (MALDI-TOF)<sup>[6]</sup>, high-pressure thin layer chromatography (HPTLC)<sup>[7]</sup>, Ultra performance liquid chromatography (UPLC)<sup>[8]</sup> and immunoassay methods<sup>[9]</sup>.

Literature also revealed some of the methods by high-performance liquid chromatography (HPLC)<sup>[10-14]</sup> in various dosage forms.

The present paper deals with a HPLC assay of lopinavir and ritonavir simultaneously. Benefits of this method are improved sensitivity and recovery compared to all other methods available in the literature. The present HPLC method is well suited for studies generating large number of samples. In addition, much improved sensitivity was achieved with lower limit of quantification (LLOQ) of 0.075  $\mu\text{g/ml}$  for lopinavir and 0.3  $\mu\text{g/ml}$  for ritonavir. Recoveries were about 99.86% and 100.16% for ritonavir and lopinavir respectively. The proposed method was subjected to validation accord-

ing to the ICH guidelines<sup>[15]</sup>.

## MATERIALS AND METHODS

### Materials and reagents

Ritonavir and Lopinavir were kindly supplied as a gift sample by Hetero drug pvt Ltd. AP. other reagents such as Acetonitrile, Methanol, Potassium dihydrogen Phosphate and water used were of HPLC and milli-q grade. All other chemicals like glacial acetic acid used were of AR grade from Merck chemicals, Mumbai.

### Chromatography conditions

chromatography separation was performed on a HPLC (Younglin with UV detector) at the wavelength of 226 nm. A reverse phase Inertsil C18, (150 X 4.6mm, i.d., 5  $\mu\text{m}$  particle size) column was used. The mobile phase consists of Buffer: Methanol: Acetonitrile (30:60:10v/v/v) with flow rate 1.2 ml/min. injection volume was 20 $\mu\text{l}$  and the chromatographic runtime of 15 min was used.

### Preparation of buffer solution

Buffer was prepared by dissolving 2.7218g of Potassium dihydrogen orthophosphate in 1000 mL of water and adjusts the pH  $6.5 \pm 0.02$  with Ortho Phosphoric acid followed by the degassing of the solution.

### Preparation of mobile phase

1000 mL of mobilephase was prepared by mixing 300ml of Potassium dihydrogen orthophosphate (0.02M), 250ml of Methanol and 450ml of Acetonitrile.

### Preparation stock solutions

Accurately weighed and transferred about 10mg of Ritanovir and Lopinavir into a separate 10 mL volumetric flasks. 5 mL of Acetonitrile was added to each volumetric flask and kept in an ultrasonic bath until it dissolved completely. Made the volume up to the mark with methanol and mixed well. This yielded solution of 1000 $\mu\text{g/ml}$  concentration.

### Preparation of sandard solution

Spiked accurately about 0.4 mL of Ritanovir and 1.6 mL of Lopinavir stock solution and transfered it into a 10ml volumetric flask. Made the volume up to

## Full Paper

the mark with mobile phase and mixed well. This yielded solution of 40 $\mu$ g/ml and 160 $\mu$ g/ml concentrations.

### Method validation

Validation experiments were performed to demonstrate System suitability, precision, linearity, Accuracy, Limit of detection and Limit of quantification.

### Precision

The precision of the method was evaluated by carrying out six independent assays of test sample against a qualified reference standard and the %RSD of assay was calculated (% RSD should not be more than 2%).

### Accuracy

Accuracy for the assay of Ritonavir and Lopinavir determined by applying the method in triplicate samples to which known amount of Ritonavir and Lopinavir standard is added at different levels (50%, 100%, and 150%). Each solution was injected thrice (n=3) into HPLC system and the average peak area was calculated from which Percentage recoveries were calculated. (% Recovery should be between 98.0 to 102.0%).

### Linearity

The Linearity of detector response was established by plotting a graph to concentration versus area of Ritonavir and Lopinavir standard and determining the correlation coefficient. A series of solution of Ritonavir and Lopinavir standard solution in the concentration ranging from about 5 - 60 $\mu$ g/ml of Ritonavir and 20 - 240 $\mu$ g/ml of Lopinavir respective levels of the target concentration were prepared and injected into the HPLC system. (Correlation coefficient should be not less than 0.999.)

### Limit of Detection (LOD) Limit of Quantification (LOQ)

LOD and LOQ for the were determined at signal to noise ratios of 3:1 and 10:1, respectively by injecting series of dilute solutions with known concentrations.

## RESULTS AND DISCUSSION

### Method development

Different chromatographic conditions were experimented to achieve better efficiency of the chromato-

graphic system. Parameters such as mobile phase composition, wavelength of detection, column, column temperature, and pH of mobile phase were optimized. Several proportions of buffer, and solvents (water, methanol and acetonitrile) were evaluated in order to obtain suitable composition of the mobile phase. Choice of retention time, tailing, theoretical plates, and run time were the major tasks while developing the method. Buffers like sodium dihydrogen orthophosphate and disodium hydrogen orthophosphate did not yield desired results. The composition of mobile phase Acetonitrile: Methanol: Buffer (50:30:20v/v/v) of pH 3.5 with flow rate of 0.8ml/min and detection at 210nm of runtime of 6min shown merging of two drugs. The composition of mobile phase Acetonitrile: Methanol: Buffer (60:30:20v/v/v) of pH 3.5 with flow rate of 1ml/min and detection at 226nm of runtime of 15 min yielded peaks with non-sink in the base line with unstable retention times.

At Acetonitrile: Methanol: Buffer (45:25:30v/v/v) of pH 6.5 with flow rate of 1.2ml/min and detection at 226nm of runtime of 12min, a perfect chromatogram was eluted. The typical chromatogram obtained from final HPLC conditions are depicted in Figure 2.




Figure 2 : typical chromatogram of Ritonavir and Lopinavir by proposed method

### Method validation

Based on International Conference on Harmonization (ICH) guidelines, the method is validated with regard to system suitability, linearity, accuracy, precision, LOD and LOQ as follows.

#### (I) system suitability

The system suitability parameter tailing factor for the proposed HPLC method from the standard injection of Ritonavir and Lopinavir are 1.03 and 1.02 re-

spectively. Theoretical Plates Obtained from the standard injection of Ritonavir and Lopinavir are 5350.4 and 6140.8 respectively. The results proved that the optimized HPLC method fulfils these requirements within the USP accepted limits.

## (II) Precision

The % R.S.D. of Ritonavir and Lopinavir assay during the method precision was found to be 0.052948% and 0.089858% respectively, indicating excellent precision of the method. The results are summarized in TABLE 1.

TABLE 1 : Results of precision

Injection number	Ritonavir (40µg/ml) Peak Area	Lopinavir (160µg/ml) Peak Area
1	416238	172271
2	415725	172418
3	415852	171989
4	415715	172184
5	415625	172103
6	415936	172319
MEAN	415848.5	172214
% RSD	0.052948	0.089858

TABLE 2 : Summary results of accuracy for ritonavir and lopinavir

RITONAVIR							
Recovery Level	Resultant solution (µg/ml)	Standard Injections				% Recovery	%RSD
		inj-1	inj-2	inj-3	inj-4		
50%	20	205254	206363	205263	205626.7	98.68485	0.310125
100%	40	415852	416365	416536	416251	99.8841	0.085517
150%	60	633524	629524	631334	631460.7	101.0174	0.317202
LOPINAVIR							
50%	80	85289	84767	85134	85063.33	98.91545	0.315152
100%	160	172045	171742	172157	171981.3	99.9938	0.124839
150%	240	259674	263745	262914	262111	101.5981	0.820651

TABLE 3 : Results of Robustness

RITONAVIR			
Chromatography changes	Area	Retention Time	%RSD
Flow rate (ml/min)			
0.6	496786	9.08	0.32
0.7*	416077	8.26	0.28
0.8	345577	7.46	0.42
Change in organic composition in the mobile phase (Buffer: Methanol: Acetonitrile)			
25:62.5:12.5	387258	7.96	0.50
(30:60:10)	416077	8.26	0.28
35:57.5:7.5	466750	8.65	0.52
LOPINAVIR			
Flow rate (ml/min)			
0.6	248443	11.11	0.48
0.7*	174020	9.97	1.08
0.8	104625	8.85	0.66
Change in organic composition in the mobile phase (Buffer: Methanol: Acetonitrile)			
25:62.5:12.5	137038	9.57	0.46
(30:60:10)	174020	9.97	1.08
35:57.5:7.5	204960	10.41	0.42

## (III) Accuracy

Percent recovery of Ritonavir samples ranged from 98.0% to 101.5%, and the Percent recovery of Lopinavir samples ranged from 98.0% to 102.0% showing the good accuracy of the method. The result is shown in TABLE 2.

## (IV) Limit of Detection (LOD) Limit of Quantification (LOQ)

The LOD of Ritonavir and Lopinavir were found to be 0.02µg/ml and 2µg/ml respectively. The LOQ was 0.05µg/ml and 1 µg/ml for Ritonavir and Lopinavir respectively. Since the LOQ and LOD values of Ritonavir and Lopinavir achieved at a very low level, this method can be suitable for cleaning validation in the pharmaceutical industry.

## (V) Linearity

The linearity of the calibration plot for the method was obtained over the calibration ranges tested, i.e., 5-60 µg/ml for Ritonavir and 20-240 µg/ml for Lopinavir three times, and the correlation coefficient obtained was

## Full Paper

0.9998 and 1 for Ritonavir and Lopinavir respectively, thus indicating excellent correlation between peak areas and concentrations of the analytes.

### (VI) Robustness

In all the deliberately varied chromatographic conditions in the concentration range for the evaluation of robustness 5- 60 µg/ml for Ritonavir and 20-240 µg/ml for Lopinavir (n=3). It can be concluded that the variation in flow rate and the variation in 10% Organic composition do not affect the method significantly. Hence it indicates that the method is robust even by change in the flow rate ±10% and change in the Mobile phase ±10%. The results are summarized in TABLE 3.

### CONCLUSION

The new, simultaneous RP-HPLC method proved to be simple, linear, precise, accurate, robust, rugged and rapid. The developed method was capable of giving faster elution, maintaining good separation more than that achieved with other available HPLC methods. The short retention times allows the analysis of a large number of samples in a short period of time and is therefore more cost-effective for routine analysis in the pharmaceutical industries. It is suitable for rapid and accurate quality control of ritonavir and lopinavir in combined dosage forms.

### REFERENCES

- [1] Product information; Kaletra Lopinavir/Ritonavir and Abbott Park, IL, Abbott Laboratories, (2000).
- [2] A.H.Corbett, M.L.Lim, A.D.Kashuba; Kaletra (lopinavir/ritonavir). *Ann Pharmacother.*, **36**, 1193-1203 (2002).
- [3] M.Ehrhardt, M.Möck, W.E.Haefeli; *J Chromatogr B.*, **850**, 249-258 (2007).
- [4] A.D.Avolio, M.Siccardi, M.Sciandra; *J Chromatogr B.*, **859**, 234-240 (2007).
- [5] R.T.Heine, H.Rosing, E.C.M.Van Gorp; *J Chromatogr B.*, **867**, 205-212 (2008).
- [6] S Notari, C Mancone and M.Tripodi; *J Chromatogr B.*, **833**, 109-116 (2006).
- [7] A.V.Sulebhavikar, U.D.Pawar, K.V.Mangoankar; *E-J Chem.*, **5(4)**, 706-712 (2008).
- [8] V.A.Marina, P.Julia, P.R.Jorge, S.Eduardo, L.Rafael.; *J Braz Chem Soc.*, **22**, 134-141 (2011).
- [9] S.Azoulay, M.C.Nevers, C.Creminon; *J Immunol Methods*, **295**, 37-48 (2004).
- [10] C.Marzolini, A.Beguin, A.Telenti, A.Schreyer, T.Buclin, J.biollaz, L.A.decosterd; *J Chromatogr B.*, **774**, 127-140 (2002).
- [11] Yoshico Usami, Tsuyoshi Oki; *Chem Pharm Bull.*, **51(6)**, 715-718 (2003).
- [12] J.Ray, E.Pang, D.Carey; *J Chromatogr B.*, **775**, 225-230 (2002).
- [13] C.M.Phechkrajang, E.E.Thin, L.Sratthaphut, D.Nacapricha, P.Wilairat; *Mahi Univ J Pharm Sci.*, **36**, 1-12 (2009).
- [14] E.M.Donato, C.L.Dias, R.C.Rossi, R.S.Valente, P.E.Froelich; *J Chromatogr B.*, **63**, 437-443 (2006).
- [15] Anonymous; ICH Guidelines. Validation of Analytical Procedures: Text and Methodology Q2 (R1), (1996).