



# **DEVELOPMENT OF METHOD FOR EXTRACTIVE SPECTROPHOTOMETRIC DETERMINATION OF Co (II) WITH 2-HYDROXY-1-NAPHTHALENE CARBOXALDEHYDE PHENYL HYDRAZONE AS AN ANALYTICAL REAGENT**

**RATNAMALA P. SONAWANE<sup>a</sup>, R. S. LOKHANDE and  
UTKARSHA M. CHAVAN**

The Institute of Science, Fort, MUMBAI – 60 (M.S.) INDIA

## **ABSTRACT**

A spectrophotometric method has been developed for the determination of Co (II) using 2-hydroxy-1-naphthalene carboxaldehyde phenyl hydrazone as an extractive reagent. The reagent forms a coloured complex, which has been quantitatively extracted into n-butanol at pH 8.4. The method obeys Beer's law over arrange from 1 to 10 ppm. The Molar absorptivity and Sandell's sensitivity were  $2.380 \times 10^4 \text{ L mol}^{-1}\text{cm}^{-1}$  and  $0.1176 \mu\text{g cm}^{-2}$ , respectively. The proposed method is very sensitive and selective. The method has been successfully applied to synthetic and commercial samples.

**Key words:** Cobalt spectrophotometric determination, n-Butanol, 2-Hydroxy-1-Naphthalene carboxaldehyde phenyl hydrazone.

## **INTRODUCTION**

A number of reagents such as hydrazone<sup>1</sup>, thiosemicarbazone<sup>2,3</sup>, oxime<sup>4,5</sup> etc. have been used for the determination of Co (II). Aromatic hydrazone derivative has also been used. The synthesis of hydrazone derivative and their applications towards metal ions have been reported<sup>6,7</sup>. Cobalt compounds have been used for centuries to impart a rich blue colour to glass, glazes and ceramics. Cobalt in small amounts is essential to many living organisms, including humans. Although cobalt is an essential element for life in minute amounts. At higher levels of exposure it shows mutagenic and carcinogenic effects. Several compounds are known to react with the metal ions to give coloured complexes and have been employed for the quantitative extraction and spectrophotometric determination of metals at trace level. In continuation to our earlier work<sup>8,9</sup>, the proposed method is a simple, rapid and selective

---

\* Author for correspondence; E-mail: ratnasonawane@yahoo.in

towards spectrophotometric determination of Co (II) using 2-hydroxy-1-naphthalene carboxaldehyde phenyl hydrazone derivative.

## EXPERIMENTAL

The reagent 2-hydroxy-1-naphthalene carboxaldehyde phenyl hydrazone was synthesized by the given procedure. The stock solution of Co (II) was prepared by dissolving a weighed amount of sulphate in double distilled water containing dilute sulphuric acid, which was diluted to the desired volume with double distilled water and standardized by nitroso-R- salt method<sup>10</sup>. Absorbance and pH measurements were carried out on a Shimadzu UV-Visible 2100 spectrophotometer with 1 cm quartz cells and digital pH meter with combined glass electrode respectively.

### Procedure for the extraction

1.0 mL of aqueous solution containing 0.1 mg of cobalt metal and 1 mL of reagent were mixed in a 50 mL beaker. The pH of the solution adjusted to 8.4 with 0.2 M boric acid and potassium chloride, keeping the volume 10 mL. The solution was transferred to 100 mL separating funnel. The beaker was washed twice with n-butanol and transferred to the same funnel. The two phases were shaken for two minutes and allowed to separate. The organic phase was collected in 10 mL measuring flask and made up to the mark with organic solvent if required. After separation of the two phases, the pH of the aqueous phase was measured and the Co (II) in each phase was determined by nitroso-R- salt method.<sup>10</sup>

## RESULTS AND DISCUSSION

The reagent HNPH forms yellow coloured complex with Co (II), which was extracted into organic phase. The extraction of Co (II) forms an aqueous phase by HNPH in n-Butanol is studied over a wide range of experimental condition. The results of various studies are discussed below.

### Extraction as a function of pH

The extraction of cobalt with 2-hydroxy-1-naphthalene carboxaldehyde phenyl hydrazone has been studied over the pH range 1-10 and was observed that percentage extraction of Co (II) is maximum at pH 8.4

### Absorption spectrum

The absorption spectrum of Co (II): 2-hydroxy-1-naphthalene carboxaldehyde phenyl hydrazone in n-butanol shows the maximum absorption at 410 nm. The absorption

due to reagent at this wavelength is nearly negligible. Hence the absorption measurements were carried out at 410 nm.

### **Influence of diluents**

The suitability of diluents was investigated using organic solvents such as chloroform, ethyl acetate, isoamyl alcohol, xylene, hexane, diethyl ether, toluene, n-butanol, carbon tetrachloride, MIBK, nitrobenzene, etc. The extraction of cobalt (II) was quantitative with HNPH in n-butanol. Hence, n-butanol was used for further extraction studies, as it gave better and quicker phase separation.

### **Effect of salting out agents**

The presence of 0.1 M salts of various alkali and alkaline metals does not show any effect over the absorbance value of Co (II): 1-naphthalene carboxaldehyde-2-hydroxy phenyl hydrazone complex extract. Therefore, no salting out agent was required during the extraction.

### **Effect of reagent concentration**


Various volumes of 0.1% reagent solution were added to the sample solution containing 80 µg of cobalt at respective pH values. The absorbance remained nearly constant when the volume of the reagent solution used was more than 1 mL. Therefore, 1 mL of 0.1% reagent was chosen for the quantitative determination of the metal.

### **Effect of equilibration time and stability of the complex**

The study of change in absorbance with variation in equilibrium time for extraction of extraction of the complex into organic solvent shows that equilibration time of 1 min is sufficient for the quantitative extraction of cobalt. The study of stability of colour of the Co (II): HNPH complex with respect to time shows that the absorbance due to extracted species is stable up to 48 hours, after which slight decrease in absorbance is observed. Throughout the experimental work, for practical convenience, the measurements have been carried out within one hour of extraction of cobalt.

### **Calibration plot**


A calibration plot of absorbance against varying cobalt concentration and fixed HNPH concentration gives linear and reproducible graph in the concentration range 1 to 10 ppm of cobalt (Fig. 1). This shows that the Beer's law is obeyed in this range. The molar absorptivity and Sandell's sensitivity were calculated to be is  $2.380 \times 10^4 \text{ L mol}^{-1}\text{cm}^{-1}$  and  $0.1176 \text{ µg cm}^{-2}$ , respectively.



**Fig. 1: Calibration plot of Co (II): HNPH complex**

### Nature of extracted species

The composition of extracted species has been determined by Job's continuous variation method, Slope ratio (Fig. 2) method and Mole ratio method. It shows that the composition of Co (II): HNPH complex is 1 : 2.



**Fig. 2: Slope ratio method for Co (II): HNPH complex**

### Effect of divalent ions and foreign ions

The effect of other ions present in various amount indicated no interference in the spectrophotometric determination of 60  $\mu\text{g}$  of cobalt. The ions which show interference in the spectrophotometric determination of cobalt were overcome by using appropriate masking agents (Table 2).

### Precision and accuracy

The precision and accuracy of the developed spectrophotometric method have been studied by analyzing five solutions each containing 40  $\mu\text{g}$  of cobalt in aqueous phase. The average of five determinations was 40.03, variance 0.006 and variation from mean at 99% confidence limit was  $\pm 0.147$ .

### Applications

Various commercial samples and synthetic mixtures containing Co (II) were prepared and analyzed according to the recommended procedure and the results were compared to those obtained by standard method. The proposed method facilitates separation of cobalt (II) from synthetic mixtures. These metal ions do not extract and remain quantitatively in the aqueous phase under the optimum extraction conditions of cobalt (II) with HNPH system facilitating separation of bivalent cobalt quantitatively by the proposed method. The results found to be in good agreement with those obtained by the standard known method<sup>10</sup> (Table 1).

**Table 1: Determination of Co (II) using HNPH from different samples**

S. No.	Sample	Amount of Co (II)	
		Standard method	Present method
<b>I</b>	Cobalt alloys		
<b>1</b>	Cobalamine	69.9%	69.8%
<b>2</b>	Steel	9.56%	9.54%
	Synthetic mixture		
<b>1</b>	Co (10) + Zn (10)	9.98 ppm	9.98 ppm
<b>2</b>	Co (10) + Mo (10)	9.98 ppm	9.96 ppm
<b>3</b>	Co (10) + Mg (10)	9.98 ppm	9.97 ppm

**Table 2: Effect of divalent ions and foreign ions**

<b>Ion</b>	<b>Tolerated Ratio</b>	<b>Ion</b>	<b>Tolerated Ratio</b>
Cl <sup>-</sup>	1 : 15	Zn <sup>2+</sup>	1 : 16
Br <sup>-</sup>	1 : 10	Ag <sup>+</sup>	1 : 12
F <sup>-</sup>	1 : 11	K <sup>+</sup>	1 : 20
ClO <sub>3</sub> <sup>-</sup>	1 : 8	Mg <sup>2+</sup>	1 : 15
BrO <sub>3</sub> <sup>-</sup>	1 : 11	Ca <sup>2+</sup>	1 : 12
IO <sub>3</sub> <sup>-</sup>	1 : 16	Ba <sup>2+</sup>	1 : 12
SO <sub>3</sub> <sup>2-</sup>	1 : 7	Bi <sup>2+</sup>	1 : 9
SO <sub>4</sub> <sup>2-</sup>	1 : 15	V <sup>+5</sup>	1 : 10
NO <sub>2</sub> <sup>-</sup>	1 : 13	Cr <sup>3+</sup>	Masked
NO <sub>3</sub> <sup>-</sup>	1 : 10	Mn <sup>2+</sup>	Masked
PO <sub>4</sub> <sup>3-</sup>	1 : 20	Ce <sup>4+</sup>	Masked
P <sub>2</sub> O <sub>7</sub> <sup>2-</sup>	1 : 19	CN <sup>-</sup>	Masked
ClO <sub>4</sub> <sup>-</sup>	1 : 15	Tartarate	Masked

## CONCLUSION

The results obtained show that HNPH in n-butanol can be effectively used for quantitative extraction from aqueous media. The proposed method is found to be quantitative as compared to other standard methods. The equilibrium time required is very little, i.e. only 1 min and the complex is stable for 48 hrs. The results show good agreement with the standard method. The method is very fast, accurate and precise. The 2-hydroxy-1-naphthalene carboxaldehyde phenyl hydrazone was used for the first time for extraction of Co (II) from various binary mixtures.

## REFERENCES

1. P. V. Jagasia and D. P. Dave, J. Ind. Chem. Soc., **80**, 145 (2003).
2. S. L. Subramanyan, K. J. Rajesh, C. J. and A. Varada, Analytical Letters, **36**, 605 (2003).

3. R. S. Lokhande, N. Shrinivasan and A. B. Chaudhari, *J. Ind. Council of Chem.*, **19**, 12 (2002).
4. H. Eskandari, H. S. Ghaziaskar and A. A. Ensaf, *Anal Sci.*, **17**, 327 (2001).
5. J. J. Desai, P. G. Desai and A. G. Mehta, *Asian J. Chem.*, **12**, 1067 (2000).
6. Byron L. Hawbecker, *J. Chem. Edu.*, **47**, 218 (1970).
7. D. E. Pearson and C. M. Greer, *J. Am. Chem. Soc.*, **71**, 1895 (1949).
8. R. S. Lokhande and S. V. Poman, *Asian J. Chem.* **13**, 1222 (2001).
9. R. S. Lokhande and A. S. Jayant, *Asian J. Chem.*, **11**, 1040 (1999).
10. G. H. Jeffery, J. Basset and R. C. Denney, *Vogel's Textbook of Quantitative Chemical Analysis*, 5<sup>th</sup> Ed<sup>n</sup>. ELBS, London (1991).

*Accepted : 07.02.2013*