



Trade Science Inc.

# Environmental Science

An Indian Journal

Ecotoxicology

ESAIJ, 4(5), 2009 [205-207]

## Effect of fresh, composted and vermicomposted parthenium and poultry droppings on yield and quality of Radish (*Raphanus sativus* var. pusa chetki)

R.Hiranmai Yadav<sup>1\*</sup>, B.Vijayakumari<sup>2</sup><sup>1</sup>Department of Botany, Avinahilingam University, Coimbatore-641043, Tamilnadu, (INDIA)<sup>2</sup>Department of Biotechnology, Karpagam University, Coimbatore-641021, Tamilnadu, (INDIA)

Tel : +091-422-2611146; Fax : +091-422-2611043

E-mail : mayahiran@yahoo.com

Received: 5<sup>th</sup> May, 2009 ; Accepted: 10<sup>th</sup> May, 2009

### ABSTRACT

A pot culture experiment was conducted to assess the effect of fresh, composted, and vermicomposted *parthenium* and poultry droppings on the yield and quality of radish. There were fifteen treatments in different dosages each with three replications. Among the treatments radish beetle length was more in T<sub>2</sub> on 45 DAS. The beetle diameter was increased due to T<sub>5</sub> on 45 and T<sub>1</sub> on 60 DAS. The beetle weight of radish was maximum in T<sub>4</sub> on 60 DAS. In radish beetles increased ascorbic acid content was in T<sub>8</sub> and T<sub>4</sub> radish had higher protein content. © 2009 Trade Science Inc. - INDIA

### KEYWORDS

Radish;  
Compost;  
Vermicompost;  
Parthenium;  
Poultry droppings;  
Yield;  
Quality attributes.

### INTRODUCTION

With the Green Revolution and its positive contribution to the nation's self sufficiency fast on the ebb, the 21st century India is in the need of another revolution in its farming sector. While a few argue that biotechnology is the panacea, a few others vociferously prescribe organic farming in the context of a second Green Revolution<sup>[1]</sup>.

The increasing cost of fertilizer, growing environmental concern and energy crisis have created considerable interest for search of alternative cheap sources of plant nutrients. The use of vermicompost, as a source of organic manure in supplementing chemical fertilizer is becoming popular among farmers of the country. Vermicompost is sustainable supplementary organic manure regenerated from organic waste using earthworms. Vermicompost is rich in nutrients and organic matter

content. It can be used by the farmers, horticulturists, floriculturists and garden lovers.

Composting is a biological process in which microorganisms, mainly fungi and bacteria, convert degradable organic waste into humus like substance. This finished product, which looks like soil, is high in carbon and nitrogen and is an excellent medium for growing plants. The process of composting ensures the waste that is produced in the kitchens is not carelessly thrown and left to rot. It recycles the nutrients and returns them to the soil as nutrients.

Though soil is the most vital input for food production, it is generally in a degraded condition due to many reasons and indiscriminate use of chemical fertilizers is the prime reason among them. Since improving the soil health by adding organic manure is a priority. High quality organic manure can be prepared by the agriculturists themselves in their farms by establishing vermicompost

## Ecotoxicology

unit using biodegradable farm waste available in their farms such as dried leaves, vegetable and fruit waste and crop remains, which is burned otherwise<sup>[2]</sup>.

The objective of the present study was to determine the effect of fresh, composted and vermicomposted parthenium and poultry droppings in improving the yield and quality of radish.

Radish is one of the most ancient and widely cultivated vegetable of temperate region in almost all altitudinal zones. This vegetable being a good source of vitamin C and minerals is grown for its tender roots as a cooking vegetable.

### MATERIALS AND METHODS

A pot culture experiment was conducted using 7kg capacity pots to study the effects of fresh, composted and vermicomposted parthenium and poultry droppings on yield and quality of radish. The radish was grown in completely randomized design with three replications. The dosages were as per the recommendations of Tamil Nadu Agricultural University, Coimbatore.

#### The treatment details

- T<sub>0</sub> - Control – Red loamy soil (7 kg)
- T<sub>1</sub> - Fresh *Parthenium* @ 35 g per pot
- T<sub>2</sub> - Composted *Parthenium* @ 26.25 g per pot
- T<sub>3</sub> - Composted *Parthenium* @ 35.0 g per pot
- T<sub>4</sub> - Composted *Parthenium* @ 43.75 g per pot
- T<sub>5</sub> - Vermicomposted *Parthenium* @ 26.25 g per pot
- T<sub>6</sub> - Vermicomposted *Parthenium* @ 35.0 g per pot
- T<sub>7</sub> - Vermicomposted *Parthenium* @ 43.75 g per pot.
- T<sub>8</sub> - Poultry droppings @ 35.0 g per pot
- T<sub>9</sub> - Composted poultry droppings @ 26.25 g per pot
- T<sub>10</sub> - Composted poultry droppings @ 35.0 g per pot
- T<sub>11</sub> - Composted poultry droppings @ 43.75 g per pot
- T<sub>12</sub> - Vermicomposted poultry droppings @ 26.25 g per pot
- T<sub>13</sub> - Vermicomposted poultry droppings @ 35.0 g per pot
- T<sub>14</sub> - Vermicomposted poultry droppings @ 43.75 g per pot
- T<sub>15</sub> - N: P: K (60:30:30 kg/ha)

#### Yield components

Bettle length of a radish is measured with a scale and expressed in cm. The diameter of a bettle was measured using a thread and recorded in cm. The weight of a bettle was measured and expressed in g. The protein content was estimated following the method of Lowry et al.<sup>[3]</sup> and ascorbic acid by the method of Sadasivam and Theymoli<sup>[4]</sup>.

#### Statistical analysis

The data collected from the different studies were subjected to statistical analysis as per the procedure outlined by Panse and Sukhatme<sup>[5]</sup>.

### RESULTS AND DISCUSSION

The yield components of radish namely bettle length, bettle diameter and bettle weight were recorded on 45 and 60 DAS (TABLE 1). The protein and ascorbic acid in the bettle were tested on 60 DAS (TABLE 2).

The bettle length of radish (*Raphanus sativus*) on 45 DAS was significantly high in T<sub>2</sub> (11.13 cm) among *Parthenium* applications and T<sub>15</sub> (10.50 cm) compared to control (6.33 cm). On 45 DAS, among *Parthenium* and poultry droppings supplied plants, the bettles from T<sub>5</sub> and T<sub>14</sub> had increased diameter of 7.97 and 7.10 cm compared to control (4.47 cm). The T<sub>1</sub> and T<sub>12</sub> treatments recorded the highest values (11.90 and 11.30 cm) among *Parthenium* and poultry droppings applications on 60 DAS compared to the lowest value in control (8.63 cm). Bettle weight of radish (*Raphanus sativus*) was observed on 60 DAS and significantly higher values were found in T<sub>1</sub> (95.00 g) and T<sub>14</sub> (98.33

**TABLE 1: Impact of fresh, composted and vermicomposted *Parthenium* and poultry droppings on yield components of *Raphanus sativus***

Treatments	Bettle length (cm)		Bettle diameter (cm)		Bettle weight (g)
	45 DAS	60 DAS	45 DAS	60 DAS	60 DAS
T <sub>0</sub>	6.33	9.70	4.47	8.63	38.67
T <sub>1</sub>	10.27	13.27	7.27	11.90	95.00
T <sub>2</sub>	11.13	12.53	7.07	10.80	83.33
T <sub>3</sub>	9.97	10.80	7.57	11.10	70.67
T <sub>4</sub>	10.23	13.70	7.83	10.50	78.33
T <sub>5</sub>	9.80	12.80	7.97	11.43	74.33
T <sub>6</sub>	9.00	12.23	6.13	9.77	73.33
T <sub>7</sub>	8.00	11.50	7.50	10.97	70.00
T <sub>8</sub>	8.17	12.50	4.50	10.70	83.33
T <sub>9</sub>	8.27	12.53	4.87	9.00	56.67
T <sub>10</sub>	8.77	11.33	5.27	10.20	70.00
T <sub>11</sub>	8.40	11.50	5.90	10.63	66.67
T <sub>12</sub>	9.23	15.40	6.37	11.30	81.67
T <sub>13</sub>	10.27	14.33	6.53	10.97	86.67
T <sub>14</sub>	9.90	14.17	7.10	11.23	98.33
T <sub>15</sub>	10.50	13.33	4.63	10.77	66.67
SEd	0.98	NS	0.66	0.78	9.97
CD (5%)	2.00		1.34	1.58	20.30

SEd – Standard error deviation; CD – Critical difference; DAS – Days after sowing

**TABLE 2: Impact of fresh, composted and vermicomposted *Parthenium* and poultry droppings on yield components of *Raphanus sativus***

Treatments	Protein (60 DAS)	Ascorbic acid (60 DAS)
T <sub>0</sub>	2.12	285.33
T <sub>1</sub>	2.70	618.67
T <sub>2</sub>	2.42	432.00
T <sub>3</sub>	3.17	720.00
T <sub>4</sub>	4.47	576.00
T <sub>5</sub>	4.46	720.00
T <sub>6</sub>	2.42	576.00
T <sub>7</sub>	2.53	642.67
T <sub>8</sub>	2.41	1032.00
T <sub>9</sub>	2.33	541.33
T <sub>10</sub>	2.44	384.00
T <sub>11</sub>	2.49	456.00
T <sub>12</sub>	2.83	456.00
T <sub>13</sub>	2.49	480.00
T <sub>14</sub>	2.64	624.00
T <sub>15</sub>	2.72	360.00
SEd	0.18	90.54
CD (5%)	0.37**	184.44**

SEd-Standard error deviation; CD- Critical difference; DAS-Days after sowing

g) among fresh and decomposed *Parthenium* and poultry droppings treatments compared to control (38.67 g).

Protein contents were found to be higher in T<sub>4</sub> (4.47 mg g<sup>-1</sup>) and T<sub>5</sub> (4.46 mg g<sup>-1</sup>) among *Parthenium* treatments. T<sub>12</sub> (2.83 mg g<sup>-1</sup>) registered the highest value among poultry droppings applications. Protein content was very low in control (2.12 mg g<sup>-1</sup>).

Maximum ascorbic acid content of radish (*Raphanus sativus*) was noticed in T<sub>3</sub> and T<sub>5</sub> (720.00 mg g<sup>-1</sup>) among *Parthenium* applied crops. T<sub>8</sub> exhibited the highest value of 1032.00 mg g<sup>-1</sup> among poultry droppings treatments. The control recorded the lowest value of 285.33 mg g<sup>-1</sup>.

As in the present study Govindan et al.<sup>[6]</sup> also obtained maximum yield of *Abelmoschus esculentus* by vermicompost application. Increased grain yield of corn due to application of beef lot manure and its compost was reported by Eghball et al.<sup>[7]</sup>. Application of compost prepared from crop residue and poultry manure resulted in highest yield of maize<sup>[8]</sup>. The increase in yield may be due to the micro nutrients in organic source and bio active compounds formed during the composting of manures.

Treatment with biofertilizer +dense organic manure recorded the highest vitamin C content of radish, FYM + dense organic manure recorded the highest protein content of radish<sup>[9]</sup>.

## REFERENCES

- [1] The Hindu; Regain your Paradise the Brand Organic Way, (2008).
- [2] The Hindu; Setting up of Vermicompost Units in farms Emphasized, (2008).
- [3] O.H.Lowry, N.J.Rosenbrough, A.L.Farr and R.J.Randall; J.Biol.Chem., **193**, 267-275 (1951).
- [4] S.Sadasivam, B.Theymoli; Practical Manual in Biochemistry, TNAU, Coimbatore, 14 (1987).
- [5] V.G.Panse, P.V.Sukhatme; Statistical methods for agricultural workers, ICAR, New Delhi, 327 (1978).
- [6] M.Govindan, P.Muralidharan, S.Sasikumaran; J. Tropic.Agric., **33**, 173-174 (1995).
- [7] Eghball, B., D.Ginting and J.E. Gilley; Agron.J., **96**, 442-447 (2004).
- [8] J.A.Adediran, L.B.Taiwo, M.O.Akande, R.A. Sobulo, O.J.Idowu; J.Plant Nutr., **27**, 1163-1181 (2004).
- [9] S.R.Singh, A.K.Singh; Agric.Sci.Digest., **21**, 115-117 (2001).