



BioTechnology

An Indian Journal

FULL PAPER

BTAIJ, 8(2), 2013 [264-269]

Establishment of a RT-lamp method for the rabbit hemorrhagic disease virus detection

Yang Zexiao*, Zeng Hui, Wang Yin, Yao Xueping, Wang Kaiyu

College of Veterinary Medicine, Key Lab of Animal Disease and Human Health of Sichuan Province, Sichuan Agricultural University, Yaan Sichuan 625014, (CHINA)

E-mail: yzxyang2003@126.com; kywangsicau@126.com

ABSTRACT

To develop a new molecular biological method for the quick detection of rabbit hemorrhagic disease virus (RHDV), a set of 7 primers were designed according to the conserved sequence of the capsid protein VP60 gene of RHDV published in GenBank, and the reverse transcriptase loop-mediated isothermal amplification (RT-LAMP) assay was established through preparation of target gene fragments, optimization of reaction conditions, sensitivity and specificity tests. Results showed the RT-LAMP method for the RHDV detection had a ladder-like pattern of amplification bands from about 213 bp incubation at 64°C for 45min by using agarose gel electrophoresis, and with good sensitivity and specificity, the detection limit could reach about 5 copies of cloned viral genomic fragments, which was more sensitive than that of traditional RT-PCR, and no amplifications for gene fragment of European brown hare syndrome virus, *Pasteurella multocida*, *E.coli* and *Salmonella* from rabbits detection by this RT-LAMP approach. © 2013 Trade Science Inc. - INDIA

KEYWORDS

Rabbit hemorrhagic disease virus;
RT-LAMP;
VP60 gene;
Detection;
RT-PCR.

INSTRUCTION

Rabbit hemorrhagic disease (RHD), an important OIE Listed disease, is a highly contagious and rapidly fatal viral disease of both domestic and wild rabbits (*Oryctolagus cuniculus*)^[1]. It was caused by rabbit hemorrhagic disease virus (RHDV), which is a non-enveloped RNA virus with a diameter of about 36-40 nm, and the genome consists of a positive-sense single-stranded RNA molecule of 7437nt in size. The disease is characterized with the typical pathological lesions including necrotic hepatitis, hemorrhaging and edema of lungs and trachea, short incubation period (24-48 h af-

ter infection), and high mortality rate of 60-100%^[2]. Since RHD was first described in China in 1984, it has been reported in many countries successively throughout the world^[3], and has resulted in the deaths of nearly a quarter billion free-living and domestic rabbits and generated significant losses to rabbit farming industries and trade. RHD is still endemic in some areas sometimes, studies on the rapid diagnosis and prevention of RHD are still important and should be paid attention^[4,5]. Although there were many common RHD detection methods were reported such as virus isolation, ELISA^[6] and RT-PCR^[5], but these techniques might delay disease diagnosis for the limits of sensitiv-

ity, or requirements of laboratory facilities, etc.

As a novel nucleic acid amplification method, which was reported recently^[7], the RT-LAMP assay has been used for the detection of many pathogens because of advantages of rapidity, sensitivity, and non-requirements of any expensive special equipment such as a thermal cycler. In this paper, based on the sequence analysis and alignment of the capsid protein(VP60) gene of RHDV published in GenBank, 2 primers were designed for the preparation of target gene fragments by RT-PCR and a set of 5 primers were designed for the RT-LAMP, and a new molecular biological method for RHDV detection was described and assessed.

EXPERIMENTAL

Reagents and sample

TIANprep Mini Plasmid Kit(DP103)and 2×Taq PCR Master Mix were provided by TIANgen Biotech(Beijing) Co.,Ltd., Betaine (Z0041-1) was provided by Shanghai Kayon Biotechnology Co.,Ltd., DNA Maker DL 2000(D501A), Primerscript RT re-

agent Kit (BK1901), TaKaRa rTaq™ (DR001 AM), *Taq I* (D1051A) and RNAiso Plus were provided by TaKaRa Biotechnology (Dalian) Co.,Ltd., Bst polymerase (M0275) was provided by Beijing New England Biolabs Co.,Ltd., RHDV-Sch01, gene fragments of European brown hare syndrome virus (pGM-T-EBHSV), *Pasteurella multocida*, *E.coli* and *Salmonella* from rabbits were provided by the animal quarantine Lab of Sichuan Agricultural University; RHDV positive samples which collected from the experimentally infected rabbits with the RHDV-Sch01 were stored at -70 °C.

Primer preparation.

Based on the sequence analysis of RHDV VP60 gene published in GenBank(accession number: NC 001543), 2 primers(P1/P2)for RT-PCR and 5 primers for RT-LAMP(listed in table 1)were carefully designed using DNASTar and Primer Explorer V4 software (<http://primerexplorer.jp/e/>),respectively. All these primers were prepared by TaKaRa Biotechnology (Dalian) Co.,Ltd.

TABLE 1 : Primers used in this study

Primer name	Sequence (5'-3')	Products Size(bp)	Reference Sequence
RT- PCR			NC_001543
P1	GGCAATGACAACAGGTGGAA (754-773)	749	
P2	CTGAGCGAAAGCCCAATTGT (1502-1483)		
RT- LAMP			
RF3 :	GGTTTTCCACGTGCAACAG (827-809)	213	
RB3 :	AATGACATGTCAGGGAAGCC (1022-1003)		
RFIP :	GATGGTCAATGTCGGCAAACCGCTGGAACCTGAACGGCAG		
RBIP :	TGGGAACAACCTCCACCAACGTGCAACCTGGGAGATAGGGT		
RBLP :	TTTGGTACGCTAATGCTGGGT		

Preparation of target gene fragments

• Isolation of RHDV RNA and RT reaction:

RHDV RNA were extracted from the positive samples of experimentally infected rabbits with RHDV-Sch01 using the total RNA isolation reagent kit(RNAiso Plus reagent,Cat.No.BK1501, TaKaRa Biotechnology (Dalian) Co.,Ltd.,China) and stored at -70 °C. A total reaction volume of 10 µL, containing 2 µL of 5×Primerscript buffer, 0.5µL of Primerscript RT Enzyme Mix, 0.5µL of Oligo dT Primer(50 µmol/L),

0.5µL of Random 6 mers(100 µmol/L), 2.5µL of total RNA extracted above, 4.0µL of RNase-free ddH₂O, was used for reverse transcription reaction following the manufacturer's instructions of Primerscript RT reagent Kit(BK1901), and carried out at 37°C for 15 min, and 85°C for 5s, then the RT products (cDNA) were collected for further study.

• PCR reaction

PCR was carried out in a total volume of 50 µL, containing 3 µL of RT products (cDNA), 1.0 µL of

FULL PAPER

each primer (P1 and P2, 10 $\mu\text{mol/L}$), 25 μL of 2 \times Taq PCR Master Mix, and 20 μL of RNase-free ddH_2O , and performed at the following reaction conditions, denaturation at 95 $^\circ\text{C}$ for 5 min followed by 30 cycles at 94 $^\circ\text{C}$ for 40 s, a gradient of annealing degree from 55 $^\circ\text{C}$ for 45 s, and 72 $^\circ\text{C}$ for 60 s, and terminated by an extension at 72 $^\circ\text{C}$ for 8 min. 5 μL of PCR products were analyzed in 1.0% agarose in TAE buffer gel containing 0.5 mg/mL greenview using electrophoresed and photographed under a UV imaging system (Bio-Rad).

• Construction of recombinant plasmids:

The PCR products were separated and purified according to the instructions of TIANgel Midi Purification Kit (DP209), directly cloned into pMD-19T Vector to construct recombinant plasmid, and transformed into the DH5 α competent cells, then the recombinant plasmids were identified using agarose electrophoresis, PCR and sequencing^[8].

Basic RT-LAMP reaction.

The RT-LAMP reaction mixture (25 μL) contained 2.5 μL of 10 \times ThermoPol buffer, 1 μL (8 units) of BstDNA polymerase, 3 μL of MgCl_2 (25 mM), 3 μL of betaine (8 mM), 6 μL of dNTP (2.5 mM), 1 μL of each primers (RF3 (5 μM), RB3 (5 μM), RFIP (50 μM), RBIP (50 μM), and RBLP (25 μM)), 1 μL of template (the recombinant plasmids) and 3.5 μL ddH_2O . The reaction mixture was incubated at 65 $^\circ\text{C}$ for 1 h then heated at 80 $^\circ\text{C}$ for 2 min in a laboratory water bath (DSY21-8, China), and the amplified products were analyzed in 15 g/L agarose in TAE buffer gel. The results could also be visualized directly with the naked eye according to the white precipitate of magnesium pyrophosphate generated in the reaction or the green color changed under 254 nm UV light by the addition of diluted SuperSYBR (61201-50, TIANDZ, Chian). To confirm the specificity of amplification products, 20 μL of the reaction mixture was digested with *Taq I* at 65 $^\circ\text{C}$ for 2 h.

Optimization of reaction conditions

In the optimum conditions assay, the RT-LAMP reaction mixture as described above were used, and the reaction conditions were determined to be as follows: a gradient of incubation degree from 60 $^\circ\text{C}$ to 65 $^\circ\text{C}$ for 30

min, 45 min and 1 h, respectively, and terminated at 80 $^\circ\text{C}$ for 2 min. In all, 5 μL of RT-LAMP products were analyzed using 15 g/L agarose gel electrophoresis.

Specificity of RHDV RT-LAMP.

The specificity of the RT-LAMP was assessed by comparing RHDV with EBHSV (recombinant plasmids constructed pGM-T-EBHSV), *Pasteurella multocida*, *E. coli* and *Salmonella* from rabbits. The DNA extraction of *Pasteurella multocida*, *E. coli* and *Salmonella* from rabbits by the common method according to the reported literature^[8]. Then amplifications of the RHDV, pGM-T-EBHSV (constructed), *Pasteurella multocida*, *E. coli* and *Salmonella* from rabbits were performed using the optimum reaction system and conditions for RT-LAMP determined above in this study.

Sensitivity of RHDV RT-LAMP.

To assess the sensitivity of RHDV RT-LAMP, the recombinant plasmids containing the RT-PCR product of RHDV (constructed above) were estimated by ND-1000 ultraviolet spectrophotometer (Nano Drop Co., Ltd, USA) and amplified in a serial of 10-fold dilution by using the optimum reaction system and conditions for RT-LAMP determined above, and compared with the RHDV RT-PCR detection method established in reported literature^[9].

Primary application of RHDV RT-LAMP


To evaluate the reliability of RHDV RT-LAMP, 5 RHDV experimentally infected samples and 30 clinical samples collected from markets were detected by using the RT-LAMP with RT-PCR established in reported literature in parallel.

RESULTS

Construction of recombinant plasmids

In order to develop the RT-LAMP for RHDV detection, a 749 bp length target DNA fragment of VP60 gene of RHDV was synthesized using RT-PCR method (see Figure 1(A)), and directly cloned into pMD-19T Vector to construct recombinant plasmid named pMD-19T-RHDV, then the recombinant plasmids were identified using agarose electrophoresis, PCR and sequencing as described in this study (see Figure

1(B)), The results showed the target DNA fragment of VP60 gene of RHDV (as shown in Figure 1) were successfully synthesized, which had 100% homology with the reference sequence published in GenBank, and pMD-19T-RHDV was extracted and stored at -20 °C for the template used in this study.



M: DNA marker DL2000; 1: RT-PCR products of RHDV-Sch01 RNA; 2: RT-PCR products of pMD-19T vector; 3: RT-PCR products of pMD-19T-RHDV; 4: pMD-19T vector; 5: pMD-19T-RHDV
Figure 1 : Agarose gel electrophoresis results of target DNA fragment of VP60 gene of RHDV(A) RT-PCR result of VP60 gene of RHDV(B) Identification results of recombinant plasmids


Basic RT-LAMP reaction

The RHDV basic RT-LAMP was carried out using pMD-19T-RHDV (constructed) as template and incubated at 65 °C for 1 h, then the amplification products were analyzed by using agarose gel electrophoresis, enzyme-digestion and visual inspection as described in the method section above. The results (as shown in Fig.2) showed the target template pMD-19T-RHDV was amplified at 65 °C with a ladder-like pattern bands from about 213 bp on the gel (Figure 2(A) lane1), which specificity were conformed by the two smaller digestion products from 100 bp to 213 bp with *Taq*^I (Figure 2(A) lane2). The positive reaction could also be visualised clearly by the white precipitate of magnesium pyrophosphate (Figure 2(B) tube3) and the green color produced after the addition of diluted Supper SYBR (Figure 2(C) tube5), while the ddH₂O or pMD-19T vector template control reactions had no amplifications and visual inspection changes (Figure 2 tube4 and 6).

Optimization of reaction conditions


The RT-LAMP reaction conditions were optimized

by different amplification temperatures and reaction time. The results (as shown in Figure 3) showed that lane3, lane4, lane5, lane6, lane7, lane8 and lane9 had more clearer ladder-like reaction bands. Then the optimal temperature and reaction times for the RT-LAMP were selected at 64 °C for 45 min for the specificity and amplification efficiency, and the 25 μL reaction volume as described in the method section above was determined, and the optimum RT reaction volume components and conditions were the same as the instructions of Primerscript RT reagent Kit.



(A) Agarose gel electrophoresis; (B) Visual inspection by turbidity; (C) visual inspection by adding Supper SYBR; M: DNA marker DL2000; 1,3,5: RT-LAMP reaction products of pMD-19T-RHDV ; 2: The positive RT-LAMP products digested with *Taq*^I; 4,6: Negative control (pMD-19T vector);

Figure 2 : Results of RHDV basic RT-LAMP




M: DNA marker DL2000; 1: 60°C 1h; 2: 61°C 1h; 3: 62°C 1h; 4: 63°C 1h; 5: 64°C 1h; 6: 65°C 1h; 7: 64°C 30min; 8: 64°C 45 min; 9: 64°C 1h

Figure 3 : The results of optimization of reaction conditions
Specificity and sensitivity of RHDV RT-LAMP

The specificity and sensitivity of RHDV RT-LAMP

FULL PAPER

were assessed in the tests as described in the method section above. There was a clear ladder-like pattern of electrophoretic bands from about 213 bp observed only for the pMD-19T-RHDV and RHDV detection (see Figure 4, lane 1 and lane 3), and no specific amplification products were obtained for the detection of pGM-T-EBHSV, *Pasteurella multocida*, *E.coli* and *Salmonella* (see Figure 4). For the sensitivity, the expected size of amplification bands of 192 bp (see Figure 5(B) lane 4, 50 copies) or ladder-like pattern from about 213 bp (see Figure 5 (A) lane 5, 5 copies) could be observed when the pMD-19T-RHDV at different concentrations were tested (as shown in Figure 5), and this result indicated the sensitivity of detection limit by RT-LAMP could reach about 5 copies of cloned viral genomic fragments, which is higher than that of RT-PCR established.




M: DNA marker DL2000; 1: pMD-19T-RHDV control; 2: ddH₂O; 3: RHDV; 4: pGM-T-EBHSV; 5: *E.coli*; 6: *Pasteurella multocida*; 7: *Salmonella*

Figure 4 : The specificity assay results of RHDV RT-LAMP on agarose gel electrophoresis

Primary application of RHDV RT- LAMP

The RHDV RT-LAMP was carried out by using 5 RHDV experimentally infected samples and 30 clinical samples with RT-PCR established in parallel. All of the 5 RHDV experimentally infected samples proved to be positive by the two methods, and there were not any amplification for the 30 clinical samples detection. The results indicated the good correlation between RHDV RT-LAMP and RT-PCR.



M: DNA marker DL2000; P: positive control of RT-PCR; N: negative control; 1-6: the copies number of pMD-19T-RHDV was about 5×10^4 , 5×10^3 , 5×10^2 ; 50, 5 and 0.5, respectively.

Figure 5 : The sensitivity assay results of RHDV RT-LAMP on agarose gel electrophoresis (A) The result obtained by RT-LAMP assay (B) The result according to RT-PCR assay by Wang et al. (2011)

DISCUSSION AND CONCLUSION

LAMP assay was regarded as a powerful molecular tool for the DNA amplification and used widely in the many fields because of its advantages of rapidity, high sensitivity, high specificity and easy manipulation^[10-12], but the requirements of the four based specially designed primers for the target amplification sequence may limit its application in the detection for all the pathogens. Yuan et al (2013) had reported a one-step RT-LAMP method for the detection of RHDV using a set of 4 basic primers and detected 7 field RHDV isolates in China successfully^[13]. In this paper, a new RT-LAMP method for RHDV detection was developed according to the VP60 gene segment using 4 basic special primers and 1 loop primer to improve the sensitivity, compared with the LAMP reported by Yuan et al. We all know European brown hare syndrome virus (EBHSV) and RHDV are both the two important members of genus Lagovirus in the family Caliciviridae, which had been confirmed by nucleotide and deduced amino acid sequence comparisons in the VP60 gene region, but many studies described recently about RHDV detection methods were few referred to EBHSV since it has not occurred in China until now, and this is the first report using a DNA fragment of 359 bp of EBHSV VP60 gene (synthesized) for the specificity evaluation of RHDV RT-LAMP in China.

In conclusion, the RT-LAMP assays described here

which performed at 64°C for 45min in a laboratory water bath and visualized the results directly with the naked eye have good sensitivity and specificity, as few as 5 copies target gene could be detected, and the primary application of RHDV RT- LAMP showed the good correlation with RT-PCR established. Therefore this RT-LAMP method supply a new convenient and useful technology for the detection of RHDV in lab and field practice in China.

ACKNOWLEDGEMENTS

This study was supported by Grants from Post-doctoral funding, Project of Sichuan Provincial Education Department (11ZB064), Sichuan Province Science and Technology Support Project (2012NZ00 05), Program for Changjiang Scholars and Innovative Research Team in University (PCS IRT0848).

REFERENCES

- [1] G.LeGall, C.Arnauld, E.Boilletot, J.P.Morisse, D.Rasschaert; *Journal of General Virology*, **79**, 11 (1998).
- [2] B.Gromadzka, B.Szewczyk, G.Konopa, A.Fitzner, A.Kesy; *Acta Biochim Pol.*, **53(2)**, 371 (2006).
- [3] F.Chen; *Veterinary Lemology*, 5th Edition. Bei jing, (2008).
- [4] M. T. McIntosh, S. C. Behan, F. M Mohamed, Z.Lu, K.E. Moran, T.G. Burrage, J.G Neilan, G.B. Ward, G. Botti, L.Capucci, S.A.Metw ally; *Virology Journal*, **4**, 96 (2007).
- [5] L. Yang, F.Wang, B.Hu, J.Xue, Y.Hu, B.Zhou, D.Wang, W.Xu; *J Virol Methods.*, **151(1)**,24 (2008).
- [6] B.D.Cooke, A.J.Robinson, J.C.Merchant, A.Nardin, L.Capucci; *Epidemiol Infect.*, **124(3)**, 563 (2000).
- [7] T.Notomi, H.Okayama, H.Masubuchi, T.Yonekawa, K.Watanabe, N.Amino, T.Hase; *Nucl. Acids Res.*, **28(12)**, E63 (2000).
- [8] Z. Yang, G.Pang, Y.Wang, X.Yao, Y. Zhang, Z. Yang, K. Wang, X.Dai; *Acta Agriculturae Boreali-occidentalis Sinica*, **19**, 8 (2010).
- [9] Y.Wang, Z. Yang, X.Han, K. Wang, X. Yao, Y.Bai; *Chinese Veterinary Science*, **41(11)**, 1165 (2011).
- [10] S.Jayawardena, C.Y.Cheung, I.Barr, K.H.Chan, H.Chen, Y.Guan, J.S.Peiris, L.L.Poon; *Emerg. Infect. Dis.*, **13**, 899 (2007).
- [11] X.Fang, W.Xiong, J.Li, Q.Chen; *J Virol Methods*, **151(1)**, 35 (2008).
- [12] S. Yin, Y.Shang, G.Zhou, H.Tian, Y.Liu, X.Cai, X.Liu; *J Biotechnol*, **146**, 147 (2010).
- [13] D. Yuan, D.Guo, J.Liu, C.Si, Q.Jiang, H.Lin, T.Yang, L.Qu; *J Virol Methods*, **187**, 274 (2013).