

Identification and Characterization of an Indigenous Hydrocarbonoclastic Bacterial Strain *Stenotrophomonas Maltophilia* Newly Isolated from Seawater and Marine Sediments of Oran Harbor, Northwestern Algeria

Faiza Bendadeche*, Mohamed Bey Baba Hamed, Sidi-Mohammed El-Amine Abi Ayad

Laboratory of Aquaculture and Bioremediation (AquaBior), Department of Biotechnology (I.G.M.O), Faculty of Nature and Life Sciences, University of Oran 1 Ahmed Ben Bella, Oran, Algeria

*Corresponding author: Faiza Bendadeche, Laboratory of Aquaculture and Bioremediation (AquaBior), Department of Biotechnology (I.G.M.O), Faculty of Nature and Life Sciences, University of Oran 1 Ahmed Ben Bella, Oran, Algeria, Tel: +213 552 641 437; E-Mail: ben.faiza.aqua@hotmail.fr

Received: May 18, 2018; Accepted: May 28, 2018; Published: June 01, 2018

Abstract

Microorganisms have an important role in the bioremediation of petroleum contaminants. That caused great concern about their persistent toxicity, carcinogenicity and difficult elimination. In this paper, a newly isolated hydrocarbon-degrading bacterium SP54N from the contaminated marine sediments and seawater at the harbor of Oran, northwestern Algeria, using Bushnell-Hass salt medium. This strain could support high concentrations of crude oil (up to 10%, v/v), and identified as *Stenotrophomonas maltophilia* by sequencing and analyzing the 16S rDNA with the BLAST program on the NCBI website. The effects of some factors on rate growth of strain SP54N, in BHSM medium supplemented with 2% (v/v) of crude oil as sole carbon and energy source, were studied. The Results show that maximum growth rate was obtained at pH 7, temperature 30°C and 12% (w/v) of salinity, at 140 rpm. Moreover, *Stenotrophomonas maltophilia* could effectively utilize crude oil as its sole carbon and energy source. Therefore, *Stenotrophomonas maltophilia* SP54N can be used as an excellent degrader to develop one eco-friendly and cost-effective method for the bioremediation of harbor of Oran, as indigenous bacteria, and marine environments polluted by oil and petroleum hydrocarbons, and could be useful for biotechnological applications.

Keywords: Bioremediation; Crude oil; 16S rDNA; Marine sediments.

Introduction

Aromatic hydrocarbons and oil are the most important environmental pollutants. Hydrocarbon contamination in seawater and sediments has been considered as an international threat for environmental reasons [1]. Hydrocarbon pollution in the harbor and marine environments emerged from the emission of waste product processing, the utilization production, drilling, transportation and storage of petroleum product and from the oil spill accidents. The discharge of petroleum products in the environment causes environmental damages affecting directly or indirectly ecological systems and human health [2,3]. Several restoration techniques have been tested to eliminate or reduce hydrocarbon contaminants in recent decades.

Citation: Bendadeche F, Hamed MBB, Sidi-Mohammed El-Amine Abiayad. Identification and Characterization of an Indigenous Hydrocarbonoclastic Bacterial Strain *Stenotrophomonas Maltophilia* Newly Isolated from Seawater and Marine Sediments of Oran Harbor, Northwestern Algeria. *Biotechnol Ind J.* 2018;14(3):164.

© 2018 Trade Science Inc.

Bioremediation is an important strategy utilizing biological activities for the rapid removal of environmental pollutants [4]. It has been confirmed that the application of to be more appropriate and practical for cleaning-up petroleum hydrocarbons in contaminated harbors and marine environments [1].

Biodegradation by natural microbial population is the best reliable mechanism to eliminate xenobiotic pollutants, as crude oil, from environment [5]. Hydrocarbon-degrading microorganisms living in marine environments would be more adapted for restoring the hydrocarbon contamination in sea. Various hydrocarbons degrading bacteria were isolated form marine environments such as *Psychrobacter sp.*, *Rodococcus sp.*, *Arthrobacter sp.*, *Micrococcus* and *Spingomonas*. Though, the number of the bacterial species capable of degrading petroleum hydrocarbons isolated from sea is less than that isolated from sediments [6-8].

Polycyclic aromatic hydrocarbons (PAHs) is a group of major environmental xenobiotic pollutants which raise significant concern over there toxicity, teratogenicity and carcinogenicity [9,10]. Previous studies confirmed that enormous bacteria, archea and fungi have the petroleum-degrading abilities [11]. However, these microorganisms were mainly isolated from terrestrial soils and sediments as well as marine sediments [12].

Our aim in present work is to isolate indigenou Hydrocarbonoclastic bacterial strains with the abilities of degrading crude oil from contaminated marine sediments and seawater of Oran harbor, and characterize the strain SP54N with its high growth rate on crude oil.

Materials and Methods

Samples collection

The samples of mixed sediment and seawater were collected from 40 m depth, at the harbor of Oran, northwestern Algeria (35°42' 44''N; 0°38'28''W) in October, 2013 (**Fig. 1**).



FIG. 1. Localization of sampling site of the Oran harbor, northwestern Algeria.

Enrichment, isolation and selection of hydrocarbon-degrading bacteria

A synthetic Bushnell Hass Mineral Salt medium (BHSM) was used for isolation of hydrocarbon-degrading bacteria. BHSM medium [13]. The pH was adjusted to 7.2, then sterilizes by autoclaving at 121 °C for 20 min. After the BHSM was cooled, it

was supplemented with 2% (v/v) of crude oil (Supplied by the Hassi-Messaoud oil Refinery, Algeria) as sole carbon and energy source. The crude oil was sterilized by 0.22 µm filter membrane.

From mixed sediments and seawater (of 40 m depth) 2 ml was taken after decantation, and added to 500 ml Erlenmeyer flask containing 100 ml BHSM, and incubated for 7 days at 30°C at 140 revolutions per minute (rpm) in a shaker incubator (Heidolph Inkubator 1000 coupled with Heidolph Unimax 1010, Germany). From the enrichment culture product, 1 ml aliquots were transferred to fresh BHSM medium for subsequent subculture. Then, 1 ml inocula from the flask were streaked on a series of four further subcultures on BHSM medium supplemented with 4%, 6%, 8% then 10% of crude oil (v/v) successively as sole carbon and energy source. Each subculture was incubated at 30°C and 140 rpm for 72 H. Inoculums from the flask of 10% (v/v) of crude oil were streaked out into BHSM agar medium supplemented with 10% of crude oil (v/v) and incubated at 30°C, for 3-7 days. Phenotypically different colonies were purified in nutrient agar medium (Sigma-Aldrich, Germany). Pure cultures were then inoculated another time in fresh BHSM medium with 10% (v/v) to be sure that it used crude oil as sole carbon and energy source, then stored at -20°C until characterization.

Morphological and biochemical characteristics

The typical biochemical and physiological characteristics of strain SP54N, including Gram staining, motility (Mannitol mobility medium), respiratory type (meat-liver agar), triple-sugar-iron test (TSI test), Citrate utilization test (Simmons' Citrate agar), were systematically analyzed according to Bergey's Manual of determination for bacteriology [14]. All mediums were purchased from Sigma-Aldrich (Germany). The oxidase activity was determined by kit oxidase test (Sigma-Aldrich, Germany) and the catalase activity was determined by bubble formation in a 3% (w/v) hydrogen peroxide solution. All biochemical tests below were performed in triplicate.

Molecular identification

DNA extraction: Genomic DNA of strain SP54N was extracted from grown cells using an EasyPure[®] Bacteria Genomic DNA Kit (TransBionovo Co., Ltd., China) by following the manufacturer's instructions. The DNA samples were stored at -20°C until use.

16S rDNA gene sequence analysis

The DNA sample were used as template for PCR amplification of the 16S rDNA gene using the universal primers ben27F (5'-AGAGTTTGATCCTGGCTC-3') and ben1492R (5'-GGTTACCTTGTTACGCTT-3'), synthesized by Sigma (Germany). The 16S rDNA gene was amplified with the following mixture: 2 µL of DNA template (20 ng), 1 µL of ben27F (25 µM), 1 µL of ben1492R (25 µM), 1.0 µL of dNTP (5 mM), 4.0 µL of MgCl₂ (25 mM), 5 µL of 10X buffer solution (20 mM), 0.5 µL of taq DNA polymerase (5 U L⁻¹), and ddH₂O up to final volume of 50 µL. PCR conditions were described as below: 95°C for 5 min; 35 cycles, 94°C for 1 min denaturation, 55°C for 1 min annealing and 72°C for 2 min extension; and final extension at 72°C for 10 min. PCR products were checked by running 1.5% agarose gel electrophoresis. This 16S rDNA was partially sequenced by Genewiz. The sequence similarity was searched by the National Centre for Biotechnology Information BLAST. Analysis of 16S rDNA gene sequence data was performed using Clustal W and the software package MEGA (Molecular Evolutionary Genetics Analysis) version 7.0 [15], and the phylogenetic tree was inferred using neighbor-joining methods.

Effects different culture conditions on the growth of the hydrocarbon degrading isolate

The effects of pH, temperature and medium salinity on growth rate of the strain SP54N were investigated using BHSM medium with 2% (v/v) of crude oil at 140 rpm shaking rate, in 250 ml sterile flasks with 50 ml BHSM medium.

In effects of culture conditions tests, initial pH in the medium was adjusted to 6.0, 7.0, 8.0 and 9.0 with 1M HCl or 1M NaOH at 30°C; the incubation temperature was controlled as 20, 25, 30, 37 and 40°C at pH 7; and the concentration of NaCl in the medium was adjusted to 0, 3, 6, 9, 12 and 15% (w/v) at 30°C and pH 7. The growth was assessed indirectly by measuring the turbidity (OD 600 nm) with spectrophotometer (Perkin Elmer Lambda 35 UV/Vis Spectrometer, USA) after 48 h of incubation.

Growth kinetic of hydrocarbon degrading isolate

The strain SP54N was grown on BHSM supplemented by 3% (w/v) of NaCl [16], and 2% of crude oil (v/v) as sole carbon and energy source, at 30°C, pH 7 and 140 rpm on an orbital shaker (Heidolph Inkubator 1000 coupled with Heidolph Unimax 1010, Germany) for about 20 days. Cultures were performed in triplicate in 250 ml Erlenmeyer flasks containing 50 ml of medium, while the growth was assessed by measuring the optical density at 600 nm every 48 h.

Nucleotide sequence submission

The 16S rDNA gene sequence of the crude oil-degrading bacteria strain SP54N was submitted in the NCBI-Gene Bank under the accession number MG738221.

Results

Isolation of hydrocarbon-degrading bacteria

Twenty four crude oil-degrading bacterial strains were isolated from sediments and seawater of Oran harbor, by enrichment culture and dilution subcultures within increasing concentration of crude oil (2-10%, v/v) and purified on plate. The degradation experiments demonstrated that these isolates exhibited their variable growth rate and degradability of crude oil in the BHSM medium with 2-10% (v/v) of crude oil as their sole carbon and energy source (data not shown). Among, strain SP54N possessed a high growth rate on crude oil as sole carbon and energy source, and supported a high concentration of crude oil (up to 10%, v/v).

Identification of stain SP54N

To determine phylogenetic relationships between Strain SP54N and other representatives strains of the genus *Stenotrophomonas*, ~1500 pb fragments of the 16S rDNA gene were amplified by PCR (**Fig. 2**) using universal primers and only 932 pb were sequenced (GenBank ID: MG738221). This continuous stretch sequence was analyzed by BLAST tool (www.ncbi.nih.gov/blast) on the National Center for Biotechnology Information (NCBI) website using the Gen Bank database. The Calculation of the sequence similarity showed that partiel 16S rDNA gene sequence of strain SP54N has a high similarity score of 99% to those of the bacterial strains *Stenotrophomonas maltophilia*. Subsequently, a phylogenetic tree was constructed using similar 16S rDNA sequences (**Fig. 3**). The phylogenetic tree was performed using the Neighbor-Joining method [17]. The bootstrap consensus tree inferred from 1000 replicates [18], using the Maximum Composite Likelihood method [19]. The analysis involved 8 nucleotide sequences, codon positions included were 1st+2nd+3 and all positions containing gaps and missing data were eliminated.




FIG. 2. Electrophoresis of the 16S rDNA PCR product of the bacterial strain SP54N on 1,5% of agarose. MT: Trans 5K DNA Marker; lane 1 and 2: 16 rDNA




FIG. 3. Phylogenetic tree of the 16S rDNA sequence of strain SP54N and the closest related species obtained from the GenBank database (NCBI). The calculation were performed according to the Neighbor-Joining method (bootstrap=1000). GenBank accession numbers are shown in parentheses. The scale bar represents 0.0020 sequence divergence.

SP54N		GTTTGATCCTGGCTCAGAGTGAACGCTGGCGGTAGGCCTAACACATGCAAGTCGAACGGC	60
<i>S. maltophilia</i> (FJ906801.1)		GTTTGATCCTGGCTCAGAGTGAACGCTGGCGGTAGGCCTAACACATGCAAGTCGAACGGC	60
<i>S. maltophilia</i> (GQ267816.1)		GTTTGATCCTGGCTCAGAGTGAACGCTGGCGGTAGGCCTAACACATGCAAGTCGAACGGC	60

SP54N		AGCACAGAGGAGCTTGCTCCTTGGGTGGCGAGTGGCGGACGGGTGAGGAATACATCGGAA	120
<i>S. maltophilia</i> (FJ906801.1)		AGCACAGAGGAGCTTGCTCCTTGGGTGGCGAGTGGCGGACGGGTGAGGAATACATCGGAA	120
<i>S. maltophilia</i> (GQ267816.1)		AGCACAGAGGAGCTTGCTCCTTGGGTGGCGAGTGGCGGACGGGTGAGGAATACATCGGAA	120

SP54N		TCTACTCTGTCTGGGGGATAACGTAGGGAAACTTACGCTAATACCGCATACGACCTACG	180
<i>S. maltophilia</i> (FJ906801.1)		TCTACTCTGTCTGGGGGATAACGTAGGGAAACTTACGCTAATACCGCATACGACCTACG	180
<i>S. maltophilia</i> (GQ267816.1)		TCTACTCTGTCTGGGGGATAACGTAGGGAAACTTACGCTAATACCGCATACGACCTACG	180

SP54N		GGTGAAGCAGGGGATCTTCGGACCTTGCAGGATTGAATGAGCCGATGTCGGATTAGCTA	240
<i>S. maltophilia</i> (FJ906801.1)		GGTGAAGCAGGGGATCTTCGGACCTTGCAGGATTGAATGAGCCGATGTCGGATTAGCTA	240
<i>S. maltophilia</i> (GQ267816.1)		GGTGAAGCAGGGGATCTTCGGACCTTGCAGGATTGAATGAGCCGATGTCGGATTAGCTA	240

SP54N		GTGGCGGGTAAAGGCCACCAGGCGACGATCCGTAGCTGGTCTGAGAGGATGATCAG	300
<i>S. maltophilia</i> (FJ906801.1)		GTGGCGGGTAAAGGCCACCAGGCGACGATCCGTAGCTGGTCTGAGAGGATGATCAG	300
<i>S. maltophilia</i> (GQ267816.1)		GTGGCGGGTAAAGGCCACCAGGCGACGATCCGTAGCTGGTCTGAGAGGATGATCAG	300

SP54N		CCCACTGGAAGTGAAGACCGTCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGG	360
<i>S. maltophilia</i> (FJ906801.1)		CCCACTGGAAGTGAAGACCGTCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGG	360
<i>S. maltophilia</i> (GQ267816.1)		CCCACTGGAAGTGAAGACCGTCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGG	360

SP54N		ACAATGGGCGCAAGCCTGATCCAGCCATACCAGCTGGGTGAAGAAGCCCTTCGGGTTGTA	420
<i>S. maltophilia</i> (FJ906801.1)		ACAATGGGCGCAAGCCTGATCCAGCCATACCAGCTGGGTGAAGAAGCCCTTCGGGTTGTA	420
<i>S. maltophilia</i> (GQ267816.1)		ACAATGGGCGCAAGCCTGATCCAGCCATACCAGCTGGGTGAAGAAGCCCTTCGGGTTGTA	420

SP54N		AAGCCCTTTTGTGGGAAAGAAATCCAGCTGGCTAATACCCGGTGGGATGACGGTACCC	480
<i>S. maltophilia</i> (FJ906801.1)		AAGCCCTTTTGTGGGAAAGAAATCCAGCTGGCTAATACCCGGTGGGATGACGGTACCC	480
<i>S. maltophilia</i> (GQ267816.1)		AAGCCCTTTTGTGGGAAAGAAATCCAGCTGGCTAATACCCGGTGGGATGACGGTACCC	480

SP54N		AAAGAATAAGCACCGGCTAACTTCGTGCGCAGCAGCCGCGGTAATACGAAAGGGTGCAGCG	540
<i>S. maltophilia</i> (FJ906801.1)		AAAGAATAAGCACCGGCTAACTTCGTGCGCAGCAGCCGCGGTAATACGAAAGGGTGCAGCG	540
<i>S. maltophilia</i> (GQ267816.1)		AAAGAATAAGCACCGGCTAACTTCGTGCGCAGCAGCCGCGGTAATACGAAAGGGTGCAGCG	540

SP54N		TTACTCGGAATTACTGGGCGTAAAGCGTGCGTAGGTGCTCGTTAAGTCCGTTGTGAAAG	600
<i>S. maltophilia</i> (FJ906801.1)		TTACTCGGAATTACTGGGCGTAAAGCGTGCGTAGGTGCTCGTTAAGTCCGTTGTGAAAG	600
<i>S. maltophilia</i> (GQ267816.1)		TTACTCGGAATTACTGGGCGTAAAGCGTGCGTAGGTGCTCGTTAAGTCCGTTGTGAAAG	600

SP54N		CCCTGGGCTCAACCTGGGCTGCGAGTGGATACTGGGCGACTAAGTGGTAGAGGGTA	660
<i>S. maltophilia</i> (FJ906801.1)		CCCTGGGCTCAACCTGGGCTGCGAGTGGATACTGGGCGACTAAGTGGTAGAGGGTA	660
<i>S. maltophilia</i> (GQ267816.1)		CCCTGGGCTCAACCTGGGCTGCGAGTGGATACTGGGCGACTAAGTGGTAGAGGGTA	660

SP54N		GCGGAATTCCTGGTGTAGCAGTGAATGCGTAGAGATCAGGAGGAACATCCATGGCGAAG	720
<i>S. maltophilia</i> (FJ906801.1)		GCGGAATTCCTGGTGTAGCAGTGAATGCGTAGAGATCAGGAGGAACATCCATGGCGAAG	720
<i>S. maltophilia</i> (GQ267816.1)		GCGGAATTCCTGGTGTAGCAGTGAATGCGTAGAGATCAGGAGGAACATCCATGGCGAAG	720

SP54N		GCAGCTACCTGGACCAAGCTGACACTGAGGCACGAAAGCGTGGGAGCAACAGGATTA	780
<i>S. maltophilia</i> (FJ906801.1)		GCAGCTACCTGGACCAAGCTGACACTGAGGCACGAAAGCGTGGGAGCAACAGGATTA	780
<i>S. maltophilia</i> (GQ267816.1)		GCAGCTACCTGGACCAAGCTGACACTGAGGCACGAAAGCGTGGGAGCAACAGGATTA	780

SP54N		GATACCCTGGTAGTCCACGCCCTAAACGATGCGAACTGGATGTTGGGTGCAATTTGGCAC	840
<i>S. maltophilia</i> (FJ906801.1)		GATACCCTGGTAGTCCACGCCCTAAACGATGCGAACTGGATGTTGGGTGCAATTTGGCAC	840
<i>S. maltophilia</i> (GQ267816.1)		GATACCCTGGTAGTCCACGCCCTAAACGATGCGAACTGGATGTTGGGTGCAATTTGGCAC	840

SP54N		GCAGTATCGAAGCTAACCGGTTAAGTTCGCCGCCCTGGGGAGTACGGTCCGCAAGACTGAAA	900
<i>S. maltophilia</i> (FJ906801.1)		GCAGTATCGAAGCTAACCGGTTAAGTTCGCCGCCCTGGGGAGTACGGTCCGCAAGACTGAAA	900
<i>S. maltophilia</i> (GQ267816.1)		GCAGTATCGAAGCTAACCGGTTAAGTTCGCCGCCCTGGGGAGTACGGTCCGCAAGACTGAAA	900

FIG. 4. Alignment sequence of partial 16S rDNA sequence gene of unknown stain SP54N and the two closest strains *Stenotrophomonas maltophilia* (code number in GenBank: in parentheses).

The phylogenetic tree indicated that the bacterial strain SP54N was related species *Stenotrophomonas maltophilia* (**Fig. 3**). Thus, strain SP54N was identified as *Stenotrophomonas maltophilia*. To highlight sequence similarity and differences between strain SP54N and the two closest strains PSM-5 (GQ267816.1) and PSM-2 (FJ906801.1) from GenBank matches, ClustalW multiple sequence alignment were used [20], and results are shown in **Fig. 4**. Results exposed in **Figs. 3 and 4** showed that strain SP54N has some differences with related strains, so that it could be a new species of *Stenotrophomonas maltophilia*. The colonies of strain SP54N were light yellow, circular, smooth, convex, with entire margins and 1-2 mm in diameter after incubation at 30°C for 24 h on nutrient agar plate (**Fig. 5a**). The gram strain showed that strain SP54N belonged one of Gram-negative rod-shaped bacteria (**Fig. 5c**). The **Fig. 2b** shows colonies of strain SP54N in BHSM agar medium with 10% of crude oil (v/v). The results of biochemical characteristics of strain SP54N are summarized in **TABLE 1**.




FIG. 5. Morphological properties of the colonies and cells of strain SP54N, (a) Colonies of strain SP54N on the nutrient agar plate; (b) Colonies of strain SP54N on the BHMS medium agar plate with 10% (v/v) of crude oil; (c) Gram-negative bacterium of strain SP54N (10 x 100).

TABLE 1. Morphological and biochemical characteristics among SP54N strain.

Characteristics	Bacteria strain SP54N
Gram staining	-
Shape	Rod shaped
Oxidase test	-
Catalase test	+
Motility	+
Mannitol	-
Respiratory type	Strict aerobia
TSI test	K/K
H ₂ S production	-
Gas production	-
Citrate utilization	-
Notes: "+", positive reaction; "-", negative	

reaction.

Effects of different biodegradation conditions

In order to understand the crude oil degradation environmental characteristics by the isolate strain SP54N, the environmental factors affecting degradation were investigated, including initial pH, temperature, and NaCl concentration.

Results obtained from the effect of different pH on growth rate of strain SP54N in crude oil as sole carbon and energy source are represented in **Fig. 6a**. As shown in this curve, the strain could grow at pH 6-9 with the optimum growth rate at pH 7.

Effects of different temperature in growth rate of strain SP54N on crude oil as sole carbon and energy source are illustrated in **Fig. 6b**. The results demonstrated that this strain could utilize crude oil at temperature 20°C-37°C, with maximum growth rate at temperature 30°C. Strain SP54N can growth in temperatures less than 20°C (data not shown).

Results obtained from the effect of different concentration of NaCl on rate growth of strain SP54N on crude oil as sole carbon and energy source are represented in **Fig. 6c**. From the curve, it can be seen that the optimum concentration of NaCl for highest growth rate of strain SP54N was 12% (w/v). However, this bacterial strain can tolerate high salt concentrations (up to 15%, w/v).




FIG. 6. Effects of pH (A); temperature (B); and medium salinity (C) and on growth rate of strain SP54N on BHSM medium supplemented with 2% crude oil (v/v) as sole carbon and energy source for 48h at 140 rpm. Experiments were performed in triplicate.

Growth Kinetic of hydrocarbon degrading isolate

The bacterial stain SP54N was cultured on BHSM medium with 2% (v/v) of crude oil as sole carbon and energy source over 20 days. The optical density at 600 nm was measured each 48 h to assessed bacterial growth, and the results are shown in **Fig. 7.** The strain started the logarithmic growth phase from the first to 13th day, and then the stationary phase was attained at day 14.




FIG. 7. Growth kinetic of the SP54N strain on crude oil (v/v) as sole source of carbon and energy over 20 days. Bars represent standard deviation.

Discussion

Strain SP54N was one of hydrocarbon-degrading bacteria belonging to *Stenotrophomonas maltophilia*, which was firstly isolated from seawater and sediment of harbor of Oran-Algeria, and had potential to develop a method to remediate marine environments polluted by petroleum hydrocarbons.

The molecular identification of the SP54N strain was considered. Therefore, the partial sequence of 16s rDNA gene (932 bp) confirmed the identification as *Stenotrophomonas maltophilia* specie with 99% of similarity. Furthermore, the morphological characteristics of strain SP54N were very similar to those of the reported strains *Stenotrophomonas maltophilia*. The results of biochemical characteristics of strain SP54N, summarized in table 01, confirmed that this strain may be a new species of *Stenotrophomonas maltophilia*.

Several strains of *Stenotrophomonas sp.* were previously reported to own their hydrocarbon-degrading capacity [21,22]. Other *Stenotrophomonas sp.* strains have demonstrated the ability to degrade organic pollutants, such as pesticides [23], insecticides [24], chitin [25], steroid hormones [26] and Keratin [27]. Also, this species possessed the ability to remediate the hydrocarbon-contaminated environments [28,29].

Strains of *Stenotrophomonas maltophilia* were capable of utilizing crude oil as a sole carbon and energy source [30], and could polycyclic aromatic hydrocarbons (PAHs), such as naphthalene, fluorene and anthracene [31], Phenanthrene [32,33], benzo[a]pyrene [33]. In addition, Guzik et al. [34] showed that *Stenotrophomonas maltophilia* (strain KB2) was capable to

degrade high concentrations of aromatic compounds, and might utilize phenol as sole carbon and energy source of, and different kind of aromatic compounds such as protocatechuic acid (3,4-DBH), 4-hydroxybenzoic acid (4-HB), benzoic acid, vanillic acid and catechol.

Mahjoubi et al. [7], showed that *Stenotrophomonas maltophilia*, isolated from petroleum contaminated sediments and seawater at the refinery harbor of the Bizerte coast, in the north of Tunisia, were the most active hydrocarbonoclastic isolated strain with significant production of biosurfactants and high emulsification activity of crude oil (up to 46%), and suggest by their results that *Stenotrophomonas maltophilia* may constitute potential candidate for bioremediation, and could be useful for biotechnological applications. That, it was demonstrate that this specie had ability to utilize a wide range of aromatic substrate as sole carbon source, and could degrade in 48 hours 80% of phenanthrene comparing to *Pseudomonas* that degraded 70% of phenanthrene in 40 days [35,36]. Moreover, *Stenotrophomonas maltophilia* is a very powerful and useful means in the biotreatment of sole and wastewater decontamination [7].

The study of effect of culture conditions on petroleum hydrocarbons degradation is very important, because it can provide information about the microorganism and their growth requirements, before any contribution for bioremediation. To reveal the crude-oil-degrading conditions of strain SP54N, pH from 5 to 9, temperature from 20°C to 40°C and NaCl concentration from 0 to 15% (w/v) were investigated, as shown in **Fig. 6a, b and c**.

Maximum growth was obtained at pH 7.0 (**Fig. 6a**). This result is in agreement with those obtained by Neethu et al. [37], and by Kumari et al. [30], who shown that the maximum growth rate of *Stenotrophomonas maltophilia*, isolated from contaminated soil refinery was at pH 7.5. According to Vandecasteele [38], extreme pH inhibited the biodegradation of the oil. However in the marine environment as fresh water, the favorable range of pH was between 7 and 8.

From the curve of **Fig. 6b**, we note that our isolate strain grow in a wide range of temperature, from 20°C to 37°C (and less than 20°C, data not shown). These results are according with those obtained with Rouf et al. [39]. On the other hand, Rodriguez-Blanco and Antoine [40] reported that the degradation of crude oil or diesel might as well be done at 25, 10 or 4°C in Mediterranean water with controlled conditions.

Temperature is major factor that influence biodegradation rate, by effecting directly bacterial metabolism and growth rate. In addition, temperature can affect the physical and chemical properties of oil [41].

Concerning salinity, it may have a positive impact on hydrocarbon degrading aptitude. There is correlation between mineralization rate of several hydrocarbons and salinity [42]. However, contamination by both salt and oil posed problem for cleaning up using bioremediation, because externally added bacteria do not support high salinity [43].

In this study, results obtained for optimum salinity test showed that strain SP54N supported a high concentration of NaCl and grown in large range of salinity (0-15%, w/v), with maximum growth rate at 12% (w/v) of NaCl (**Fig. 6c**). These results consist with those obtained by Arulazhagan et al. [31] and Bayat et al. [44]. Although, Roder et al. [45] showed that *Stenotrophomonas maltophilia* decreased growth rate when exposed to more than 3% (w/v) of NaCl. However, other studies reported similar range of salinity for growth of *Alcanivorax sp.* and *Marinobacter sp.* strains [46].

The isolate strain from contaminated marine sediments and sea water in our study tolerate significant concentrations of NaCl (up to 15%, w/v) and crude oil (up to 10%, v/v) because of the polluted area and mechanisms of induction of developed enzymes of interest and this tolerance may reflect the evolutionary adaptation. Thus, strain SP54N may be a new species of *Stenotrophomonas maltophilia*.

This adaptation translates in high stability which gives bacteria the ability to answer more quickly to new source of hydrocarbons then bacteria without spoilage [47-50].

For results obtained from kinetic of biodegradation of crude oil by the isolated bacterial strain (**Fig. 7**), the stationary phase was attained at day 14. These results indicate that this bacterial strain had effectively the ability to degrade crude oil. Salinity was taken only at 3% and not at 12% of NaCl (w/v) which represented the optimal salinity of the strain SP54N, in order to use this bacterial strain with other bacterial strains in mixed culture to obtained better results for bioremediation (in future studies), and these other bacterial strains may not support a high salinity concentration.

Abbreviations

(PAHs): Polycyclic Aromatic Hydrocarbons; (BHSM): Bushnell Hass Mineral Salt Medium; (TSI): Triple-Sugar-Iron

Conclusion

In the present study, a newly isolated marine petroleum hydrocarbon-degrading bacterium strain SP54N identified as *Stenotrophomonas maltophilia*, from mixture seawater and sediment of the Oran harbor-Algeria, according to its phenotypic and phylogenetic characteristics, possessed an outstanding growth capacity in crude oil, and excellent adaptability in pH and salinity, and could be used as a good degrader, to establish one eco-friendly, and cost-effective method for the removal of hydrocarbon contaminants in diverse marine environments polluted by crude oil, and oil spills, particularly in harbor of Oran.

Compliance with Ethical Standards

Conflict of interest: On behalf of all authors, the corresponding author states that there is no conflict of interest.

Ethical approval: This article does not contain any studies with human participants or animals performed by any of the authors.

Acknowledgments: This study was funded by Laboratory of (AquaBior), Department of Biotechnology, University of Oran 1 Ahmed Ben, Faculty of Nature and Life Sciences, Oran - Ministry of Higher Education and Scientific Research (Algeria). The authors thank scientific group of Aquaculture and Bioremediation Laboratory (AquaBior) for their fruitful help for the development of this work.

REFERENCES

1. Deng MC, Li J, Liang FR, et al. Isolation and characterization of a novel hydrocarbon-degrading bacterium *Achromobacter* sp. HZ01 from the crude oil-contaminated seawater at the Daya Bay, southern China. *Marine Poll Bull.* 2014;83(1):79-86.
2. Jiang Z, Huang Y, Xu X, et al. Advance in the toxic effects of petroleum water accommodated fraction on marine plankton. *Acta Ecologica Sinica.* 2010;30(1):8-15.
3. Gong Y, Zhao X, Cai Z, et al. A review of oil, dispersed oil and sediment interactions in the aquatic environment: influence on the fate, transport and remediation of oil spills. *Marine Poll Bull.* 2014;79(1-2):16-33.
4. Hassanshahian M, Emtiazi G, Cappello S. Isolation and characterization of crude-oil-degrading bacteria from the Persian Gulf and the Caspian Sea. *Marine pollution bulletin.* 2012;64(1):7-12.
5. Hassanshahian M, Emtiazi G, Cappello S. Isolation and characterization of crude-oil-degrading bacteria from the Persian Gulf and the Caspian Sea. *Mar Pollut Bull.* 2012; 64: 7-12.

6. Cappello S, Caruso G, Zampino D, et al. Microbial community dynamics during assays of harbour oil spill bioremediation: a microscale simulation study. *J Appl Microbiol.* 2007;102(1):184-94.
7. Hii YS, Law AT, Shazili NA, et al. Biodegradation of Tapis blended crude oil in marine sediment by a consortium of symbiotic bacteria. *Int Biodeterior Biodegradation.* 2009;63(2):142-50.
8. Mahjoubi M, Jaouani A, Guesmi A, et al. Hydrocarbonoclastic bacteria isolated from petroleum contaminated sites in Tunisia: isolation, identification and characterization of the biotechnological potential. *New Biotechnol.* 2013;30(6):723-33.
9. Varjani SJ. Microbial degradation of petroleum hydrocarbons. *Bioresource technology.* 2017 Jan 1;223:277-86.
10. Lee RF, Anderson JW. Significance of cytochrome P450 system responses and levels of bile fluorescent aromatic compounds in marine wildlife following oil spills. *Marine Poll Bull.* 2005;50(7):705-23.
11. Song X, Xu Y, Li G, et al. Isolation, characterization of *Rhodococcus* sp. P14 capable of degrading high-molecular-weight polycyclic aromatic hydrocarbons and aliphatic hydrocarbons. *Marine Poll Bull.* 2011;62(10):2122-8.
12. Liu R, Zhang Y, Ding R, et al. Comparison of archaeal and bacterial community structures in heavily oil-contaminated and pristine soils. *J Biosci Bbioeng.* 2009;108(5):400-7.
13. Liu SH, Zeng GM, Niu QY, et al. Bioremediation mechanisms of combined pollution of PAHs and heavy metals by bacteria and fungi: A mini review. *Bioresour Technol.* 2017;224:25-33.
14. Mehdi H, Giti E. Investigation of alkane biodegradation using the microtiter plate method and correlation between biofilm formation, biosurfactant production and crude oil biodegradation. *Int Biodeterior Biodegradation.* 2008;62(2):170-8.
15. Holt SG, Kriey NR, Sneath PHA, et al. *Bergey's Manual of determination for Bacteriology.* Williams and Wilkins, New York. 1998.
16. Kumar S, Stecher G, Tamura K. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol.* 2016;33(7):1 870-4.
17. Mohanram R, Jagtap C, Kumar P. Isolation, screening, and characterization of surface-active agent-producing, oil-degrading marine bacteria of Mumbai Harbor. *Mar Pollut Bull.* 2016;105(1):131-8.
18. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol.* 1987;4(4):406-25.
19. Felsenstein J. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution.* 1985;39(4):783-91.
20. Tamura K, Nei M, Kumar S. Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc Natl Acad Sci U S A.* 2004;101(30):11030-5.
21. Sievers F, Higgins DG. Clustal Omega, accurate alignment of very large numbers of sequences. In *Multiple sequence alignment methods 2014*;105-116. Humana Press, Totowa, NJ.
22. Lee EY, Jun YS, Cho KS, et al. Degradation characteristics of toluene, benzene, ethylbenzene, and xylene by *Stenotrophomonas maltophilia* T3-c. *J Air Waste Manage Assoc.* 2002;52(4):400-6.
23. Tiwari B, Manickam N, Kumari S, et al. Biodegradation and dissolution of polyaromatic hydrocarbons by *Stenotrophomonas* sp. *Bioresour Technol.* 2016;216:1102-5.

24. Chen S, Yang L, Hu M, et al. Biodegradation of fenvalerate and 3-phenoxybenzoic acid by a novel *Stenotrophomonas* sp. strain ZS-S-01 and its use in bioremediation of contaminated soils. *Appl Microbiol Biotechnol.* 2011;90(2):755-67.
25. Tang H, Li J, Hu H, et al. A newly isolated strain of *Stenotrophomonas* sp. hydrolyzes acetamiprid, a synthetic insecticide. *Process Biochem.* 2012 Dec 1;47(12):1820-5.
26. Kobayashi DY, Reedy RM, Bick J, et al. Characterization of a chitinase gene from *Stenotrophomonas maltophilia* strain 34S1 and its involvement in biological control. *Appl Environ Microbiol.* 2002;68(3):1047-54.
27. Li Z, Nandakumar R, Madayiputhiya N, et al. Proteomic analysis of 17 β -estradiol degradation by *Stenotrophomonas maltophilia*. *Environ Sci Technol.* 2012;46(11):5947-55.
28. Fang Z, Zhang J, Liu B, et al. Biochemical characterization of three keratinolytic enzymes from *Stenotrophomonas maltophilia* BBE11-1 for biodegrading keratin wastes. *Int Biodeterior Biodegradation.* 2013 Aug 1;82:166-72.
29. Mukherjee P, Roy P. Genomic potential of *Stenotrophomonas maltophilia* in bioremediation with an assessment of its multifaceted role in our environment. *Front Microbiol.* 2016;7:967.
30. Venkidusamy K, Megharaj M. Identification of Electrode Respiring, Hydrocarbonoclastic Bacterial Strain *Stenotrophomonas maltophilia* MK2 Highlights the Untapped Potential for Environmental Bioremediation. *Front Microbiol.* 2016;7:1965.
31. Kumari S, Regar RK, Manickam N. Improved polycyclic aromatic hydrocarbon degradation in a crude oil by individual and a consortium of bacteria. *Bioresour Technol.* 2018;254:174-9.
32. Arulazhagan P, Vasudevan N, Yeom IT. Biodegradation of polycyclic aromatic hydrocarbon by a halotolerant bacterial consortium isolated from marine environment. *Int J Environ Sci Tech.* 2010;7(4):639-52.
33. Gao S, Seo JS, Wang J, et al. Multiple degradation pathways of phenanthrene by *Stenotrophomonas maltophilia* C6. *Int Biodeterior Biodegradation* 2013;79:98-104.
34. Shuona C, Hua Y, Jingjing C, et al. Physiology and bioprocess of single cell of *Stenotrophomonas maltophilia* in bioremediation of co-existed benzo [a] pyrene and copper. *J Hazard Mat.* 2017;321:9-17.
35. Urszula G, Izabela G, Danuta W, et al. Isolation and characterization of a novel strain of *Stenotrophomonas maltophilia* possessing various dioxygenases for monocyclic hydrocarbon degradation. *Braz J Microbiol.* 2009;40(2):285-91.
36. Wojcieszynska D, Hupert-Kocurek K, Greń I, et al. High activity catechol 2, 3-dioxygenase from the cresols–Degrading *Stenotrophomonas maltophilia* strain KB2. *Int Biodeterior Biodegradation.* 2011;65(6):853-8.
37. Kahlon RS. Biodegradation and bioremediation of organic chemical pollutants by *Pseudomonas*. In *Pseudomonas: Molecular and Applied Biology* 2016;343-417.
38. Neethu CS, Rahiman KM, Rosmine E, et al. Utilization of agro-industrial wastes for the production of lipase from *Stenotrophomonas maltophilia* isolated from Arctic and optimization of physical parameters. *Biocataly Agric Biotechnol.* 2015;4(4):703-9.
39. Vandecasteele JP. Microbiologie pétrolière: Concepts implications environnementales applications

- industrielles. *Technip*. 2005;1: 534-35.
40. Rouf R, Karaba SM, Dao J, et al. *Stenotrophomonas maltophilia* strains replicate and persist in the murine lung, but to significantly different degrees. *Microbiology*. 2011;157(7):2133-42.
 41. Rodríguez-Blanco A, Antoine V, Pelletier E, et al. Effects of temperature and fertilization on total vs. active bacterial communities exposed to crude and diesel oil pollution in NW Mediterranean Sea. *Environ Pollut*. 2010;158(3):663-73.
 42. Wang Y, Feng J, Lin Q, et al. Effects of crude oil contamination on soil physical and chemical properties in Momoge wetland of China. *Chin Geogr Sci*. 2013;23(6):708-15.
 43. Shiaris MP. Phenanthrene mineralization along a natural salinity gradient in an urban estuary, Boston Harbor, Massachusetts. *Microb Ecol*. 1989;18(2):135-46.
 44. Malkawi HI, Fatmi LM, Al-deeb TM. Mutational analysis of oil degrading genes in bacterial isolates from oil contaminated soil at the Jordanian oil refinery. *World Appl Sci J*. 2009;6(2):208-20.
 45. Bayat Z, Hassanshahian M, Hesni MA. Enrichment and isolation of crude oil degrading bacteria from some mussels collected from the Persian Gulf. *Mar Pollut Bull*. 2015;101(1):85-91.
 46. Roder A, Hoffmann E, Hagemann M, et al. Synthesis of the compatible solutes glucosylglycerol and trehalose by salt-stressed cells of *Stenotrophomonas* strains. *FEMS Microbiol Lett*. 2005;243(1):219-26.
 47. Jurelevicius D, Alvarez VM, Marques JM, et al. Bacterial community response to petroleum hydrocarbon amendments in freshwater, marine, and hypersaline water-containing microcosms. *Appl Environ Microbiol*. 2013;79(19):5927-35.
 48. Girvan MS, Campbell CD, Killham K, et al. Bacterial diversity promotes community stability and functional resilience after perturbation. *Environ Microbiol*. 2005;7(3):301-13.
 49. Bordenave S, Goni-Urriza MS, Caumette P, et al. Effects of heavy fuel oil on the bacterial mat. *Appl Environ Microbiol*. 2007;73(19): 6089-97.
 50. Maila MP, Randima P, Drønen K, et al. Soil microbial communities: Influence of geographic location and hydrocarbon pollutants. *Soil Biol Biochem*. 2006;38(2):303-10.