



MOLECULAR INTERACTION STUDIES IN TERNARY GLYCINE + (AQUEOUS THIOUREA) SOLUTIONS AT 298.15 K

S. D. DEOSARKAR*, A. R. BAN, P. D. TAWDE and R. T. SAWALE

School of Chemical Sciences, Swami Ramanand Teerth Marathwada University,
NANDED – 431606 (M.S.) INDIA

ABSTRACT

Partial molar volumes (φ_v^0) and viscosity B -coefficients of glycine + aqueous thiourea solutions (0.5 and 1.5 mol·dm⁻³) were determined at 298.15 K using Masson ($\varphi_v = \varphi_v^0 + S_v \times \sqrt{c}$) and Jones-Dole relation ($\eta_r - 1/\sqrt{c} = A + B \times \sqrt{c}$), respectively. Partial molar volumes of transfer ($\Delta_{tr}\varphi_v^0$) and B -coefficients of transfer of glycine at infinite dilution from pure water to aqueous-thiourea solutions were calculated. Strong solute-solvent interactions strengthening with thiourea concentration and hydrophilic-hydrophilic as well as ion-hydrophilic or ion-polar interactions were observed in the these systems.

Key words: Glycine, Thiourea, Partial molar volumes, Viscosity B -coefficients.

INTRODUCTION

Amino acids are building blocks of protein and they contain both amino (NH₂) and acid (COOH) groups. Glycine is a smallest amino acid found in proteins and it is precursor to proteins as well as it is a building block to number of natural products and has pharmaceutical, industrial and research applications. Glycine can fit into both; hydrophilic and hydrophobic environments. It can form H-bond with water through -NH₂ and -COOH groups. Physico-chemical investigation of dissolution and solvation of amino acid like glycine in aqueous solution of thiourea are of great significance. Numbers of workers have studied the aqueous glycine system and hydration structure of glycine in different environments¹⁻⁵. Densities, partial molar volumes and heat capacities of glycine in aqueous magnesium chloride solutions at different temperatures have been studied by Lark et al.⁶, transfer volumes of glycine, from water to 1,2-butanediol-water mixtures at 298.15 K were

* Author for correspondence; E-mail: sandeo24@yahoo.co.in

studied by Xu et al.⁷ and influence of hydrotropic agents on the solute-solvent interactions in aqueous solutions of glycine at different temperatures were studied by Pattnaik et al.⁸ Physico-chemical properties of amino acids in aqueous, aqueous electrolytes and organic solutes have been extensively investigated⁹⁻¹¹. Volumetric methods are widely used to understand the solute-solvent interactions because of their high sensitivity^{12,13}. Intermolecular interactions have been largely investigated through thermodynamic methods like density, viscosity and the refractive index of solutions^{14,15}.

In continuation with our program to understand the structure and molecular interactions in solution¹⁶⁻²¹, we report the systematic study of volumetric and viscometric behavior of glycine with different concentrations in 0.5 and 1.5 mol·dm⁻³ aqueous thiourea solutions at 298.15 K.

EXPERIMENTAL

Glycine (sd fine, AR grade, > 99.0%) was used. Aqueous stock solutions of 0.5 and 1.5 mol·dm⁻³ thiourea were prepared in calibrated volumetric flasks by dissolving accurate quantity of thiourea in double distilled water. Glycine solutions with different concentrations were prepared in aqueous thiourea stock solutions by dissolving accurate amounts of glycine. The solutions were always kept in airtight flasks. Density measurements were carried out using single capillary pycnometer and weighing was done on single pan, electronic balance (± 0.001 g) using a weighing scoop. Pycnometer was calibrated with benzene and distilled water at 298.15 K. Pycnometer was kept in transparent walled constant temperature water bath for 15 min to attain thermal equilibrium. In order to get the accurate results, three sets of density measurements were performed. Viscosity measurements were carried out using Ostwald viscometer by flow time method at 298.15 K and atmospheric pressure. Viscometer was kept in the transparent walled constant temperature water bath at experimental temperature for 15 min to attain the thermal equilibrium. Flow time was recorded using electronic digital stop watch (± 0.01 s) and average of three flow times was considered for calculation of viscosity.

RESULTS AND DISCUSSION

Variation in density (ρ) and viscosity (η) with concentration of glycine for glycine + aqueous thiourea solutions at 298.15 K are graphically presented in Fig. 1 and 2. It is seen that, density and viscosity increased with increase in concentration of glycine in aqueous 0.5

and $1.5 \text{ mol}\cdot\text{dm}^{-3}$ thiourea, which is due to change in volume, modification in the structure and existence of molecular interactions in solutions. Apparent molar volume (ϕ_v) of glycine was calculated from density data²²⁻²⁵. Calculated ϕ_v values with glycine concentration are presented in Fig. 3. The ϕ_v values are positive and increased with concentration of both; glycine in 0.5 and $1.5 \text{ mol}\cdot\text{dm}^{-3}$ aqueous thiourea solutions, which indicate strong solute-solvent interactions strengthening with glycine concentration. The ϕ_v of glycine in water ($43.16 \text{ cm}^3 \text{ mol}^{-1}$) is less than its ϕ_v in aqueous thiourea solutions. Hydrophilic-ionic or hydrophilic-hydrophilic interactions in glycine + aqueous thiourea solutions results in reduction of electrostriction effect and enhancement of overall water structure²⁶.




Fig. 1: Variation in density (ρ) with concentration of glycine in ternary glycine + {aqueous-thiourea} solutions at 298.15 K




Fig. 2: Variation in viscosity (η) with concentration of glycine in ternary glycine + {aqueous-thiourea} solutions at 298.15 K




Fig. 3: Variation in apparent molar volume (ϕ_v) with concentration of in ternary glycine + {aqueous-thiourea} solutions at 298.15 K

Concentration dependence of ϕ_v was fitted to Massons linear relation^{27,28} and from the plots of ϕ_v and \sqrt{c} (Fig. 4), S_v and ϕ_v^0 was determined as slope and intercept, respectively and reported in Table 1. The ϕ_v^0 is partial molar volume at infinite dilution and it represents solute-solvent interactions and S_v represents solute-solute interactions. The ϕ_v^0 of aqueous-glycine solution at 298.15 K was found to be $43.16 \text{ cm}^3 \cdot \text{mol}^{-1}$, which is in good agreement with earlier work²⁹. The ϕ_v^0 of glycine increased with addition of thiourea in solution compared to in water and further increased with thiourea concentration, which is due to perturbation in the arrangement of water molecules in bulk water by polar effects of thiourea.




Fig. 4: Graphical determination of ϕ_v^0 and S_v for glycine + {aqueous-thiourea} solutions from ϕ_v vs. \sqrt{c} plot using Masson's relation at 298.15 K

The ϕ_v^0 values for all studied systems are positive, which indicates the presence of strong solute-solvent interactions in solution and these interactions strengthens upon addition of thiourea³⁰. Solute-solute interactions are present but these interactions are weak,

which can be recognized from small positive values of S_v compared to ϕ_v^0 . Glycine has zwitterionic structure and therefore, positive values of S_v suggests that pairwise interactions dominates over charged functional groups.

Table 1: The ϕ_v^0 , S_v and $\Delta_{tr}\phi_v^0$ for glycine + {aqueous-thiourea} solutions at 298.15 K

Property/System	Glycine + {Aqueous 0.5 mol·dm ⁻³ thiourea}	Glycine + Aqueous 1.5 mol·dm ⁻³ thiourea}
ϕ_v^0 (cm ³ ·mol ⁻¹)	43.92	45.08
S_v (cm ³ ·Kg ^{3/2} ·mol ^{-3/2})	9.88	11.59
$\Delta_{tr}\phi_v^0$ (cm ³ ·mol ⁻¹)	0.76	1.92

Partial molar volumes of transfer ($\Delta_{tr}\phi_v^0$) at infinite dilution from pure water (43.16 cm³ mol⁻¹) to aqueous thiourea were calculated³⁰. The $\Delta_{tr}\phi_v^0$ is due to resultant electrostriction because of solute-solvent interaction³¹. Values $\Delta_{tr}\phi_v^0$ are reported in Table 1. The $\Delta_{tr}\phi_v^0$ values are positive, which is attributed to decrease in volume of shrinkage because of interactions between glycine and thiourea and existence of hydrophilic-hydrophilic interaction as per cosphere overlap model developed by Gurney³² and ion-hydrophilic or ion-polar interactions.

This is also supported from positive value of slope (S_v) of Massons plots. Reduction in electrostriction occurs due to less hydration of glycine in presence of thiourea as thiourea takes water molecule to hydrate and glycine will leave with less water molecules, which enhances structure of water. Hydration of glycine decreases due to existence of solute-co-solute ionic/polar-hydrophilic interactions.

The result also indicates absence of hydrophilic-hydrophobic interactions. Positive $\Delta_{tr}\phi_v^0$ values indicate dominating hydrophilic-hydrophilic or ion-hydrophilic interactions during the overlap of hydration cospheres and strengthening of these interactions with thiourea concentration.

Relative viscosity ($\eta_r = \eta/\eta_0$) was calculated and fitted to Jones-Dole relation and viscosity-A and B coefficients were determined as intercept and slope of the plot η_r-1/\sqrt{c} versus \sqrt{c} (Fig. 5). Graphical A and B viscosity coefficients are reported in Table 2.

Strong solute-solvent interactions are confirmed from the positive values of B-coefficient. The values of viscosity A-coefficient indicates weak solute-solute interactions compared to solute-solvent interactions.




Fig. 5: Jones-Dole plots ($\eta_r - 1/\sqrt{c}$ versus \sqrt{c}) for ternary glycine + {aqueous-thiourea} solutions at 298.15 K

Table 2: Viscosity A and B -coefficients for ternary glycine + {aqueous-thiourea} solutions at 298.15 K

Property/System	Glycine + {Aqueous 0.5 mol·dm ⁻³ thiourea}	Glycine + {Aqueous 1.5 mol·dm ⁻³ thiourea}
A (dm ^{3/2} mol ^{-1/2})	0.006	-0.196
B (dm ³ mol ⁻¹)	0.260	0.624
ΔB (dm ³ mol ⁻¹)	0.038	0.402

The B -coefficient of transfer from pure water (0.222 dm³ mol⁻¹) to aqueous thiourea was calculated³⁰. Values of ΔB (Table 2) are in line with the $\Delta_{tr}\phi_v^0$ and therefore, it is concluded that the ion/charged-hydrophilic group interactions are much stronger and dominates over the ion-hydrophobic group interactions³⁰. Glycine showed preference for contact with water than thiourea and its hydrogen bond sites get solvated by water rather than thiourea. Therefore, glycine-thiourea interactions are weaker than glycine-water.

CONCLUSION

Partial molar volumes (ϕ_v), partial molar volume of transfer ($\Delta_{tr}\phi_v^0$) and viscosity B -coefficients of glycine + aqueous-thiourea solutions of 0.5 and 1.5 mol·dm⁻³ were determined at 298.15 K. Strong solute-solvent interactions strengthening with thiourea concentration and hydrophilic-hydrophilic as well as ion-hydrophilic or ion-polar interactions were observed in the studied systems. The ϕ_v^0 was found to increase with increase in concentration of thiourea. Standard transfer volume of glycine, $\Delta_{tr}\phi_v^0$ from pure

water to aqueous thiourea solutions indicated that the hydrophilic-hydrophilic and ion-hydrophilic interactions exist in studied solutions. The dependence of ΔB over concentration of co-solute is in line with the results obtained for the $\Delta_{tr}\phi^0_v$. Ion/charged-hydrophilic group interactions are stronger and dominates over the ion-hydrophobic group interactions. Strong solute-solvent interactions strengthening with concentration of thiourea were observed.

ACKNOWLEDGEMENT

Authors are thankful to the Director, School of Chemical Sciences, Swami Ramanand Teerth Marathwada University Nanded (MS) India for providing necessary facilities.

REFERENCES

1. M. Sasaki, Y. Kameda, M. Yaegashi and T. Usuki, Bull. Chem. Soc. Jpn., **76**, 2293 (2003).
2. Y. Kameda, H. Ebata, T. Usuki, O. Uemura and M. Misawa, Bull. Chem. Soc. Jpn., **67**, 3159 (1994).
3. J. L. Alonso, E. J. Cocinero, A. Lesarri, M. E. Sanz and J. C. Lopez, Angew. Chem. Int. Ed., **45**, 3471 (2006).
4. J. H. Jensen and M. S. Gordon, J. Am. Chem. Soc., **117**, 8159 (1995).
5. H. Basch and W. J. Stevens, Chem. Phys. Lett., **169**, 275 (1990).
6. B. S. Lark, P. Patyar, T. S. Banipal and N. Inshore, J. Chem. Eng. Data, **49**, 553 (2004).
7. L. Xu, C. R. Ding and R. S. Lin, J. Solution Chem., **35**, 191 (2006).
8. S. Pattnaik and U. N. Dash, J. Chem. Pharm. Res., **4**(9), 4364 (2012).
9. R. K. Wadi and R. K. Goyal, J. Solution Chem., **21**(2) 163 (1992).
10. T. Banerjee and N. Kishore, J. Solution Chem., **34**(2), 137 (2005).
11. Y. Akhtar and S. F. Ibrahim, Arab. J. Chem., **4**, 487 (2011).
12. R. Badarayani and A. Kumar, Fluid Phase Equilib., **201**, 321 (2002).
13. R. Bhat and J. C. Ahluwalia, J. Phys. Chem., **89**, 1099 (1985).
14. J. G. Baragi, M. I. Aralaguppi, T. M. Aminabhavi, M. Y. Kariduraganavar and A. S. Kittur, J. Chem. Eng. Data, **50**, 910 (2005).

15. S. L. Oswal, J. S. Desai and S. P. Ijardar, *Thermochim. Acta*, **449**, 73 (2006).
16. S. D. Deosarkar and U. B. Shaikh, *Russ. J. Phy. Chem. A*, **83(12)**, 2392 (2013).
17. S. D. Deosarkar, *Russ. J. Phy. Chem. A*, **86(10)**, 1507 (2012).
18. S. D. Deosarkar, R. T. Sawale, A. R. Ban and A. L. Puyad, *J. Chem. Pharm. Res.*, **6(2)**, 390 (2014).
19. S. D. Deosarkar, H. G. Jahagirdar and V. B. Talwatkar, *Rasayan J. Chem.* **3**, 755 (2010).
20. S. D. Deosarkar and M. L. Narwade, *Rasayan J. Chem.* **3**, 55 (2010).
21. S. D. Deosarkar and A. S. Ghatbandhe, *Russ. J. Phy. Chem. A*, **88(1)**, 32 (2014).
22. M. Nath Roy, B. Sinha, R. Dey and A. Sinha, *Int. J. Thermophys.*, **26(5)**, 1549 (2005).
23. B. Hawrylak, R. Palepu and P. R. Tremaine, *J. Chem. Thermodyn.*, **38**, 988 (2006).
24. V. R. Karanth and D. K. Bhat, *Thermochimica Acta*, **572**, 23 (2013).
25. K. Rajagopal and S. E. Gladson, *J. Chem. Thermodyn.*, **43**, 852 (2011).
26. S. Chauhan, P. Chaudhary, K. Sharma, K. Kumar and Kiran, *Chem. Pap.*, **67(11)**, 1442 (2013).
27. D. O. Masson, *Philos. Mag. Ser.*, **7**, **8(49)**, 218 (1929).
28. D. Choudhary and A. Aswar, *J. Therm. Anal. Calorim.*, **107**, 21 (2012).
29. A. Ali, S. Khan and S. Hyder, *J. Chin. Chem. Soc.*, **52(2)**, 512 (2005).
30. B. Sinha, P. K. Roy and M. N. Roy, *Acta Chim. Slov.*, **57**, 651 (2010).
31. F. Shahidi, P. G. Harreil and J. T. Edward, *J. Solut. Chem.*, **5**, 807 (1976).
32. R. W. Gurney, *Ionic Processes in Solution*, McGraw Hill, New York (1953).

Accepted : 17.07.2014