



Trade Science Inc.

Organic CHEMISTRY

An Indian Journal

Short Communication

OCAIJ, 2(4), 2006 [56-59]

N, N, N', N'-Tetraacetythylenediamin And Sodium Percarbonate Reagents As An *In Situ* Producing Of Peracetic Acid For Baeyer-Villiger Oxidation Of Ketones



Corresponding Author

Mohammed M.Hashemi
Chemistry Department,
Sharif University of Technology,
P.O.BOX: 11365-9516 Tehran (IRAN)
Fax : 0098216005718
E-mail: mhashemi@sharif.edu

Received: 11th July, 2006Accepted: 14th August, 2006Web Publication Date : 14th November, 2006

Co-Author

S.Hadi Khezri
Chemistry Department, Sharif University of Technology,
P.O.BOX: 11365-9516, Tehran (IRAN)

ABSTRACT

In-situ production of per acetic acid using N, N, N', N'-tetraacetythylenediamin and sodium percarbonate for Baeyer-Villiger oxidation of cyclic and acyclic ketones to lactones and esters in ethylacetate at reflux condition is experienced in low reaction time and high yields. © 2006 Trade Science Inc. -INDIA

KEYWORDS

Baeyer-Villiger;
Peracetic acid;
N,N,N',N'-tetraacetyl
ethylenediamin(TAED);
sodium percarbonate(SPC).


INTRODUCTION

The Baeyer-Villiger oxidation of ketones to lactones or esters is one of the most important reactions in organic chemistry^[1,2]. This oxidation is carried using m-chloro perbenzoic acid catalyzed by different Lewis acids such as sulfonic acids^[3], Nafion-H^[4], CF₃COOH^[5], hydrotalcite^[6], SnCl₄^[7], scandium triflate^[8], boron trifluoride^[9], and bismuth triflate^[10]. Recently, we reported the Baeyer-Villiger oxidation of ketones using molecular oxygen with acidic resin^[11].


Also the reaction is accelerated by bases^[12], or metal complexes^[13]. However, many of these methods suffer from drawbacks such as unsatisfactory yields and the use of large amounts and very expensive catalysts and impermanency of peracids. Thus, the development of general and facile new reagent for Baeyer-Villiger oxidation is an active field of research^[13].

N, N, N', N'-tetraacetythylenediamin (TAED) was synthesised for the first time in 1911 by Franchimont et al^[14] and since then there are so many

Short Communication



SCHEME 1: Production of TAED




SCHEME 2: In situ production of peracetic acid using TAED and SPC

reports about synthesis^[15] and modification of producing methods for it^[16]. But one of the best methods for synthesis of TAED is carried in two steps^[17]. In which the first step, the 1,2-diammoniumethan diacetate salt is converted to N, N'-diacetyethylenediamin (DAED) on heating and removing of water during the reaction. In the second step, DAED is converted to TAED in the presence of excess amount of acetic anhydride at reflux condition by removing acetic acid (SCHEME 1).

After synthesis of TAED many research have been done about application of TAED in the presence of peroxy scaffold compounds such as sodium perborate (SPB) and sodium percarbonate (SPC) in designing new detergent formulation for low temperature bleaching purposes^[18], the mechanism of this bleaching as shown in (SCHEME 2) involve nucleophilic attack of peroxide anion to electrophilic carbonyl group of (TAED) and in situ production of peracetic acid that equilibrated with its anion^[19].

RESULTS AND DISCUSSION

In situ production of peracetic acid and its anion allowed us using this reagent (TAED and SPC) for Bayer-Villiger oxidation of cyclic and acyclic ketones



SCHEME 3: Bayer-villiger oxidation of ketones using TAED and SPC

to related lactones and esters (SCHEME 3).

In order to optimize the oxidant activity of TAED and SPC, we have studied the Bayer-Villiger oxidation of acetophenone (TABLE 1, entry 3) as a model substrate with 1 equivalent of TAED using

TABLE 1: The oxidation of ketones by TAED and sodium percarbonate in acetonitril at reflux condition^a

Entry	R ¹	R ²	Time(h)	Yield (%) ^b
1		(-CH ₂) ₄	3.30	81
2		(-CH ₂) ₅	3.30	78
3	CH ₃	C ₆ H ₅	2.30	80
4	CH ₃	4-Cl-C ₆ H ₄	2.45	79
5	CH ₃	4-Br-C ₆ H ₄	2.45	84
6	C ₆ H ₅	C ₆ H ₅	4	80
7	CH ₃	C ₆ H ₅ -CH ₂	2	91
8	C ₆ H ₅	C ₆ H ₅ -CH ₂	2.15	83

^aAll products were identified by comparison of their physical and spectral data with those of authentic samples. ^bIsolated yields are based on ketones.

Short Communication

various amount of SPC. The most satisfactory result was obtained when 1.33 equivalent of SPC was used to give the desired ester quantitatively in a much shorter reaction time. Good yields were obtained when dichloromethane was used as solvent; however, it was found that ethylacetate was the best solvent for this reaction.

The reaction procedure was simple and the generality of this process was illustrated by the wide range of cyclic and acyclic aromatic and aliphatic ketones. The results are shown in TABLE 1.

In summary, we have introduced a simple, fast and chemoselective TAED and SPC as new reagent procedure for Baeyer-Villiger oxidation of ketones.

Taking advantages of availability, mild reaction condition via not direct using of peracids and low cost of reagent and accelerating the oxidation. The method provides a simple and efficient alternative to existing procedures for conducting Baeyer-Villiger reaction.

EXPERIMENTAL

Melting points were measured on the Electrothermal 9100 apparatus and are uncorrected. IR spectra were measured on a Bomen FT-IR-MB 100 spectrometer. ^1H and ^{13}C NMR spectra were measured with a Bruker DRX-500 Avance spectrometer at 500.1 and 125.8 MHz using TMS as internal standard. All chemicals were obtained from Merck or Fluka, and used without further purification.

General procedure: Ketone (2 mmol), TAED (1mmol, 0.23g), sodium percarbonate (1.33 mmol, 0.21g) and ethylacetate (10 ml) were placed in a round bottomed flask with a magnetic stirrer. The reaction was refluxed in a period of time (TABLE 1). After the completion of the reaction (the progress of the reaction was monitored by TLC), the mixture was cooled to room temperature, the reaction mixture was washed with water for extraction by-products of the reactions (DAED, Na_2CO_3 , sodium acetate) and unreacted SPC. The organic extract dried over anhydrous NaSO_4 . All the organic phase was evaporated; the crude product was isolated and purified

(silica gel column chromatography). All products are known compounds which were characterized by IR, ^1H NMR, ^{13}C NMR and mass spectral data and their melting points compared with literature reports.

ACKNOWLEDGEMENTS

We are thankful to the Sharif University of Technology Research Council for the partial support of this work.

REFERENCES

- [1] G.J.Brink, I.W.C.E.Arends, R.A.Sheldon; *Chem.Rev.*, **104**, 4105-4123 (2004).
- [2] T.Punniyamurthy, S.Velusamy, J.Iqbal; *Chem.Rev.*, **105**, 2329-2363 (2005).
- [3] B.Pandey, P.V.Dalvi; *J.Org.Chem.*, **54**, 2968-2970 (1989).
- [4] G.A.Olah, T.Yamato, P.S.Iyer, N.J.Trivedi, B.P.Singh, G.K.S.Prakash; *Mater.Chem.Phys.*, **17**, 21-30 (1987).
- [5] S.S.C.Koch, A.R.Chamberlin; *Synth.Comm.*, **19**, 829-833 (1989).
- [6] K.Kaneda, T.Y.Mashita; *Tetrahedron Lett.*, **37**, 4555-4558 (1996).
- [7] T.Sugimura, Y.Fujiwara, A.Taj; *Tetrahedron Lett.*, **38**, 6019-6022 (1997).
- [8] H.Kotski, K.Arimura, T.Araki, T.Shinohara; *Synlett.*, **4**, 462- 464 (1999).
- [9] P.A.Grieco, T.Oguri, Y.Yokoyama; *Tetrahedron Lett.*, **19**, 419-420 (1978).
- [10] M.A.Mujahid, R.Varala, S.R.Adapa; *Synth.Comm.*, **33**, 3035-3040 (2003).
- [11] M.M.Hashemi, Y.Beni; *J.Chem.Res.*, 196-197 (2000).
- [12] H.Hagiwara, H.Nagatomo, S.I.Kazayama, H.Sakai, T.Hoshi, T.Suzuki, M.Ando; *J.Chem.Soc.Perkin. Trans.I*, 457-460 (1999).
- [13] a) J.S.Yadav, B.V.S.Reddy, A.K.Basak, A.V.Narsairh; *Chem.Lett.*, **33**, 248-249 (2004).
b) P.Goswami, S.Hazarika, A.M.Das, P.Chowdhury; *Indian J.Chem.Sect.B Org.Chem.Incl.Med.Chem.*, **43**, 1275-1281 (2004).
c) G.Strukul; *Angew.Chem.Int.Ed.*, **37**, 1198-1209 (1998).
d) M.Boronat, A.Corma, M.Renz, G.Sastre, P.M. Viruela; *Chem.Eur.J.*, **11**, 6905-6915 (2005).
e) C.Bolm, O.Beckmann; *Chirality*, **12**, 523-525 (2000).

Short Communication

- f) X.Hao, O.Yamazaki, A.Yoshida, J.Nishikido; Tetrahedron Lett., **44**, 4977-4980 (2003).
- [14] A.P.N.Franchimont, J.V.Dubski; Rec.Trav.Chim.Pays-Bas, 183-185 (1911).
- [15] a) G.Kunstle, H.Spes, H.Siegl; US Patent Application, US 3 886 212 (1975).
b) R.P.Mariella, K.H.Brown; J.Org.Chem., **36**, 735-737 (1971).
- [16] a) S.J.Hannam, J.S.Saynor, A.D.Walling; UK Patent Application, GB2 078 716 A (1982).
b) G.Roscher, W.Pressler; US Patent Application, US 4 219 500 (1980).
- c) H.John, S.J.Stewart; DE Patent Application, DE 3024694 A1 (1982).
- d) P.B.Morris, H.P.A.Theodore, M.J.Crosby; UK Patent Application, GB2 106 903 A (1983).
- e) K.Koster, F.J.Carduck; DE Patent Application, DE 3609735 A1(1987).
- f) E.Pontoglio, G.Donelli; EP Patent Application, EP 0 484 634 (1992).
- [17] S.H.Khezri; MS Thesis, Chemistry Department, Tehran University, Tehran, Iran (2003).
- [18] T.J.W.Van Thoor; 'Materials and Technology', Longmans-Green-De Bussy, Amsterdam (1968).
- [19] J.Mathews; JSDC, **15**, 155-157 (1999).