



Trade Science Inc.

ISSN : 0974 - 7532

Volume 5 Issue 1

Research & Reviews in

BioSciences

Review

RRBS, 5(1), 2011 [21-26]

## New insight into the relationship between TGF- $\beta$ superfamily and noggin in hair cycle

Dongliang Zhang<sup>1</sup>, Lijuan Gu<sup>1</sup>, Qijun Wang<sup>2</sup>, Jingjie Li<sup>1</sup>, Zheng Li<sup>1</sup>, Chunyan Wang<sup>1</sup>,  
Zhen Wang<sup>1</sup>, Lei Liu<sup>1</sup>, Mira Li<sup>1</sup>, Changkeun Sung\*<sup>1</sup>

<sup>1</sup>Department of Food Science and Technology, College of Agriculture and Biotechnology, Chungnam National University, Daejeon 305-764, (REPUBLIC OF KOREA)

<sup>2</sup>College of Light Industry & Food Science, South China University of Technology, Guangzhou, 510640, (CHINA)  
E-mail: kchsung@cnu.ac.kr

Received: 9<sup>th</sup> January, 2011 ; Accepted: 19<sup>th</sup> January, 2011

### ABSTRACT

Members of TGF- $\beta$  superfamily participate in controlling hair apoptosis, and the roles of those factors in the hair cycle have not been sufficiently clear. BMPs which belong to the TGF- $\beta$  superfamily also play critical roles in governing hair growth. In the process of controlling hair growth through the interacting between BMPs and noggin, noggin not only acts as an inhibitor of BMP-2 and 4 but influences the wavy expressions of BMP-2 and 4. Furthermore, BMPs are similar with TGF- $\beta$ 1, 2, 3 in structures and biologic characters. We present the positions of TGF- $\beta$ 1, 2, 3, and among those factors TGF- $\beta$ 1, 2 have some consistency with the positions of noggin in the hair regeneration cycles. Furthermore, TGF- $\beta$ 1, 2 are known as apoptosis-inducing factors, they can cease the cell proliferation through downstream signaling factors. After checking the positions of TGF- $\beta$ 1, 2 and Noggin, we inferred that TGF- $\beta$ 1, 2 may induce hair enter into catagen phase by inhibiting cell division or noggin expression, the wavy expressions of TGF- $\beta$ 1, 2 probably specify the hair cycle.

© 2011 Trade Science Inc. - INDIA

### KEYWORDS

Transforming growth factor beta;  
Bone morphogenetic protein;  
Noggin;  
Hair cycle;  
Bone morphogenetic protein receptor.

### INTRODUCTION OF TGF- $\beta$ FAMILY

Growth factors are involved in hair morphogenesis and cycle, among those growth factors, transforming growth factor beta (TGF- $\beta$ ) family plays an important part in the hair biologic cycle. TGF- $\beta$  superfamily includes bone morphogenetic proteins (BMPs), growth and differentiation factors (GDFs), activins, inhibins, et al<sup>[1]</sup>. is important in regulating differentiation, proliferation and apoptosis of many kinds of cells. Furthermore, TGF- $\beta$  superfamily is one of those superfamilies that

have been most deeply studied. Binding to TGF- $\beta$  receptors I and II, they specify their functions by two main downstream pathways (Smad and TAK1) on many biological processes which are essential for hair, tooth and T-cells to maintain their normal functions<sup>[1,2]</sup>.

### ROLE OF TGF-1, 2, 3 IN HAIR MORPHOGENESIS AND CYCLE

In 18 days old TGF- $\beta$ 1-/- mouse, hair follicles are in early catagen, whereas in the same age of TGF- $\beta$ 1+/-

## Review

+ mouse, hair follicles are in telogen phase. More fast-proliferation cells besides hair follicles appear in TGF- $\beta$ 1<sup>-/-</sup> mouse than TGF- $\beta$ 1<sup>+/+</sup> mouse, which means TGF- $\beta$ 1 inhibits hair growth.<sup>[3]</sup> In vitro culture, TGF- $\beta$ 1 derives from dermal papilla cells and has responsibility for inhibiting epithelial cell growth.<sup>[4]</sup> This means TGF- $\beta$ 1 could cease hair growth and decide the time of hair cycle stages.

In early anagen, TGF- $\beta$ 1 is expressed only in the bulge region of the out root sheath (ORS), this part bellows the sebaceous gland.<sup>[3]</sup> At this growth phase, the stem cells are activated in the bulge area and then reinforce the dermal papilla cells, which give birth to the new hair. TGF- $\beta$ 1<sup>-/-</sup> mice lack the downstream targets of TGF- $\beta$ 1 and have fewer label-retaining cells than normal mice, most of those label-retaining cells are stem cells.<sup>[5]</sup> Those discoveries show tight relationship between stem cells and TGF- $\beta$ 1.

High expression of TGF- $\beta$ 1 residents in out root sheath at full anagen phase and catagen phase, the expression part of TGF- $\beta$ 1 extends to upside of hair follicle.<sup>[6]</sup> TGF- $\beta$ 1 expression increases, the immunostaining results show that expression locations of TGF- $\beta$ 1 are seen both in the ORS and epithelial strand during late anagen and catagen stages, nevertheless in those two growth phases, TGF- $\beta$ 1 is rarely detected in the inner root sheath (IRS).<sup>[6]</sup> In vitro culture, TGF- $\beta$ 1 has the function of blocking cell division and hair growth.

TGF- $\beta$ 2 aggregates when hair follicles are undergoing the growth phase between anagen and catagen. In this aggregating process, strong TGF- $\beta$ 2 deposition is demonstrated in the lower part of the boundary area between the dermal papilla and the germinative

matrix cells. In vitro, when hair follicles are occupying catagen-like morphological changes, TGF- $\beta$ 2 deposits in the bulb area.<sup>[2]</sup>

TGF- $\beta$ 2-deficient mice delay hair follicle development and have less hair follicle than normal mice, the reduction of hair follicle number has no relationship with the hair apoptosis process induced by TGF- $\beta$ 2. In vitro TGF- $\beta$ 2 is isolated from the basal layer of the epidermis and regulated by phosphorylated SMAD-2 which is the downstream effector of TGF- $\beta$ 2 signaling. Employing mutant mouse models and culturing keratinocytes, Fuchs, E. et al found that TGF- $\beta$ 2 signaling is necessary to transiently induce the transcription factor Snail and activate the Ras-mitogen-activated protein kinase (MAPK) pathway in the bud.<sup>[3]</sup> This may be the explanation that less developed hair follicles appear in the embryonic period in TGF- $\beta$ 2-deficient mice. However, for another two factors, TGF- $\beta$ 1 has slight influence on hair formation through inhibiting keratinocyte growth, and TGF- $\beta$ 3 has no effect on hair formation.<sup>[5]</sup>

During anagen to catagen phase, TGF- $\beta$ 2 appears in the outermost cell layer of the outer root sheath, and in this transient phase, strong TGF- $\beta$ 2 immunoreactivity appears in the lower bulb matrix cells which are adjacent to the dermal papilla.<sup>[6]</sup> TGF- $\beta$ 2 could induce catagen phase of murine and human hair follicles through inhibiting cell proliferation or inducing apoptosis of hair matrix keratinocytes.<sup>[7]</sup> In addition, TGF- $\beta$ 2 and TGF- $\beta$ 2 type II receptor resident in the regressing epithelial strands<sup>[6]</sup> and TGF- $\beta$ 2 deposits in the boundary area in early catagen, so TGF- $\beta$ 2 may decide the regression of hair at the catagen phase (Figure 1). After catagen phase, hair follicles come into the telogen phase. Molecular media-




Figure 1 : The location of TGF- $\beta$ 1, 2<sup>[2,4,24]</sup>

TABLE 1 : Molecular mediators of TGF- $\beta$  superfamily<sup>[2,4,22-25]</sup>

Factors	Location in hair cycle	Function
TGF- $\beta$ 1	In inner root sheath, out root sheath, of in mature follicle	Plays a role in inducing catagen; blocking anagen in vivo
TGF- $\beta$ 2	All expressed in developing follicle, aggregates at dermal papilla at catagen to telogen phase	Controls cell apoptosis, induces catagen
TGF- $\beta$ 3	Hair cortex and the hair cuticle in the keratogenous zone (KZ) of the upper hair bulb.	Promotes epithelial cell adhesion
TGF- $\beta$ -RI	In Out root sheath at late anagen/catagen	Signal receptor of TGF- $\beta$ isoforms; plays a role in catagen development
TGF- $\beta$ -RII	Same as TGF- $\beta$ -RI	Specifies the differentiation inner root sheath.
BMP2	Anagen bulb; subcutaneous layer	Suppresses proliferative activity and supports differentiation
BMP4	Lower follicle mesenchyme	Suppresses hair growth
Noggin	Follicular mesenchyme; dermal papilla	Suppresses activity of BMP4 and induces hair growth




Figure 2 : The location of BMP-2 and BMP-4

tors of TGF- $\beta$  superfamily are shown in TABLE 1, locations of TGF-1, 2 are demonstrated in Figure 1.

### INTERACTIONS OF BMP-2, 4 AND NOGGIN IN HAIR MORPHOGENESIS

Stem cells play an essential role in cellular specialization and pattern formation during embryogenesis and in tissue regeneration in adults.<sup>[8]</sup> At early embryonic stage, the stem cells are specified into daughter cells such as hair follicle stem cells, dermis and epidermis.<sup>[9]</sup>

Bmp signaling specify ectodermal cells to differentiate into epidermis, it begins in the neuroepithelium of embryonic period. Once the embryonic skin stem cell (SC) progenitor cells have been confirmed, the next crossroads for signaling appears to be at the juncture of hair placode formation. Placode formation is dependent

on Noggin, in the presence of excess BMPs, or the absence of the BMP-inhibitor noggin or the follicle density is reduced.<sup>[3,4]</sup> Conditional absence of the *BMPR1A* gene also results in the accumulation of large masses of undifferentiated, *Lef1*-expressing, placode-like cells, further emphasizing a role for Bmp inhibition in the early stages of HF morphogenesis.<sup>[3]</sup>

The appearance of the *BMPR1A* gene is also investigated as a positive role for Bmp signaling, especially in the differentiation of matrix cells into inner root sheath (IRS) and hair shaft lineages. Several markers of matrix cell differentiation are strongly reduced or absent following the *BMPR1A*-null mouse.<sup>[3]</sup> Using the *BMPR1A*-absence mutant mouse, Munehiro Yuhki *et al.* found the differentiation of inner root sheath, but not outer root sheath is severely impaired. The number of HFs (hair follicles) was reduced in the dermis and

## Review


subcutaneous tissue, and cycling epithelial cells were reduced in mutant mice HFs. BMPR1A is essential for inner root sheath differentiation.<sup>[5]</sup>

Nuclear  $\beta$ -catenin is also influenced in the matrix cells of BMPR1A-deficient mouse, which means that BMP signaling lies upstream of  $\beta$ -catenin signaling during matrix cell differentiation. From those evidences, Bmp signaling is required for SC activation; furthermore BMP signaling is required for the differentiation of activating SCs in adopting one or more of its six different lineages which compose the mature HF.<sup>[10]</sup>

### ROLES OF NOGGIN IN HAIR CYCLE

Noggin should be referred no matter as the inhibitor of BMP or its functions in hair development. In E15.5-17.5 mice, expression of noggin which is the

antagonist of BMP is seen in mesenchymal cells under the basal membrane of the epidermis. In adult mouse noggin is restricted to the dermal papilla and connective tissue sheath of the hair follicle.<sup>[11]</sup> Noggin functions in hair morphogenesis through inhibiting BMPs. (The expression of noggin displays in Figure 3) Enhanced expression of BMP4 or targeted inactivation of noggin results in significantly blocking the hair follicle formation and inducing the hair follicle apoptosis. The highest noggin expression level is in the early anagen phase and the expression locations of noggin coincide with the positions of epithelial stem cells marker-K15. Variety of noggin expression may influence the expression level of BMPs,<sup>[10]</sup> which relate to the expression of Lef1 and Wnt signaling, which specify hair follicle morphogenesis and cycle by exerting their functions on hair follicle stem cells.<sup>[3]</sup>



**Figure 3 : The location of noggin in hair cycle.**<sup>[11,12,23]</sup>

In early anagen there is highest noggin expression; this increased noggin expression influences the BMPs signaling in K15-positive epithelium stem cells.<sup>[10]</sup> Those stem cells are induced by the intensive noggin expression which inhibits BMPs in early anagen. Noggin's mRNA is seen in the dermal papilla and distal outer root sheath at the secondary hair germ and it also can be seen at all the anagen at the same position.<sup>[12]</sup> Those results strongly support the immunohistochemistry (IHC) results, for the locations of Noggin's mRNA expression are almost the same as the positions of Noggin. Noggin's expression in epithelium stem cells coincides with the activation of Wnt/ $\beta$ -catenin signaling at the anagen phase of HF growth cycle. Noggin induces hair growth phase in postnatal skin, the noggin-treated mice show the hair grow out but the untreated group have no new hair.<sup>[12]</sup> Mis-expression of

Noggin in transgenic mice induces premature onset of catagen in second hair follicles cycle.<sup>[13]</sup> Figure 3 shows the location of noggin in hair cycle.

### DOWNSTREAM SIGNALING FACTORS OF TGF- $\beta$ SUPERFAMILY

Members of TGF- $\beta$  signaling family exert their functions through type I and type II receptors, and those factors have been identified in vertebrates. Each member of the TGF- $\beta$  superfamily binds to a characteristic combination of type I and type II receptors.<sup>[14]</sup>

During hair follicle development in embryonic period, initially TGF- $\beta$ -RI appears in all basal epidermal layers. In later stage TGF- $\beta$ -RI immunoreactivity is demonstrated strongly in epithelia cells, and sebocytes

and interfollicular epidermal keratinocyte cells also display clear TGF- $\beta$ -RI immunoreactivity.<sup>[15]</sup> TGF- $\beta$ -RII transcripts during skin development exclusively in the mesenchyme,<sup>[16]</sup> at early stage of hair follicle morphogenesis TGF- $\beta$ -RII was restricted in epithelial cells. Within the first 3 weeks there is no TGF- $\beta$ -RII immunoreactivity in dermal papilla, fibroblasts or their precursor cells, only the epidermal region that subsequently formed a hair placode which generates a new hair follicle displays TGF- $\beta$ -RII immunoreactivity. Later, after dermal papilla formation, TGF- $\beta$ -RII appears in out root sheath and infundibulum.<sup>[15]</sup>

Postnatal mouse, TGF- $\beta$ -RI emerges in sebaceous gland in all of hair cycle phases. Early anagen, TGF- $\beta$ -RI can be detected in transit-amplifying cells and the part adjacent to dermal papilla. Later anagen, TGF- $\beta$ -RI immunoreactivity was in out root sheath and inner root sheath. At catagen phase, TGF- $\beta$ -RI expression residents in the hair bulb disappears together with the out root sheath regression, the immunoreactive cell population just emerges proximally in epithelial strand.

At telogen, TGF- $\beta$ -RI expression is restricted to the sebaceous gland and the epidermis.<sup>[15,17]</sup>

No TGF- $\beta$ -RII is detected in skin in telogen and early anagen; subsequently it can be seen in inner root sheath and lightly expressed in out root sheath. At late anagen, TGF- $\beta$ -RII reaches up to out root sheath border with epidermis, and absents in dermal papilla and hair bulb. During the regression phase, TGF- $\beta$ -RII stays in hair bulb and inner root sheath, it doesn't scatter anywhere of skin after this stage.<sup>[15]</sup> (Figure 4 is the locations of TGF- $\beta$ -RI and II in hair cycle.) TGF- $\beta$  superfamily members convey signals through type I and type II receptors and other downstream factors, termed Smads. After ligand binding, the activated receptor/ligand combines receptor-regulated Smad proteins (R-Smad), which bind to the common partner proteins (Co-Smad) and subsequently translocate to the nucleus to regulate the transcription of target genes.<sup>[1]</sup> And Smad6 and Smad-7 are inhibitory Smad proteins (I-Smad) which function in the cytoplasm as negative regulators of the intracellular signal transduction network.<sup>[18]</sup>




Figure 4 : The location of TGF- $\beta$ -RI and II in hair cycle.<sup>[15,18,26]</sup>

Receptor-regulated Smad proteins (R-Smad) include Smad-1, 2, 3, 5, 8, Smad-2 and -3 are regulated by TGF- $\beta$  and activin, whereas Smad-1, -5, and -8 are primarily activated by BMPs.<sup>[19]</sup> So far Smad-2, 4 and 7 are the mostly concerned Smads by published research of hair biology. In the wild-type mouse, however, there is very low level of endogenous Smad-2 expression in the epidermis.<sup>[20]</sup> Yang Chai, *et al.* developed transgenic mice that overexpress Smad-2 in epidermis under the control of keratin 14 promoter. Overexpression Smad-2 enhanced the expression of

Smad-4, which discovers the interaction of Smad-2 and Smad-4 in conducting or controlling TGF- $\beta$  signaling during skin development.<sup>[20]</sup> Owing to the failure of germ layer and streak formation, Smad-2 mouse dies before hair follicle germ appearance.<sup>[19,21]</sup> Overexpression of Smad-2 elevates the endogenous TGF- $\beta$ 1 level while heterozygous loss of Smad-2 reduced TGF- $\beta$ 1 expression, when Smad-2 transgene expression in epidermis, its effects on keratinocyte proliferation and differentiation.<sup>[20]</sup> Conditional Smad-4 knockout mice develop epidermal hyperplasia, progressive hair loss beginning

## Review

at the first catagen phase on P16, and spontaneous skin tumor formation later in life. Whether the elimination of Smad-4 causes the consequence of hair follicle degeneration and affects epidermal differentiation is unknown.<sup>[21]</sup>

### CLOSING REMARK

The appearance of noggin coincided with the wavy emergence of TGF- $\beta$ . In anagen phase, the noggin resident in the same parts of hair follicles with TGF- $\beta$ 1, but just noggin appears in the dermal papilla, which produces new cells to complement the hair. In TGF- $\beta$ 1 -/- mouse, hair has a longer anagen phase than normal mouse. And culture in vitro, TGF- $\beta$ 1 inhibits the hair growth, makes the hair come in to catagen phase in advance. BMPs and TGF- $\beta$ s share 30% amino acid homology and some downstream signaling pathways, for example Smad-dependent pathways.<sup>[1]</sup> Roles of TGF- $\beta$ 1 in hair cycle may be explained as follow, TGF- $\beta$ 1 inhibits the extension of out root sheath and then controls the hair growth, and this process may be achieved through suppressing noggin.

For TGF- $\beta$ 2, in the early anagen phase, the high immunohistochemistry staining shows TGF- $\beta$ 2 concentrates at the dermal papilla. TGF- $\beta$ 2 control cell apoptosis, TGF- $\beta$ 2 may stop the dermal papilla growth by blocking the noggin expression. The late telogen can also be called competent telogen. In this phase, the hair is going to regenerate. And at this phase there is no expression of TGF- $\beta$ 1 or 2, at this phase noggin was activated and stimulus the hair follicle stem cells to give birth the new hair.

### ACKNOWLEDGEMENT

I would like to express my gratitude to Dr. Chunyan Wang, Dr. Zhen Wang, Dr. Lei Liu, MD. Zheng Li, MD. Jingjie Li, whose expertise, understanding, and patience, added considerably to my graduate experience. I appreciate his vast knowledge and skill in many areas. I would like to thank the other members of my colleague, Dr. Mira Li, and Dr. Mao for the assistance they provided at all levels of the research project.

### REFERENCES

- [1] B.Klopčič, et al.; *Eur.J.Cell Biol.*, **86(11-12)**, 781-799 (2007).
- [2] T.Hibino, T.Nishiyama; *J.Dermatol.Sci.*, **35(1)**, 9-18 (2004).
- [3] C.Blanpain, E.Fuchs; *Annu.Rev.Cell Dev.Biol.*, **22**, 339-373 (2006).
- [4] E.Fuchs; *Nature*, **445(7130)**, 834-842 (2007).
- [5] M.Yuhki, et al.; *Development*, **131(8)**, 1825-1833 (2004).
- [6] T.Soma, Y.Tsuji, T.Hibino; *J.Invest.Dermatol.*, **118(6)**, 993-997 (2002).
- [7] K.Foitzik, et al.; *J.Invest.Dermatol.*, **124(6)**, 1119-1126 (2005).
- [8] J.W.Zhang, L.H.Li; *Dev.Biol.*, **284(1)**, 1-11 (2005).
- [9] J.M.Waters, G.D.Richardson, C.A.B.Jahoda; *Semin.Cell Dev.Biol.*, **18(2)**, 245-254 (2007).
- [10] J.W.Zhang, et al.; *Stem Cells*, **24(12)**, 2826-2839 (2006).
- [11] V.A.Botchkarev; *J.Invest.Dermatol.*, **120(1)**, 36-47 (2003).
- [12] V.A.Botchkarev, et al.; *FASEB J.*, **15(12)**, 2205-2214 (2001).
- [13] U.Guha, et al.; *Am.J.Pathol.*, **165(3)**, 729-740 (2004).
- [14] P.Ten Dijke, C.S.Hill; *Trends Biochem.Sci.*, **29(5)**, 265-273 (2004).
- [15] R.Paus, et al.; *J.Invest.Dermatol.*, **109(4)**, 518-526 (1997).
- [16] Y.Q.Wang, et al.; *Mech.Dev.*, **52(2-3)**, 275-289 (1995).
- [17] U.Wollina, et al.; *Histol.Histopathol.*, **11(2)**, 431-436 (1996).
- [18] H.Chang, A.L.Lau, M.M.Matzuk; *Mol.Cell. Endocrinol.*, **180(1-2)**, 39-46 (2001).
- [19] P.Owens, et al.; *J.Invest.Dermatol.*, **128(4)**, 783-90 (2008).
- [20] Y.Ito, et al.; *Dev.Biol.*, **236(1)**, 181-94 (2001).
- [21] P.Owens, et al.; *Dev.Biol.*, **322(1)**, 156-66 (2008).
- [22] K.S.Stenn, R.Paus; *Physiol.Rev.*, **81(1)**, 449-494 (2001).
- [23] M.V.Plikus, et al.; *Nature*, **451(7176)**, 340-4 (2008).
- [24] K.Foitzik, et al.; *FASEB J.*, **14(5)**, 752-760 (2000).
- [25] J.Li, et al.; *J.Biol.Chem.*, **274(7)**, 4213-4219 (1999).
- [26] T.Soma, et al.; *J.Invest.Dermatol.*, **121(5)**, 969-975 (2003).