



# BioTechnology

An Indian Journal

FULL PAPER

BTALJ, 10(1), 2014 [01-04]

## Optimization of the extraction of essential oils from *Areca catechu* L. flowers using response surface methodology

Liang Li<sup>1,2</sup>, Jianbang Ji<sup>1\*</sup>, Xiaoning Kang<sup>1</sup>, Ping Dai<sup>1</sup>, Chuanxian Fu<sup>1,2</sup>

<sup>1</sup>Institute of Processing & Design of Agroproducts, Hainan Academy of Agricultural Sciences, Haikou 571100, (CHINA)

<sup>2</sup>College of Food, Hainan University, Haikou 570228, (CHINA)

E-mail: jwllok@sina.com; jjb1578@sina.com; 13136055153@163.com; 912519503@qq.com; 370979009@qq.com

### ABSTRACT

The present work was to optimize a water vapor distillation procedure to extract essential oils from *Areca catechu* L. Flowers. A three-variable, three-level Box-Behnken experimental design (BBD) was conducted to analyse the effects of soaking time, distillation time and solid to liquid ratio on the extraction efficiency. The experimental data were fitted to a second-order polynomial equation using multiple regression analysis and also analyzed using the appropriate statistical methods. The optimisation suggested that extraction with the soaking time for 4.8h, distillation time for 5.0h and the ratio of solid to liquid 1:2.7 were the best solutions for this combination of variables, under the optimum operating conditions, the extraction yield was 0.0196%. © 2014 Trade Science Inc. - INDIA

### KEYWORDS

Areca flower (*Areca catechu* L.);  
Essential oils;  
Water vapor distillation;  
Response surface analysis (RSA).

### INTRODUCTION

Areca (*Areca catechu* L.) is one of the traditional herbal medicines used in Hainan and widely distributed in East Africa, Southeast Asia and Pacific islands. Its flower contain essential oils, arecoline<sup>[1]</sup>, phenolics<sup>[2]</sup>, amino acids<sup>[3]</sup>, is a favorite healthcare products for its nutritional content<sup>[4]</sup>. Essential oils compounds are the main body of the aroma.

The water vapor distillation, which has been one of the most frequently used methods for the isolation of volatiles, has already shown its better reliability for the extraction of volatile compounds of various spices, such as rose, clove and rosemary<sup>[5]</sup>.

The technology of areca flower essential oils extraction is analysed with the response surface methodol-

ogy (RSM), in order to optimize the technical parameters process. It will obtain more suitable essential conditions for extraction of areca flower for development and utilization, aiming at providing the scientific experiment basis for areca flower and its odoriferous substances extraction<sup>[6]</sup>.

### MATERIALS AND METHODS

#### Materials

Fresh areca flower was obtained from the Coconut Research Institute, Chinese Academy of Tropical Agricultural Sciences, Wenchang, Hainan Province, China. The samples were comminuted by a pulverizer using high speed.

## FULL PAPER

### Experimental design

On the basis of single-factor experimentation, soaking time, distillation time and solid to liquid ratio were determined. A three-factor and three-level Box-Behnken design (BBD) consisting of seventeen experimental runs was used, including three replicates at the center point<sup>[7]</sup>. Soaking time (*A*, h), distillation time (*B*, h) and solid to liquid ratio (*C*, volume per mass) were chosen for independent variables, the dependent variable was extraction yield of areca flower (*Y*, %). Other factors in the extraction were kept constant: the amount of areca flower (200 g). The three variables and their levels in coded and natural values are shown in TABLE 1. The whole design consisted of 17 experimental points generated by Box Behnken design are presented in TABLE 2.

**TABLE 1 : Independent variables and their coded and actual values used for optimization**

Independent variables	Symbol		Code levels		
	Coded	Uncoded	-1	0	1
Soaking time/h	$X_1$	$x_1$	1	3	5
Extraction time/h	$X_2$	$x_2$	1	3	5
Solid to liquid ratio (volume per mass)	$X_3$	$x_3$	1	3	5

### Essential oils compounds extraction

The water vapor distillation conditions, such as extraction temperature, amount of areca flower, water and solvent, have been optimized in our laboratory. The essential oils extraction compounds were extracted by diethyl ether. 200g of areca flowers was laced in a 1000ml round-bottom flask with purified water. The samples in flasks were heated to boiling. Collected the distillate, and extraction with diethyl ether. After cooling to ambient temperature for some hours, the diethyl ether extract was dried over anhydrous sodium sulfate. The extract was carefully concentrated at 40°C with a rotary evaporator at atmospheric pressure, and then dichloromethane was used to metered volume to 10mL.

In the present work, the extraction yield (*Y*, %) was the ratio of extract mass and test sample mass (weight).

$$Y = (W_1/W_0) \times 100\% \quad (1)$$

Where *Y* is the ratio of extraction mass in test sample areca flowers mass,  $W_1$  is the extraction mass from each run of the extraction,  $W_0$  is the test sample mass.

**TABLE 2 : Experimental designs and the results of Box-Behnken design**

Run Number	Coded Factor Values			Uncoded Factor Values			Response Values
	$X_1$	$X_2$	$X_3$	$x_1$	$x_2$	$x_3$	<i>Y</i>
1	0	0	0	3	3	3	0.0149
2	0	1	1	3	5	5	0.0167
3	1	0	-1	5	3	1	0.0149
4	0	-1	1	3	1	5	0.0051
5	-1	1	0	1	5	3	0.0140
6	1	-1	0	5	1	3	0.0061
7	1	0	1	5	3	5	0.0152
8	0	0	0	3	3	3	0.0151
9	-1	-1	0	1	1	3	0.0050
10	0	1	-1	3	5	1	0.0161
11	0	0	0	3	3	3	0.0189
12	0	0	0	3	3	3	0.0161
13	0	0	0	3	3	3	0.0122
14	0	-1	-1	3	1	1	0.0041
15	-1	0	1	1	3	5	0.0081
16	1	1	0	5	5	3	0.0195
17	-1	0	-1	1	3	1	0.0131

### Data analysis

Analysis of each set of experimental design and calculation of predicted responses were carried out using Design Expert software (version 8.0).

A second-order polynomial equation was used to fit the experimental data of the studied variables. The generalized second-order polynomial model used in the response surface analysis is shown in Eq. (2):

$$Y = \beta_0 + \sum_{i=1}^k \beta_i X_i + \sum_{i=1}^k \beta_{ii} X_i^2 + \sum_{i < j=1}^k \beta_{ij} X_i X_j \quad (2)$$

Where  $\beta_0$ ,  $\beta_i$ ,  $\beta_{ii}$ , and  $\beta_{ij}$  are the regression coefficients for intercept, linear, quadratic and interaction terms, respectively, and  $X_i$  and  $X_j$  are the independent variables. The significances of all terms in the polynomial were judged statistically by computing the *F* value at a probability (*P*) of 0.05 and 0.1.

## RESULTS AND DISCUSSION

### Fitting the models

Mean values of all responses by water vapor distillation are presented in TABLE 2. The following regression equation was obtained.

$$Y = 0.015 + 0.001938 * X_1 + 0.005775 * X_2 - 0.004125 * X_3 + 0.001125 * X_1 * X_2 + 0.00125 * X_1 * X_3 - 0.000075 * X_2 * X_3 - 0.000975 * X_1 * X_1 - 0.0033 * X_2 * X_2 - 0.001675 * X_3 * X_3 \quad (3)$$

Where  $Y$ , extraction yield of areca flower (%),  $X_1$ , soaking time(h);  $X_2$ , extraction time(h);  $X_3$ , ratio of liquid to solid. The quadratic model indicated that the linear variables  $X_1$ ,  $X_2$ ,  $X_1X_2$ ,  $X_1X_3$  demonstrated positive effects on the extraction yield, whereas the linear variables  $X_3$ , the quadratic variables  $X_1^2$ ,  $X_2^2$ ,  $X_3^2$ , and the two variable interactions  $X_2X_3$  showed negative effects.

Analysis of variance (ANOVA) was performed to investigate the adequacy of the suggested models and identify the significant factors (TABLE 3). The fitness of the quadratic polynomial model was inspected by the regression coefficient  $R^2$ .  $F$ -value and  $p$ -value can check the significances of experimental variables. Moreover, the regression model can well fit the response values when generated small  $P$  value and high  $R^2$  values.  $R^2$  was calculated to be 0.9222 for the extraction efficiency. This implies that the sample variation of 92.22% could be attributed to the independent variables and the model did not explain only 7.78% of the total variations (TABLE 3). The model's  $p$ -value of 0.0039 was less than 0.05 implying that the model is significant.

TABLE 3 : Analysis of variance for BBD

Source	SS	DF	MS	F	Pr>F
Model	0.00037645	9	0.00004183	9.21528051	0.0039
$X_1$	0.00003003	1	0.00003003	6.61637422	0.0369
$X_2$	0.00026681	1	0.00026681	58.78149343	0.0001
$X_3$	0.00000136	1	0.00000136	0.29990558	0.6010
$X_1X_2$	0.00000506	1	0.00000506	1.11535133	0.3260
$X_1X_3$	0.00000625	1	0.00000625	1.37697695	0.2790
$X_2X_3$	0.00000002	1	0.00000002	0.00495712	0.9458
$X_1^2$	0.00000400	1	0.00000400	0.88184502	0.3790
$X_2^2$	0.00004585	1	0.00004585	10.10208265	0.0155
$X_3^2$	0.00001181	1	0.00001181	2.60263137	0.1507
Residual	0.00003177	7	0.00000454		
(Lack of Fit)	0.00000657	3	0.00000219	0.34775132	0.7940
(Pure Error)	0.00002520	4	0.00000630		
Total	0.00040822	16			

### Optimization of water vapor distillation and verification of models

In common, the efficiency of the extraction is influ-

enced by multiple parameters such as temperature, time and solvent polarity, among others, and their effects may be either interactive or independent<sup>[8]</sup>. Selection of appropriate conditions is crucial for the extraction of volatile compounds. The three factors and lower, middle and upper design points for RSM in coded and uncoded values are shown in TABLE 1.

According to the experimental design, 17 experimental results are shown in TABLE 2. By ANOVA (analysis of variation) on the results, the different significances of all variation sources were obtained (TABLE 3). The linear variables  $X_2$  was statistically very significant at  $P < 0.01$ ; the linear variables  $X_1$  had significant influences ( $P < 0.05$ ) on the extraction yield; and the quadratic variables  $X_2^2$  had significant influences ( $P < 0.05$ ) on the extraction yield of aniseed; whereas the linear variables  $X_3$ , the quadratic variables  $X_1^2$  and  $X_3^2$ , the two-variable interactions  $X_1X_2$ ,  $X_1X_3$  and  $X_2X_3$  had no significant influence ( $P > 0.05$ ) on the extraction yield.

To compare the predicted results with the practical values, verification experiments were performed using the deduced optimal conditions. The mean value of extraction yields obtained from experiments was 0.0201%. Under the same conditions, the model predicted that the maximum result was 0.0196%, it was not significantly different between predicted values and experimental values within 95% confidence interval, confirming the validity and adequacy of the predicted models.

## CONCLUSIONS

The water vapor distillation showed the best extraction rates for all kinds of volatiles. An extraction method of areca flower was developed with the application of the water vapor distillation. On the basis of the preliminary investigation, optimal extraction parameters for the water vapor distillation were: the soaking time 4.8h, extraction time 5h, and the ratio of solid to liquid 1:2.7. These conditions not only could increase the extraction yield, but also greatly shortened the total experiment time. Thus, the method of the water vapor distillation has a promising application for the extraction of areca flower.

## ACKNOWLEDGEMENTS

This research work was supported by the Key

**FULL PAPER**

---

Science and Technology Project of Hainan province (ZDXM20120016).

**REFERENCES**

- [1] W.M.Zhang, B.Li, L.Han, H.D.Zhang; African Journal of Biotechnology, **16**, 3887-3892 (2009).
- [2] Yogita Chavan, Rekha S.Singhal; Innovative Food Science and Emerging Technologies, **17**, 106-113 (2012).
- [3] En-Shyh Lin, Chia-Ching Li; Journal of Medicinal Plants Research, **10**, 975-981 (2010).
- [4] F.F.Cheng, W.J.Chen, Y.L.Huang; Food Research International, **44**, 98-102 (2011).
- [5] Phineas Masango; Journal of Cleaner Production, **13**, 833-839 (2005).
- [6] B.Z.Baya, H.Aicha, B.Colette et al; Journal of Chromatography. A, **1217**, 6134-6142 (2010).
- [7] J.E.Cacace, G.Mazza; Journal of Food Engineering, **59**, 379-389 (2003b).
- [8] D.C.Montgomery; Design and analysis of experiments (5th Edition). New York: Wiley, (2001).