



# Natural Products

Trade Science Inc.

*An Indian Journal***Full Paper**

NPAIJ, 3(3), 2007 [182-186]

## Pharmacognostic evaluation of *Calendula officinalis* L. flowers

Madhurima\*<sup>1</sup>, S.H.Ansari<sup>2</sup>, Mohd.Ali<sup>2</sup>, Prawez Alam<sup>2</sup>, Md.Sohail Akhtar<sup>2</sup>, Sayeed Ahmad<sup>2</sup><sup>1</sup>Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy,  
Jamia Hamdard, Hamdard University, New Delhi-110 062, (INDIA)<sup>2</sup>Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Jamia Hamdard, New Delhi-110062, (INDIA)

Tell : +91- 9810537914; Fax: +91-11-26059663

E-mail : madhurima\_pharma@rediffmail.com

Received: 24<sup>th</sup> July, 2007 ; Accepted: 29<sup>th</sup> July, 2007

### ABSTRACT

*Calendula officinalis* is a most common medicinal and ornamental plant. The flowers of the plant has been used since long, in traditional system of medicine especially in Homeopathic system for the treatment of skin diseases as well as for its wound healing property. In the present investigations an attempt has been made for the pharmacognostic standardization of flowers of *C.officinalis* using macroscopic and microscopic characters, physico-chemical constants, phytochemical screening, heavy metal analysis and TLC fingerprint profile. Results of the study indicates that besides the characteristic features like solitary inflorescence, median veins in ligulate florets, glandular trichomes, thick dark tannin filled cells and secretory canals in involucre etc., fluorescence analysis and thin layer chromatographic pattern of different extract can serve as finger prints of drug. Hence, the data generated will have potential utilization to determine correct identity, adulterants as well as for standardization and quality control of raw material used in the development of medicaments.

© 2007 Trade Science Inc. - INDIA

### KEYWORDS

*Calendula officinalis*;  
Pharmacognostic  
evaluation;  
HPTLC;  
Microscopy;  
Physico-chemical values;  
Heavy metal analysis.

### INTRODUCTION

*Calendula officinalis* L. (Compositae) commonly known as Marigold, is an annual herb, cultivated for ornamental and medicinal properties in India, Europe and America. The plant has been employed for long time in folk therapy. Tincture and extracts of florets have been used topically to promote wound healing and to reduce inflammation; systemically, in the treatment of dysmenorrhea, angina, fever, gastritis, hypotension, jaundice and rheumatism<sup>[1-2]</sup>. The flowers have been used for the treatment of smallpox, measles, jaundice, wounds, ulcers and skin diseases<sup>[3]</sup> and are the important ingredient of several homeopathic formulations. The plant has been found to contain number of phytocon

stituents like, triterpenoides<sup>[4-5]</sup>, carotenoids<sup>[6]</sup>, volatile oils<sup>[7]</sup> and flavonoides<sup>[8]</sup>. It has been reported to possess antiinflammatory<sup>[9]</sup>, antitumour<sup>[10]</sup>, immuno stimulant<sup>[11]</sup>, hypoglycemic<sup>[12]</sup> and antidiarrheal activities<sup>[13]</sup>. In light of the importance of flowers of *C.officinalis* in traditional and modern system of medicine, it was thought worthwhile to develop quality standard for the same. Hence, in the present investigation an attempt has been made to standardize *C.officinalis* by using microscopic characters, physico-chemical values, heavy metal analysis and TLC fingerprint profile.

### MATERIALS AND METHODS

#### Chemicals and collection of plant material

All the chemicals and reagents used were of analytical grade purchased from Sigma Chemical Co. (St Louis, MO, USA) and Merck (Darmstadt, Germany).

Flowers of *C.officinalis* were collected from Herbal garden of Jamia Hamdard, New Delhi (February 2006), which was identified by Taxonomist, Department of Botany, Hamdard University. The voucher specimen (PC-015/2006) was deposited in phytochemistry research laboratory.

### Morphological studies

The morphological study was carried out for shape, size, color, odour, taste and fracture of the drug<sup>[14]</sup>.

### Microscopic studies

The microscopic studies carried out using the method described by O'Brien et al.<sup>[15]</sup>. Microphotography on different magnifications was carried out with nikon labphot 2 microscopic unit. The polarized light was used for the study of crystals, starch grains and lignified cell.

### Physico-chemical studies

Different physico-chemical values such as ash value<sup>[16]</sup>, extractive values<sup>[16]</sup>, loss on drying<sup>[14]</sup>, total tannin<sup>[17]</sup>, total resin and total fat<sup>[14]</sup> were determined.

Preliminary phytochemical screening of drug was carried out as per method described by Peach and Tracy<sup>[18]</sup>. The 5g of dried and powdered leaf was extracted in a Soxhlet apparatus with hexane, chloroform, methanol and water, successively. The extracts were dried and weighed. The presence or absence of different phyto-constituents viz. triterpenoids, steroids, alkaloids, sugar, tannins, glycosides and flavanoids etc, were detected.

Fluorescence analysis study of powdered drug material with different reagents was carried out to observe the color reactions<sup>[19]</sup>.

The bitterness value, swelling index and foaming index were determined as per the WHO protocols<sup>[20]</sup> to determine the presence of bitters, mucilage, gum and saponins, respectively.

The presence of heavy metals analysis was determined<sup>[20]</sup> to ensure the safety of drug, for its use in human being.

### High performance thin layer chromatographic fin-

### gerprints

HPTLC fingerprint profile of the drug was developed as per the TLC method described by Stahl<sup>[21]</sup>. Three extracts namely; hexane, chloroform and methanol; obtained from dried and coarse powdered of flowers *C.officinalis* were subjected to HPTLC analysis to find out the nature and approximate numbers of compounds present.

The extracts were applied (5µl each) on TLC plate (precoated silica gel G60 F<sub>254</sub> aluminium sheets, (10cm × 10cm) in triplicate with bandwidth of 8mm using CAMAG Linomat V applying device on separate plates. The chromatograms were developed upto distance of 80mm at room temperature in different solvent systems as given in TABLE 4 using previously saturated twin trough chamber (CAMAG). The developed chromatograms were scanned at different wavelengths using CAMAG TLC Scanner III and deuterium and tungston lamp in absorbance/fluorescence mode to determine best suitable wavelength showing maximum number of compounds.

## RESULTS AND DISCUSSION


### Macroscopic characters

The capitula (inflorescence) are solitary, heterogamous having ligulate florets and tubular florets. Ligulate florets were orange or orange-yellow in colour, 5-6mm in width, with 3-toothed apex and yellowish brown tube with projecting style and 2-lobed stigma while tubular florets about 7-8mm long, yellow or orange red 5-lobed corolla in colour and with orange-brown tube. The anthers are two lobed with thin connective tissue. The involucre 5-6mm in length, concave on the inner side, thick in middle and tapering towards margins. Morphological studies on the flowers were in accordance with previous data<sup>[1]</sup>.


### Microscopic characters

Outer epidermal layers of tubular floret composed with tangentially oblong dilated cells and inner epidermis with thin walled parenchyma cells. Small nest of vascular strands were seen in parenchyma zone. Glandular trichomes were mainly on inner side (Figure 1). Ligulate floret was folded twice and had three major


## Full Paper




**Figure 1: T.S. of tubular corolla**  
Abbreviations: TC, tubular corolla; St, stigma




**Figure 2: T.S. of ligulate corolla**  
Abbreviations: LV, lateral vein; MV, mid vein; Tr, trichome; IS, inner side



**Figure 3: T.S. of involucre bract**  
Abbreviations: GT, ground tissue; IE, inner epidermis; SC, secretory canal; OE, outer epidermis; VB, vascular bundle



**Figure 4: Venation pattern in the petal**  
Abbreviations: LV, lateral vein; MV, mid vein; CV, central vein



**Figure 5: Marginal vein with secretory canal and glandular trichomes**  
Abbreviations: MV, mid vein; GT, ground tissue; SC, secretory canal



**Figure 6: Pollen grains with exine**

median veins and small veinlets. It was folded along the three major veins. The veins were prominent and projected as hemispherical lump on the outer part with small vascular bundles. The outer epidermis of ligulate floret consist of three or four layers of parenchyma zones and dense outgrowth of trichomes on inner side (Figure 2). The outer epidermis of involucre composed of rectangular or squarish cells with thick dark tannin fitted cells and inner epidermis with hemispherical cells. All along the involucre, there were wide, circular secretory canals (Figure 3).

### Powder microscopy

The powdered material of the flower showed fragments of petals and pollen grains. The petal showed prominent central vein, thin lateral veins and marginal veins (Figure 4). The veins were invariably associated with the wide, undulate secretory canals. Glandular trichomes were frequently seen in the powder (Figure 5). Abundant pollen grains, spherical in shape with dense spiny exine were found in drug powder (Figure 6).

### Physico-chemical values



**TABLE 1: Percentage of loss on drying, ash and extractive value of *C.officinalis* flower**

Parameters	<i>C.officinalis</i> (mean <sup>a</sup> ±S.D)
Loss on drying	5.17±0.19
Total ash	9.12±0.02
Water soluble ash	3.67±0.59
Acid insoluble ash	1.35±0.09
Water soluble extractive	11.08±0.26
Alcohol soluble extractive	18.15±0.18

<sup>a</sup>mean value of six readings

**TABLE 2 : Preliminary phytochemical screening of *C. officinalis* flower extracts**

Extract	Chemical constituents	<i>C.officinalis</i>
Hexane	Triterpenoids	++
	Resins	-
Chloroform	Steroids	+++
	Triterpenoids	+
Alcohol	Alkaloids	-
	Flavonoids	++
	Alkaloids	-
	Reducing sugar	-
	Glycoside	+++
Water	Tannin	+
	Alkaloids	-
	Reducing sugar	-
	Saponin	++
	Tannin	-


**TABLE 3: Fluorescence analysis study of *C.officinalis* flower powder**

Solvent Used	UV light (254nm)	UV light (366nm)
Drug powder as such	Brown	Green
NaOH in methanol	Light Green	Green
NaOH in water	Cherry Red	Light Red
Benzene	Yellowish Brown	Reddish Brown
Acetone	Yellowish Brown	Light Brown
Ethyl acetate	Red	Green
Chloroform	Red	Light Green
Dil. H <sub>2</sub> SO <sub>4</sub>	Light Green	Light Green
Con. HCl	Reddish Brown	Green
Distilled water	Brown	Dark Green
50% HCl	Light Brown	Green
Dil. HNO <sub>3</sub>	Reddish Brown	Green
Conc. H <sub>2</sub> SO <sub>4</sub>	Brown	Green


The total ash, acid insoluble ash, water soluble ash and mean values of different solvent extractives have

**TABLE 4: TLC fingerprint profiles of *C.officinalis* flower extracts**


Extract	Solvent system	Wavelength(nm)	No. of spots (R <sub>f</sub> values)
Hexane	Toluene:chloroform: ethylacetate (4:4:1.0)	366	19 (0.02, 0.08, 0.15, 0.18, 0.23, 0.28, 0.33, 0.34, 0.38, 0.48, 0.64, 0.67, 0.70, 0.78, 0.80, 0.85, 0.89, 0.91, 0.93)
Chloroform	n-butanol: glacial acetic acid: water (5.6: 1.0: 1.6)	366	6 (0.02, 0.08, 0.66, 0.70, 0.76, 0.96)
Methanol	Toluene: ethyl acetate (6: 1.0)	366	17 (0.02, 0.04, 0.09, 0.13, 0.22, 0.25, 0.31, 0.40, 0.45, 0.48, 0.61, 0.65, 0.67, 0.72, 0.80, 0.90, 0.96)



**Figure 7: Successive extractive values of *C.officinalis* flower**



**Figure 8 : Physico-chemical studies of *C. officinalis* flower**



**Figure 9 : Heavy metal analysis study of *C.officinalis* flowers**

been tabulated in TABLE 1. Presence and absence of different phyto-constituents were detected (TABLE 2). Result of fluorescence analysis study was tabulated in TABLE 3. The percentage of successive extractives were calculated and depicted in histogram (Figure 7). Quantitative estimation of total tannin, resin, fat, bitterness value and swelling index were also determined and

## Full Paper

results are presented in histogram (Figure 8). The foaming index was found to be less than 100. The percentage of heavy metals were quantified and depicted in histogram (Figure 9).

### HPTLC fingerprints of flower extracts

Hexane, chloroform and methanol successive extracts were analysed by HPTLC for development of fingerprints. The chromatograms obtained after development in different solvent system followed by scanning at 366nm in absorbance mode depicted presence of number of substances in the extracts. Hexane, chloroform and methanol showed presence of 19, 6, and 17 spots respectively, with different  $R_f$  values as given in TABLE 4.

### CONCLUSION

The results of present investigation reveals that the data generated can be used for determining correct identity of flowers and detection of adulterants as well. The quality pharmacognostic standards developed could be helpful for standardization of raw materials used for development of single drug and compound formulations as well as for their quality control.

### REFERENCES

- [1] K.R.Kirtikar, B.D.Basu; 'Indian Medicinal Plants' third revised and enlarged edition, LMB Prakashan, Allahabad, India, 1951-1952 (2000).
- [2] Anonymous; 'British Herbal Pharmacopoeia', British Herbal Medicine Association, London, (1996).
- [3] Anonymous; The Wealth of India, New Delhi, Raw Materials, Publications and Information Directorate, CSIR, 2, 55 (1988).
- [4] G.Adler, Z.Kasprzyk; Phytochemistry, 14, 627-631 (1975).
- [5] J.St.Pyrek; Polish J.Chem, 53(12), 2465-2490 (1979).
- [6] H.Neukirch, M.D.Ambrosio, V.J.Dalla, A.Guerriero; Phytochem.Anal., 15(1), 30-35 (2004).
- [7] E.Bako, J.Deli, G.Toth; J.Biochem.Biophys., 53, 241-250 (2002).
- [8] L.Gracza, X.Szasz; Acta.Pharm.Hung., 38, 118-125 (1986).
- [9] E.Vidal-oliver, R.Elias, F.Faure, A.Babadjamian, F.Crespin, G.Balansard, G.Boudon; Planta Medica, 55, 73-75 (1989).
- [10] Y.Boucard-Maitre; Pharmazie, 43, 220 (1988).
- [11] H.Wagner, A.Proksch, I.Riess-Maurer, A.Vollmar, S.Odenthal, H.Stuppner, K.Jurcic, M.Turdu, J.N.Fang; Arzneimittel-Forschung, 35(7), 1069-1075 (1985).
- [12] M.Yoshikawa, T.Murakami, A.Kishi, T.Kageura, H.Matsuda; Chem.Pharm.Bull., 49(7), 863-870 (2001).
- [13] V.S.Rosario, A.Miguel, S.Zavala, P.G.Cuauhtemo, M.Rosa, G.Perez, P.G.Salud; Phytother.Res., 12, S47-S48 (1998).
- [14] Anonymous; 'Indian Pharmacopoeia', Government of India, New Delhi (1996).
- [15] T.P.O'Brien, N.Feder, M.E.Mc-Cull; Protoplasm, 59, 364-373 (1964).
- [16] Anonymous; 'Indian Pharmacopoeia', Government of India, New Delhi (1966).
- [17] F.Feigl; 'Spot test in organic Analysis', Elsevier Publishing Company, Amsterdam, (1966).
- [18] K.Peach, M.V.Tracy; 'Modern Methods of Plant Analysis', Springer-Verlag, Heidelberg, (1955).
- [19] C.J.Kokoshi, R.J.Kokoshi, P.J.Sharma; J.Am. Pharm.Assoc., 47, 715-717 (1958).
- [20] Anonymous; 'Quality control methods for medicinal plant materials', World health organization, Geneva, (1998).
- [21] E.Stahl; 'Thin Layer Chromatography', A Laboratory, Handbook, Springer, New York, (1969).