



Studies on the antimicrobial potential of *Hygrophylla spinosa* (Acanthaceae)

J.K.Das*, P.Chatterjee, Md.Azaharruddin, A.Mondal, N.Sarkar, S.Adhikary,
C.C.Kandar, S.Choudhury

Institute of Pharmacy, Jalpaiguri, West Bengal, PIN – 735 101, (INDIA)

E-mail : mailme_jayantadas@rediffmail.com

Received: 16th August, 2009 ; Accepted: 26th August, 2009

ABSTRACT

Hygrophylla spinosa is used widely in ayurveda to treat a number of ailments; it is also used as vegetable in the study area and is also used in the folk medicine of different casts and tribes in the vicinity of the study area. In our present work, we have investigated the basic phytochemical profile of the plant along with its antimicrobial potential. The plant extract showed significant antimicrobial activity as compared with standard ciprofloxacin. The preliminary phytochemical screening showed the presence of glycoside, alkaloid, phenolic compounds, proteins, tannins etc. in the extracts tested. © 2009 Trade Science Inc. - INDIA

KEYWORDS

Hygrophylla spinosa;
Antimicrobial Activity;
Phytochemical Screening.

INTRODUCTION

The quest for new antimicrobial agents seems perpetual, as microorganisms are getting resistant to the available antimicrobial agents day by day and are producing antibiotic resistant strains. So, we are screening the antimicrobial potential of locally available medicinal plants, whose use are evident in ayurveda or in folk medicine of the local communities to treat infectious disease or to accelerate the wound healing procedure. Our present work is a part of this bigger search. *Hygrophylla spinosa* (Acanthaceae) is commonly known as 'Kulekara' in Bengali, and is commonly found in India, Srilanka, Burma, Malaysia, and Nepal in water logged lands^[1,2]. Traditionally the plant is used for treating urinary tract infection, as haematinic, in jaundice, in rheumatic arthritis, in diarrhoea, as analgesic and

anti inflammatory etc^[2]. The search of earlier literature revealed that the plant is having antioxidant and free radical scavenging activity, antinoceptive activity^[3], hepatoprotective^[4], antihyperglycemic^[5] and antitumor^[6] activity. In our present work we have studied the preliminary phytochemical profile of the plant along with its antimicrobial activity. Although, the presence of proteins, polyphenolic compounds, reducing sugars, glycosides etc., has already been reported in the earlier literature^[7].

MATERIALS AND METHODS

Plant material

About 5 kg of the aerial plant material was procured from the local market in October, 2008 and was screened for defects and abnormalities; only healthy plant material was used in the study. A voucher speci-

Short Communication

men of the plant is preserved in the herbarium of Department of Pharmacognosy, Institute of Pharmacy, Jalpaiguri for future reference. 3 kg of the aerial plant part was shade dried, cut into pieces and was grounded to form plant powder.

As a part of the pharmacognostic study we have performed ash analysis^[8,9] and extractive value determination^[8,9].

Extraction procedure

The finely powdered plant material was macerated using ethanol for 48 hrs with occasional stirring, the extract was then filtered and the plant material was re-extracted with the same solvent for a further period of 48 hrs in the same manner and was filtered, the filtrates were combined and was evaporated to dryness under reduced pressure at a temperature not exceeding 40°C to get the crude extract.

Preliminary phytochemical screening

The crude ethanolic extract was subjected to preliminary phytochemical screening using specific reagents^[8,10,11]; the extract was also subjected to TLC using different solvent systems and specific derivatizing reagents^[11] were used to confirm the identity of the phytoconstituents.

Antimicrobial study

The antimicrobial study was conducted by cup-plate agar diffusion method^[10]. The different antibacterial stains used in the study were *Shigella Sonnei*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Bacillus subtilis*. Culture media was prepared and after sterilization, was taken in sterilized Petri dishes and the microorganisms were grown by pouring microbial suspension on the solidified media in Petri dishes and incubating them at 30°C temperature for 24 hrs. After growth of the microorganism, pours were made using borer and different concentration of the extracts were placed in each hole, marketed standard disks were used as standard. The Petri dishes were further incubated at 30°C for a period of 24 hrs and the diameter of the zone of inhibition was measured. The diameter of zone of inhibition for each concentration was measured thrice against each microbial strain and the result shows the average diameter of zone of inhibition.

RESULTS AND DISCUSSION

The results of different pharmacognostic studies are shown in TABLE 1; preliminary phytochemical screening of the extract revealed the presence of polyphenolic compounds, glycosides, sugars, alkaloids etc. The results of the antibacterial activity study is shown in TABLE 2, and shows that the ethanolic extract of *Hygrophylla spinosa* at a dose of 100 mg/ml has almost equally potent antibacterial activity as compared with that of standard drug (Ciprofloxacin).

TABLE 1 : Ash Analysis and Extractive Values of *Hygrophylla spinosa*

1. Ash values	
(i) Total ash	18.7% of Dry weight
(ii) Acid insoluble ash	3.67 % of Total ash
(iii) Water soluble ash	16.07 % of Total ash
(iv) Sulphated ash	21.07 % of Total ash
2.Extractive values	
(i) Pet. Ether soluble extractive values	2.1 %
(ii) Chloroform soluble extractive values	4.85 %
(iii) Ethanol soluble extractive values	2.3 %
(iv) Water soluble extractive values	6.3 %

TABLE 2 : Antimicrobial Activity of the ethanolic extract of *Hygrophylla spinosa*

Microorganism	Diameter of Zone of Inhibition (mm)			
	25 mg/ml	50 mg/ml	100 mg/ml	Standard
<i>Shigella Sonnei</i>	18	21	25	26
<i>Escherichia coli</i>	12	14	18	27
<i>Pseudomonas aeruginosa</i>	16	19	20	29
<i>Staphylococcus aureus</i>	-	08	13	23
<i>Bacillus subtilis</i>	14	21	27	28

- diameter of zone of inhibition is negligible

As the plant extract contains a number of different significant phytoconstituents, so further study should be conducted to identify and isolate the active principle from the extract.

ACKNOWLEDGEMENT

The authors like to thank the authority of Institute of Pharmacy, Jalpaiguri for providing all necessary facilities to undertake the study.

REFERENCES

- [1] Anonymous; 'The Wealth of India'. 1st supplement series (Raw Material). NISCOM, New Delhi, **3**, 319-320 (2002).
- [2] R.N.Chopra, S.L.Nayar, I.C.Chopra; 'Glossary of Indian Medicinal Plants'. CSIR, New Delhi, (1986).
- [3] P.Shanmugasundaram, S.Venkataraman; Afr.J.Trad.CAM, **2(1)**, 62- 69 (2005).
- [4] Anubha Singh, S.S.Handa; J.Ethanopharmacol, **49**, 119-126 (1995).
- [5] Prachi Garodia, Haruyo Ichikawa, Nikita Malani, Gautam Sethi, Bharat B.Aggarwal; Journal of the Society for Integrative Oncology, **5(1)(Winter)**, 001-016 (2007).
- [6] U.K.Mazumdar, M.Gupta, S.Maiti, D.Mukherjee; Indian J.Experiment Biol., **35**, 473-477 (1997).
- [7] A.Dewanji, S.Chanda, L.Si, S.Barik, S.Maiti; Plant food Human Nutr., **50**, 349-357 (1997).
- [8] C.K.Kokate; 'Practical Pharmacognosy', Pub.Vallabh Prakashan, India, (2000).
- [9] Anonymous; 'Indian Pharmacopoeia'. The Controller of Publications, Delhi, (1996).
- [10] J.H.Doughari, A.S.Ioryue; 'Pharmacologyonline', **1**, 1333-1340 (2009).
- [11] J.B.Harborne; 'Phytochemical methods – A guide to modern techniques of plant analysis', Springer (India) Pvt. Ltd, 3rd Edition, Reprint (2007).