

Study on Association of Coomassie Brilliant Blue G250 and CTAB at Different Temperatures

Juan Guo, Jilei Jing, Zhaoran Wang, Hong Lin and Nong Wang*

School of Chemical and Biological Engineering, Lanzhou Jiaotong University, Lanzhou, P. R. China

*Corresponding author: Nong Wang, School of Chemical and Biological Engineering, Lanzhou Jiaotong University, Lanzhou 730070, P R China, E-mail: wangnong07@163.com

Received: April 18, 2017; Accepted: May 05, 2017; Published: May 10, 2017

Abstract

In this paper, the association of anionic dyes Coomassie brilliant blue G250 (CBBG250) with cationic surfactant CTAB was studied by spectroscopy. It can be seen from the spectra that the characteristic absorption peak of CBBG250 is changed obviously by the addition of CTAB. We establish a reasonable association reactions physical model to obtain a series of thermodynamic parameters such as the equilibrium constant, Gibbs free energy and entropy of the association reaction at different temperatures. The experimental results show that the elevated temperature is favorable for the association reaction. The higher the temperature, the greater will be the degree of molecule association.

Keywords: CBBG250; CTAB; Equilibrium constant; Gibbs free energy; Entropy

Introduction

As the surfactant with sterilization, wetting, solubilization, emulsification, dispersion and a series of functions, it can be widely used in the leather industry [1], dyeing auxiliaries [2,3] and so on. Dyes are materials that color fibers and other materials. In recent years, dyes are not limited to the dyeing and printing of textiles, and are used in many sectors such as paint, plastics, leather, and food. The interaction of dyes with surfactants plays a very important role in textile dyeing, micellar catalysis, micellization, photography, printing inks, and some chemical studies such as biochemistry, analytical chemistry, and light perception [4]. Therefore, more and more people pay attention to the interaction between dyes and surfactants [5-12]. Coomassie brilliant blue G250 (CBBG250) is an anionic dye and is commonly used for the determination of protein content [13-15]. Cationic surfactant cetyltrimethylammonium bromide (CTAB) is widely used as an adjuvant in textile finishing processes which can improve the dyeing process [16,17]. Therefore, the exploration of the interaction between the dye and the surfactant is very valuable. In addition, many factors, such as pH [18], alcohol [19], salts [20], etc [4,21-24], may have a significant impact on the reaction process between dye and surfactant, which make the reaction mechanism extremely complicated. Usually, most of the researchers [25,26] take advantage of conductometry [19,27], UV-Vis spectroscopy [28], surface tension method [29], cyclic voltammetry, potentiometry [30], fluorescence method [31] to investigate the reaction mechanism. Many scientific literatures [32,33] point out that dye-surfactant is more likely linked by

Citation: Guo J, Jing J, Wang Z, et al. Study on Association of Coomassie Brilliant Blue G250 And CTAB at Different Temperatures. Sci Revs Chem Commun. 2017;7(1):104.

© 2017 Trade Science Inc.

hydrogen bonding, ionic bonding, van der Waals force or hydrophobic interaction. But the study of these mechanisms is still imperfect, so it is important to delve deeper into it.

Experimental

Apparatus and materials

LabTech UV-2100 UV-Vis spectrophotometer was used to measure the absorption spectra and absorbance. In order to weigh chemicals, analytical balances, model FA12048 was used. For maintaining different temperatures, a thermostat was used. The dye CBBG250 was obtained from Sinopharm Chemical reagent co., LTD. CTAB was obtained from Shanghai Zhongqin Chemical Reagent Co., Ltd, China, LTD.

Methods




FIG. 1. CBBG250 spectra: (1) CBBG250 in aqueous solution, (2) CBBG250 in CTAB aqueous solution ($C_{\text{CBBG250}} = 2 \times 10^{-5} \text{ mol/L}$, $C_{\text{CTAB}} = 1.2 \times 10^{-5} \text{ mol/L}$), (3) CTAB in aqueous solution.

Results and Discussion

The association equilibrium constant and spectrogram between CBBG250 and CTAB in aqueous solution

Assumptions the association equilibrium constants K can be expressed as

$$K = \frac{C_x}{(C_{\text{CTAB}}^0 - C_x)(C_{\text{CBBG250}}^0 - C_x)} \quad (1)$$

Where C_x is the concentration of reaction product. C_{CTAB}^0 , C_{CBBG250}^0 are respectively initial concentrations of CTAB and CBBG250.

Based on the Beer-Lambert Law, the absorbance of unit thickness absorbing layer can be expressed as

$$A = \varepsilon_{\text{CBBG250}} C_{\text{CBBG250}}^0 \quad (2)$$

Where A is the absorbance of CBBG250, $\varepsilon_{\text{CBBG250}}$ is molar absorptivity of CBBG250.

When the reaction reaches equilibrium, the measured absorbance should be expressed as

$$A' = \varepsilon_{\text{CTAB}} (C_{\text{CTAB}}^0 - C_x) + \varepsilon_{\text{CBBG250}} (C_{\text{CBBG250}}^0 - C_x) + \varepsilon_x C_x \quad (3)$$

Where $\varepsilon_{\text{CTAB}}$, ε_x is respectively molar absorptivity of CTAB and reaction product. As shown in **FIG. 1**, the molar absorption coefficient of CTAB can be ignored at the wavelength of 584 nm that is $\varepsilon_{\text{CTAB}} \approx 0$.

Then the Equation (3) can be translated into the following expression

$$A' = \varepsilon_{\text{CBBG250}} (C_{\text{CBBG250}}^0 - C_x) + \varepsilon_x C_x \quad (4)$$

From the above, the following equation can be obtained:

$$\frac{C_{\text{CTAB}}^0 \cdot C_{\text{CBBG250}}^0}{A - A'} = \frac{1}{\varepsilon_{\text{CBBG250}} - \varepsilon_x} \cdot \left[\frac{1}{K} + (\varepsilon_{\text{CTAB}} + \varepsilon_{\text{CBBG250}}) \right] - \frac{A - A'}{(\varepsilon_{\text{CBBG250}} - \varepsilon_x)^2} \quad (5)$$

At a certain temperature, $(\varepsilon_{\text{CTAB}} + \varepsilon_{\text{CBBG250}})$ is set up to be constant value.

According to chemical reaction so thermal equation, the Gibbs free energy $\Delta_r G_m$ may be expressed as

$$\Delta_r G_m = -RT \ln K \quad (6)$$

According to Gibbs - Helmholtz equation:

$$\Delta_r G_m = \Delta_r H_m - T \Delta_r S_m \quad (7)$$

$\Delta_r H_m$ and $\Delta_r S_m$ are obtained from the intercept and slope of the plot of $\Delta_r G_m$ versus T .




FIG. 2. The absorbance of CBBG250 in aqueous solutions of different concentrations at 584 nm at different temperatures.

[CBBG250]: 1.0×10^{-5} mol/L, 1.6×10^{-5} mol/L, 2.2×10^{-5} mol/L, 2.8×10^{-5} mol/L, 3.4×10^{-5} mol/L, 4.0×10^{-5} mol/L, 4.6×10^{-5} mol/L

FIG. 2 shows the absorbance of different concentrations of CBBG250 in 584 nm at different temperatures. By linear fitting we get $\epsilon_{\text{CBBG250}}$ in 584 nm at different temperatures, as shown in **TABLE 1**. It can be seen that the $\epsilon_{\text{CBBG250}}$ decreases with increasing temperature.

<i>T</i>	298.15K	308.15K	318.15K	328.15K	338.15K
$\epsilon_{\text{CBBG250}}$	42109.7 ± 84.1	41618.3 ± 160.0	41512.2 ± 196.7	41447.8 ± 130.6	41186.0 ± 163.9

TABLE 1: The $\epsilon_{\text{CBBG250}}$ in 584 nm at different temperatures

According to the model requirements, the sum of the initial concentrations of CBBG250 and CTAB is set up to be constant value (3.2×10^{-5} mol/L) at different temperatures (308.15 K, 318.15 K, 328.15 K, 338.15K) measure the absorbance of the system. At different temperature, a plot between $\frac{C_{\text{CTAB}}^0 \cdot C_{\text{CBBG250}}^0}{A - A'}$ and $A - A'$ get different straight line as shown in **FIG.**

3-6.




FIG. 3. Plot of $\frac{C_{\text{CTAB}}^0 \cdot C_{\text{CBBG250}}^0}{A - A'}$ vs. $(A - A')$ at 308.15K.




FIG. 4. Plot of $\frac{C_{CTAB}^0 \cdot C_{CBBG250}^0}{A - A'}$ vs. $(A - A')$ at 318.15K.




FIG. 5. Plot of $\frac{C_{CTAB}^0 \cdot C_{CBBG250}^0}{A - A'}$ vs. $(A - A')$ at 328.15K.




FIG. 6. Plot of $\frac{C_{CTAB}^0 \cdot C_{CBBG250}^0}{A - A'}$ vs. $(A - A')$ at 338.15K

Then K and ε_x are obtained at different temperatures from the slope and intercept based on FIG. 3-6. By the Equation (6) the obtained K can be used to calculate $\Delta_r G_m$, the results are shown in TABLE 2.

$T(K)$	$\varepsilon_x \times 10^4$	$K \times 10^5 (L/mol)$	$\Delta_r G_m (kJ/mol)$
308.15	3.328 ± 0.073	6.82 ± 0.031	-34.41 ± 0.089
318.15	3.215 ± 0.068	8.30 ± 0.067	-36.05 ± 0.213
328.15	3.191 ± 0.060	10.81 ± 0.042	-37.90 ± 0.157
338.15	3.154 ± 0.077	14.89 ± 0.180	-39.96 ± 0.205

TABLE. 2. The thermodynamic constants of CBBG250 associated with CTAB at different temperatures.

By analyzing the data in TABLE. 2, it can be seen that the association equilibrium constant K increases with increasing temperature, indicating that the temperature rise is conducive to the positive direction of the reaction. At the same time, the reaction of Gibbs free energy $\Delta_r G_m < 0$, indicating that the reaction is spontaneous, and the higher the temperature, the greater the absolute value of Gibbs free energy, it indicates that the elevated temperature plays an important role in the reaction, the higher the temperature, the greater the degree of association.

According to the Equation 7 and using the Origin software for linear fitting to get FIG. 7. Finally, we obtained the $\Delta H_m^0 = 22.703 \pm 2.147$ kJ/mol and $\Delta S_m^0 = 0.185 \pm 0.007$ kJ/mol/K from the intercept and slope of the line, respectively. $\Delta H_m^0 > 0$ shows that this reaction is an endothermic process. In general, the enthalpy change of the chemical bond is greater than 60

kJ/mol. Therefore, the association between CBBG250 and CTAB is not only the chemical bond; it is more likely linked by hydrogen bonding, ionic bonding, van der Waals force or hydrophobic interaction, and so on. $\Delta S_m^0 > 0$ indicates that the process is entropy driven.




FIG. 7. Plot of $\Delta_r G_m$ vs. T .

Conclusion

The interaction between CBBG250 and CTAB in aqueous solution can be illustrated by the redshift of the spectrum at 584 nm. According to association reactions physical model, the relevant thermodynamic parameters of CBBG250 and CTAB association at different temperatures were obtained. It can be seen that the association equilibrium constant K increases with increasing temperature, indicating that the temperature rise is conducive to the positive direction of the reaction. At the same time, the reaction of Gibbs free energy $\Delta_r G_m < 0$, indicating that the reaction is spontaneous; $\Delta H_m^0 > 0$ shows that the formation of the CTAB micelle is an endothermic process and $\Delta S_m^0 > 0$ indicates that the process is entropy driven. With the value of ΔH_m^0 , we derive the force between CBBG250 and CTAB is not only the chemical bond, it is more likely linked by hydrogen bonding, ionic bonding, van der Waals force or hydrophobic interaction, and so on.

Acknowledgements

This work was supported by the Lanzhou Municipal Science and Technology Bureau Program (Grant No.2015-4-137) and the Program Project of Science and Technology Department of Gansu Province (Grant No. 1407GCCA013).

REFERENCE

1. JE Witman and ZG Wang. A Lattice Model of Vitrification and Gelation. *J Phys Chem B*. 2006;110: 6312-6324.
2. M Weiss, RC Darton, T Battal et al. Surfactant Adsorption and Marangoni Flow in Liquid Jets. 3. Modeling in the Presence of Micellar Surfactant. *Ind Eng Chem Res*. 2006;45: 2235-2248.
3. AA Karyakin, YN Ivanova, E. E. Karyakina. Equilibrium (NAD⁺/NADH) potential on poly(Neutral Red) modified electrode. *Electrochem. Commun.* 2003;5: 677-680.
4. J Yang. Interaction of surfactants and aminoindophenol dye. *J Colloid Interface Sci*. 2004;274: 237-243.
5. N Li, SY Tong. Spectrophotometric study of the interaction of tetraphenylporphyrin tetrasulfonate (TPPS4) with proteins. *Talanta* 1994;41: 1657-1662.
6. H Akbaş, Ç Kartal. Conductometric studies of the interaction of C.I. Reactive Orange 16 with cationic alkyltrimethylammonium bromide surfactants. *Dyes Pigm.* 2007;72: 383-386.
7. H Akbaş, Ç Kartal. CI Reactive Orange 16-dodecylpyridinium chloride interactions in electrolytic solutions. *Spectrochim Acta*. 2006;65: 95-99.
8. T Shen, R Xiao, Q Wang, et al. Study of Interactions and Association Thermodynamics between Sudan Red and Sodium Dodecyl Sulfate in Microemulsions. *J Disper Sci Technol*. 2014;35: 435-440.
9. RK Dutta, SN Bhat. Interaction of Methyl Orange with Submicellar Cationic Surfactants. *B Chem Soc Jpn*. 1993;66: 2457-2460.
10. P Forte Tavčer. Interactions between some anionic dyes and cationic surfactants with different alkyl chain length studied by the method of continuous variations. *Dyes Pigm.* 2004;63: 181-189.
11. SM Ghoreishi, M Behpour, AG Farsani. Study of interaction between a cationic surfactant and two anionic azo dyes by ion-selective electrode technique and spectrophotometry. *Dyes Pigm.* 2007;74: 585-589.
12. N Wang, M Zhao. Study on Thermodynamics and Kinetics of Association Interactions between Malachite Green and OP-10 in Aqueous Solutions. *J Disper Sci Technol*. 2016;37: 190-195.
13. XX Han, Y Xie, B Zhao, et al. Highly Sensitive Protein Concentration Assay over a Wide Range via Surface-Enhanced Raman Scattering of Coomassie Brilliant Blue. *Anal Chem*. 2010;82: 4325-4328.
14. LT Jin, J K Choi. Usefulness of visible dyes for the staining of protein or DNA in electrophoresis, Electrophoresis. 2004;25: 2429-2438.
15. BM Löffler, H Kunze. Refinement of the Coomassie brilliant blue G assay for quantitative protein determination. *Anal Biochem*. 1989;177: 100-102.
16. RT Buwalda, JBFN Engberts. Aggregation of Dicationic Surfactants with Methyl Orange in Aqueous Solution. *Langmuir*. 2001;17: 1054-1059.
17. N Wang, SJ Xiao, CW Su. Preparation of molecularly imprinted polymer for methylene blue and study on its molecular recognition mechanism. *Colloid Polym Sci*. 2016;294: 1305-1314.
18. LDL Miranda, CR Bellato, MPF Fontes, et al. Preparation and evaluation of hydrotalcite-iron oxide magnetic organocomposite intercalated with surfactants for cationic methylene blue dye removal. *Chem Eng J*. 2014;254, 88-97.
19. K Małysa, R Cohen, D Exerowa et al. Steady-state foaming and the properties of thin liquid films from aqueous alcohol solutions. *J Colloid Interface Sci*. 1981;80: 1-6.

20. PK Behera, SN Panda, S Sahu, Effect of NaCl on the binding constant of styryl pyridinium dyes in anionic surfactant system. *J Mol Liq.* 2013;177: 110-113.
21. RK Dutta, SN Bhat. Dye–Surfactant Interaction in Submicellar Concentration Range. *B Chem Soc Jpn.* 1992;65: 1089-1095.
22. SS Shah, R Ahmad, SWH Shah, et al. Synthesis of cationic hemicyanine dyes and their interactions with ionic surfactants. *Colloids Surf.* 1998;137: 301-305.
23. Y Miyashita, S Hayano. Kinetic Study of the Penetration of an Anthraquinoid Acidic Dye into Cationic Micelles. *B Chem Soc Jpn.* 1981;54: 3249-3252.
24. H Lin, H Zhu, J Guo, et al. Wang, Interactions between bromophenol blue and cetyl- trimethylammonium bromide in aqueous solutions and microemulsions. *J Disper Sci Technol.* 2017;38: 1410-1414.
25. S. Chandravanshi, SK Upadhyay. Natural dye– surfactant interactions: thermodynamic and surface parameters. *Color Technol.* 2012;128: 300-305.
26. W Liu, R Guo. Interaction of flavonoid, quercetin with organized molecular assemblies of nonionic surfactant. *Colloids Surf.* 2006;274: 192-199.
27. N Wang, H Lin, H Zhu. Study of the Association Behavior Between Bromophenol Blue and Octylphenol Polyoxyethylene Ether (10) in Aqueous Solution and the Solubilization of Bromophenol Blue by Micelles. *J Solution Chem.* 2016;45: 1689-1700.
28. ZH Ren. Interacting Behavior between Amino Sulfonate Surfactant and Octylphenol Polyoxyethylene Ether in Aqueous Solution and Effect of Hydrophilicity. *Ind Eng Chem Res.* 2014;53: 10035-10040.
29. S Bračko, J Špan. Anionic dye–cationic surfactant interactions in water–ethanol mixed solvent. *Dyes Pigm.* 2001;50: 77-84.
30. B Simončič, M Kert. A study of anionic dye–cationic surfactant interactions in mixtures of cationic and nonionic surfactants. *Dyes Pigm.* 2002;54: 221-237.
31. W Liu, R Guo. The interaction between morin and CTAB aggregates. *J Colloid Interface Sci.* 2005;290: 564-573.
32. B Gohain, S Sarma, RK Dutta. Protonated dye-surfactant ion pair formation between neutral red and anionic surfactants in aqueous submicellar solutions. *J Mol Liq.* 2008;142: 130-135.
33. H Margenau. Van der waals forces. *Rev Mod Phys.* 1939;11: 1-35 (1939).