



## Zeolites: Potential soil amendments for improving nutrient and water use efficiency and agriculture productivity

Vijay Sandeep Jakkula and Suhas P Wani

International crops Research Institute for the Semi-Arid Tropics, Patancheru, Hyderabad, Telangana, India

\*Corresponding author: Vijay Sandeep Jakkula, International crops Research Institute for the Semi-Arid Tropics, Patancheru, Hyderabad, Telangana - 502324, India, Tel: +91-40-30713457; E-mail: [j.vijay@cgiar.org](mailto:j.vijay@cgiar.org)

Received: February 23, 2018; Accepted: May 07, 2018; Published: May 10, 2018

### Abstract

The fertilizer use in developing countries has shown a steady increase over the last few decades, and the use and manufacture of N fertilizers contributes to about 60% of the total release of reactive N. Higher farm subsidies and lower N fertilizer prices have further increased N inputs. Inappropriate fertilization patterns and excessive use of N fertilizer have resulted in considerable N losses through ammonia  $\text{NH}_3$  volatilization and  $\text{NO}_2$  leaching. This has meant that NUE has been as low as ~35%. An efficient crop nutrient management is important practice and thus, new designer or smart N fertilizers technologies are needed to support the increasing demand and avoid the low nitrogen use efficiency (NUE). The ammonia nitrogen volatilization and nitrate leaching can be reduced or prevented by the use of zeolite carrier material applications which have N in their framework and act as slow/controlled release fertilizers. These materials will reduce ammonia volatilization and nitrate leaching and at the same time increase crop yield. Zeolites are also known for their water holding capacity and in drylands they are the most suitable fertilizers to prolong moisture levels in severe drought like conditions. In addition to macronutrients, micronutrients can also be introduced into zeolites which can supplement nutrient deficient soils. Thus, zeolites along with increasing yield can also increase the nutrient and water use efficiency of drylands.

**Keywords:** Zeolites; Nutrient use efficiency; Water holding capacity; Soil conditioners; Crop yield

### Introduction

Zeolites were first introduced by a Swedish mineralogist, A.F. Cronstedt in 1756, with the discovery of the mineral Stilbite [1]. By heating with a blowpipe flames this mineral lost water. He called this mineral “zeolite” from the Greek ‘zeo’, to boil and ‘lithos’, stone [2]. Since their discovery, zeolites are one of the most abundant minerals on earth which have been recognized as a separate group of minerals. After the discovery of zeolites work was carried out on ion exchange, hydration and dehydration. Zeolites are aluminosilicate minerals which have a molecular sieve action due to their open channel network, and are composed of  $\text{TO}_4$  tetrahedra linked with oxygen sharing the negative charge created by the presence of  $\text{AlO}_2^-$  which is balanced by cations that neutralize the charge deficiency. These cations include: the alkaline ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Rb}^+$ ,  $\text{Cs}^+$ ), the alkaline earth ( $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ) cations,  $\text{NH}_4^+$ ,  $\text{H}_3\text{O}^+$  ( $\text{H}^+$ ),  $\text{TMA}^+$  (Tetramethylammonium) and other nitrogen containing organic cation, and the rare earth and noble metal ions [2]. After the discovery of zeolites, work was carried out to study their physical and chemical properties. In 1930 “Molecular sieve” was a term introduced for materials that exhibited selective adsorption properties [3]. These materials contain other elements in addition to, or in lieu of, silicon and aluminum. In the

**Citation:** Jakkula VS, Wani SP. Zeolites: Potential soil amendments for improving nutrient and water use efficiency and agriculture productivity. *Sci Revs Chem Commun.* 2018;8(1):119.

© 2018 Trade Science Inc.

past limited classes of zeolites were known, however work by researchers on different classes of zeolites paved way for silicates, metallosilicates, metalloaluminates,  $AlPO_4$ s and silico-and metalloaluminophosphates.

It is simple to compare adsorptive and catalytically properties of zeolites by describing them in terms of their pore openings and channel systems. A need to relate and compare structural properties of a large number of zeolite structures has led to development of secondary building units (SBU). These units are also used in efforts to understand formation of individual structures from complex mixtures used in the synthesis [2]. As the individual tetrahedral  $TO_4$  unit is the basic building unit of a zeolite, a secondary building unit (SBU) consists of selected geometrical groupings of those tetrahedrons. Pore size and channel system play a vital role within a zeolite framework. Instead of visualizing zeolites in terms of interconnection of voids by these 8, 10, and 12 ring pore openings; the structure can be viewed in terms of one, two or three dimensional tubes or channels. For some zeolites “channels” are very short, similar to portholes or windows connecting large cavities within the structure [2].

This review is an overview of application of zeolites in various fields with emphasis on their use in agriculture as effective soil conditioners, their influence on cation exchange capacity, their soil nutrient interactions and their impact on soil properties, nutrient and water use efficiency and crop yield. Most of the research carried out on zeolites relates to irrigated and temperate conditions. It is therefore envisaged that through this review an understanding of zeolites for drylands and semi-arid tropical climatic conditions is established, which will then pave way for synthesizing these materials as slow/controlled release fertilizers to minimize the use of urea in developing countries and also as effective macro/micro nutrient carriers for enriching degraded soils of drylands.

## **Applications of Zeolites**

Zeolites have numerous applications i.e. in catalysis, in gas adsorption, industrial gas separation, water treatment (wastewater and drinking water), agriculture, and metal immobilization in soils, ion exchange, aquaculture, odour control, and desiccation and as phosphate substitutes in detergents. Zeolites have various applications, based on their cost and ion exchange behavior the main areas where zeolites are widely used are in detergents; in ammonia/ammonium removal from wastewater effluent; in radioactive isotope removal from spent pile effluent and in agriculture. By far the most important of these is in detergents, where zeolites are employed as water softeners, partially replacing tri-polyphosphate builders [4]. The annual turnover of Zeolite A for this purpose alone in the U.S is several billions of dollars. The adsorption capacity of various zeolites i.e. Mordenite, Chabazite, Clinoptilolite, Ferrierite and Phillipsite, have been tested for ammonia over methane in the gasification procedure and demonstrated that Phillipsite would not adsorb  $CO$ ,  $CO_2$  and  $CH_4$  in a gasification stream [5] but at  $25^{\circ}C$ , Phillipsite adsorbs  $NH_3$  rapidly. Synthetic zeolites are potentially useful additives to bind heavy metals, study by Lewis et al. [6] showed that out of the many zeolites i.e. Zeolite A, Mordenite, Faujasite, Zeolite X, Zeolite P; Zeolite A has the highest capacity to bind heavy metals such as cadmium and zinc. Zeolite A has the highest binding capacity between pH 5 and 6.5 and was stable  $pH > 5.5$  [6].

Many authors have studied the possibilities of separation of ammonium ions from drinking water and from wastewaters by zeolites [7-9]. The application of synthetic Phillipsite prepared by hydrothermal alteration of power plant

ashes, when compared to a natural Slovakian Clinoptilolite, showed that the former significantly reduced the concentration of ammonium ions in wastewaters [10]. Various systems are proposed for removal of pollutants including heavy metals and radioactive ions from industrial effluents, in Denver, Colorado (USA) natural zeolites are used for removal of ammonium in potable water systems.  $\text{Na}^+$  present in wastewater can also be removed using natural zeolites, thereby allowing this water to be used for agricultural applications. In a study by Santiago et al. [11], it was observed that only 14.3% of the exchangeable  $\text{Na}^+$  adsorption capacity i.e. 22.04 mEq/100 g  $\text{Na}^+$  was for natural zeolite. In order to increase the adsorption capacity, zeolites were treated with  $\text{NH}_4\text{C}_2\text{H}_3\text{O}_2$  at 1 M concentration which increased the adsorption capacity to 41.08 mEq/g. This increase represents 26.6 % of the theoretical exchangeable capacity of zeolite material. Since synthetic zeolites were first used commercially, they have found many uses in adsorbent and catalytic applications based on their unique physical structure. As adsorbents, zeolites have wider applications in chemical, petroleum and natural gas operations. They are used to dry and purify gaseous and liquid streams along with wide use in drying and purifying both gaseous and liquid streams along with separating olefins from paraffin's, oxygen from air and Para xylene from mixed xylenes.

Work has been carried out in introducing metals into zeolite framework, which serves as both oxidation and reduction catalysts due to the chemical reactions that take place within the internal cavities. In principle the high internal surface area allows high reactivity and the cation exchange capacity of zeolites permits facile introduction of acidic or transitional metal catalytic function [12]. Some examples of catalysis uses are: titanium ZSM-5 in the production of caprolactum, and copper zeolites in  $\text{NO}_x$  decomposition. Zeolites are widely used in catalysis and are improved heterogeneous catalysts as compared to amorphous catalysts. Because of the shape selectivity zeolites are used in cracking and alkylation processes i.e. in hydrocracking, toluene alkylation, and methanol dehydration. Due to the ban on production and use of chlorofluorocarbons (CFCs) and Halons, which resulted in ozone depletion, there is a greater need for safe disposal of these banned materials and/or their conversion into alternative materials, which are environmentally acceptable. Development of zeolite-based catalysts can act as potential agents for dehalogenation processes for environmental protection [12].

In agriculture zeolites have numerous applications, as slow release fertilizers, as heavy metal removers, as soil conditioners, and as increasing the nutrient and water use efficiency along with increasing crop yield. Soils throughout the world have been contaminated with heavy metals (i.e. Cu, Cd, Pb and Zn) and with radionuclides (i.e.,  $^{134}\text{Cs}$ ,  $^{137}\text{Cs}$  and  $^{90}\text{Sr}$ ). Studies have shown that zeolites are widely used in heavy metal uptake. There was a significant decrease in metal uptake by plants grown in zeolite amended soils [13]. In barren soils, contaminated with heavy metals zeolite addition completely eliminated metal phytotoxicity and allowed establishment of vegetation [14,15]. The simple concept of use of zeolites in metal contaminated soils shows that soils get alkalized and may help to reduce metal mobility and phytotoxicity. All of these depend upon zeolites ion exchange capability.

### **Use of zeolites in agriculture**

Nitrate contamination of both underground and surface waters results from nitrogen loss from irrigated cropland, particularly sandy soils. Increase in crop yield and decrease in nitrate pollution of water systems can be achieved by addition of zeolites (both natural and synthetic) as a fertilizer amendment to soil, thereby saving the cost for future improvement. Zeolites have been used since 1960s, with the use of natural zeolites for plant growth being first reported in Japan [16]. Soil

macro nutrients i.e. nitrogen, phosphorus and potassium could be supplied to plants in the form of ammonium ( $\text{NH}_4^+$ ) and Phosphate (P) exchanged zeolites [17-19]. The zeolites can act as a macro nutrient supplement. One of the most important applications of zeolites in agriculture is the slow/controlled-release fertilizer aspect. Slow release is a term that is interchangeable with delayed-release, controlled-release, controlled-availability, slow acting and metered-release [20]. The widespread abundance Clinoptilolite, Chabazite, Phillipsite and Mordenite in nature and their selectivity for certain cations (i.e.  $\text{NH}_4^+$  and  $\text{K}^+$ ) makes them suitable for slow-release fertilizer aspects.

Study by Jakkula et al. [21] has shown that Linde type F is one such zeolite which has a very high affinity towards  $\text{NH}_4^+$ . The synthesized Linde type F (LTF) which is a synthetic form of edingtonite (EDI) was ion exchanged with  $\text{NH}_4\text{NO}_3$  and introduced as a soil amendment to look at growth of maize plant (*Zea mays*). The selectivity and affinity of Linde type F for  $\text{NH}_4^+$  was higher, which resulted in lower plant growth. In a study to assess the Cd uptake of two test crops wheat and spinach, Puschenreiter et al. [22] found that slow-release synthetic zeolite-bound Zinc and Copper fertilizers may help to reduce Cd accumulation in edible parts of crops. Most of the previous studies carried out on the slow/controlled release fertilizer aspects of zeolites were focused on natural Clinoptilolite and to a certain extent on natural Phillipsite. Zeolites when applied with conventional fertilizers aids in an increase in seed germination and an overall increase in yield of spinach (*Spinaciaoleracea*) [23]. Zeolite also have an effect on yields of plums (*Prunuspersica*) and vines (*Vitisvinifera*), in a study in southern Italy by Burriesci et al. [24] it was found that on adding zeolites to traditional fertilizers there is an increase in fruit size and yield. Jordanian zeolite tuffs have been used for various applications. Studies on strawberries (*Fragariaoxananassa*) [25] showed by adding Jordanian Chabazite-Phillipsite tuff-soil mixture in 2:5 ratio increased crop yield ( $\leq 72\%$ ) and plant height and decreased mortality. Although zeolites have been used since ages still there is huge scope for these materials to penetrate into agriculture as slow release fertilizers, soil conditioners and drought proof materials. Research done in the past focused on natural zeolites, current research is looking at both natural and synthetic zeolites for improving soil properties and increasing crop yield. The main focus of research is to develop zeolites which are eco-friendly, cost effective and potential fertilizer substitutes for increasing crop yield and reducing nitrate leaching.

### **Zeolites as effective soil conditioners**

Zeolites are widely used as soil conditioners to improve soil physio-chemical properties [26]. The increase in soil pH and exchangeable K on adding Clinoptilolite to soils is significant, and has no influence on soil humus content and soil chemical composition [27]. Addition of zeolites to soil helps to control soil pH and improve ammonium retention. Out of the many properties of zeolites some of the prominent ones include increase in soil CEC, act as a reservoir of  $\text{K}^+$  [28] and increase in the water-holding capacity of loamy sand soils [29]. Studies on synthetic Phillipsite and natural Clinoptilolite supplemented soils showed an increase in water holding capacity and CEC [26]. Natural zeolite-enriched soils increased water holding capacity by (18-19%) and CEC by 30-40%, whereas the synthetic zeolite enriched soils increased water-holding capacity by 25-30% and CEC by 40-50%. Zeolites once saturated with  $\text{NH}_4^+$  can be used as effective fertilizers. Ion exchange plays an important role in introducing  $\text{NH}_4^+$  into the framework, by exchanging for cations present in the zeolite framework. The use of conventional fertilizers used intensively by farmers for increased crop production can be replaced by ammonium exchanged zeolites which are valuable source for controlled release of N. Soils low in K can be replaced with, zeolites such as Linde type F synthesized in K form which can be used for replenishing K in these soils.

Increase of soil pollution by heavy metals is one of the main problems of modern agriculture. The increased accumulation of metals in soils is highly influenced by human activity and the related development of industry and irrational use of fertilizers. Heavy metals may persist in the soil for hundreds of years, which is associated with the risk of their inclusion in a trophic chain. The main factor determining the solubility of heavy metals in soil is low pH [30]. Another important factor that determines the retention of heavy metal ions is the sorption capacity of soils, which is determined by the number and quality of soil sorption complex. Natural and synthetic zeolites are being used among other additives to reduce the bioavailability of heavy metals in the soil [16]. Ion exchanged zeolites can be used as carriers of nutrient elements in fertilizers, and can be exploited to trap undesirable metals and prevent their uptake into the food chain. The transfer of fertilizer-added heavy metals, such as copper, cadmium, lead, and zinc, from soils to plants can be reduced by application of pulverized zeolites [31].

The use of zeolites in acidic soils causes an increase in pH that significantly affects reduction in heavy metals solubility and bioavailability for plants. Except of ion exchange reactions, the pH increase promotes the adsorption of heavy metals on the surface of zeolites and their oxides precipitation [32]. The use of zeolites significantly reduced the uptake of cadmium and lead in wheat [33]. Studies concerning the effects of various additives on a change of the solubility of lead, cadmium and zinc have shown that zeolites significantly reduced the solubility of lead and cadmium in polluted soils. Zeolites from Italy and New Zealand have been used for ammonium removal from wastewater [8,33]. After researching Turkish zeolites studies confirmed that nitrification decreased after zeolite application to soil [34]. The protection of groundwater at waste disposal sites can be achieved by using natural zeolites [4]. Another important use for natural zeolites is reduction in soil contamination risk associated with heavy metals [35] and decrease in the heavy metal content of plants [36].

### **Use of clinoptilolite in agriculture**

The pronounced selectivity of Clinoptilolite for  $\text{NH}_4^+$  and  $\text{K}^+$  was exploited in Japan as slow-release chemical fertilizers. The high cation exchange property of Clinoptilolite towards  $\text{NH}_4^+$  makes it particularly valuable when applied to finely texture sandy soils to restrict leaching of  $\text{NH}_4^+$ , as high retention of this ion would control its accessibility to nitrifying bacteria [37]. Other studies showed that an organo-zeolitic substrate incorporated in a composted mixture of poultry manure and Clinoptilolite-zeolite tuff is an effective fertilizer and soil conditioner [38]. Some of the soil properties which can be improved by incorporating zeolites as part of organic fertilizers are soil quality by improving soil structure, ion exchange capacity of soils, and moisture and nutrient retention. The nutrient loss to the environment can also be inhibited by application of zeolites to soil. Studies showed that nitrogen can be supplied to plants from soils amended with ion exchange form of  $\text{NH}_4^+$ - saturated Clinoptilolite [6,39]. A substantial reduction of  $\text{NO}_3^-$  leaching and increased uptake of N-fertilizer was observed on addition of 10% (w/w) Clinoptilolite to sand used in the construction of golf-course greens [40]. Studies on sweet corn (*Zea mays*) showed that amending soils with ammonium-Clinoptilolite [41], N leaching reduced considerably, while sustaining plant growth and increasing N-use efficiency, when compared to ammonium sulphate amended soils. Also sweet corn grown in soil amended with ammonium-Clinoptilolite assimilated significantly more N than sweet corn grown in soil amended with ammonium sulphate. Researchers developed a kinetics theory that, controlled release of P,  $\text{NH}_4^+$  and  $\text{K}^+$  can be accomplished in synthetic soil through dissolution and cation exchange reactions between phosphate rock and

Clinoptilolite [18]. On the other hand, mixing  $\text{NH}_4^+$ -saturated Clinoptilolite with phosphate rock can act as a supplement of N and P to plants [42,43].

### **Use of phillipsite in agriculture**

The exchange properties of natural Phillipsite tuff obtained from the Aritain area of Jordan were evaluated by studying the exchange properties of this zeolite in the  $\text{NH}_4^+\text{-Na}^+$  system. The tuff has a higher selectivity for ammonium ions than sodium ions. Studies on tuff revealed that release of  $\text{NH}_4^+$  from the saturated tuff occurs over a long period of time (630 days) [44]. These properties are favorable for using Jordanian Phillipsite tuff as a potential ammonium-based slow release fertilizer. Studies on the supply rate of P from the zeolitic fertilizer (saturated Phillipsite tuff from Tenerife) compared with  $\text{KH}_2\text{PO}_4$  alone indicated that zeolite fertilizers supply available P after 70 days of continuous percolation, whereas P from  $\text{KH}_2\text{PO}_4$  is exhausted after 50 days. Zeolitic fertilizers supplied available K throughout the whole experiment period [45]. Jakkula et al. [46] carried out research to study the effect of both synthetic zeolite Phillipsite and its natural counterpart on the high selectivity and affinity of these zeolites towards ammonium ( $\text{NH}_4^+$ ) and its potential as a slow release fertilizer. Results were promising for synthetic Phillipsite which showed predominantly higher affinity towards  $\text{NH}_4^+$ . The exchange capacity of Phillipsite is around  $3.5\text{-}3.6 \text{ meq.g}^{-1}$ , which makes Phillipsite suitable for agriculture due to the presence of high levels of potassium among exchangeable cations. The use of  $\text{NH}_4^+$ -Phillipsite tuff offers an alternative option to the widely used soluble  $\text{NH}_4$  fertilizers in agriculture.

Study was carried out in UK on synthetic and natural Phillipsite to study their effectiveness as slow release fertilizers [46]. Na-K form of Phillipsite (PHI) was synthesized and  $\text{NH}_4^+$  was introduced into the framework. Ion exchanged synthetic zeolite was compared with its natural counterpart. In a glasshouse experiment, maize was grown in zeolite amended at different loadings. A control comprising NPK fertilizer added to soil was used for comparison. Zeolite mineral Phillipsite amended soils at various growth stages revealed the affinity of this zeolite for  $\text{NH}_4^+$ . Results demonstrated that natural zeolite Phillipsite has lower affinity towards  $\text{NH}_4^+$  on comparison to synthetic Phillipsite when introduced as a soil amendment. The ion exchange interactions in soil were promising for synthetic zeolite Phillipsite, with cations in soil exchanging more freely with  $\text{K}^+$  and  $\text{NH}_4^+$  present in the synthetic Phillipsite framework [47].

### **Zeoponic substrates**

Co-operative programmes between NASA and industry have developed zeoponic plant growth media and fertility systems. A zeoponic plant growth system is defined as cultivation of plants in artificial soils, which contain zeolites as a major component [20,48]. Zeoponic substrates have been used on the Russian Mir Space Station and U.S. Space Shuttles to grow various plants, including radishes, wheat and brassicas. Space gravitational biologists have considered Zeoponic substrates as preferred material for plant growth in a microgravity environment. However, the effectiveness of zeoponic substrates for plant growth in macro gravity is unclear. Between 1994-1997, Zeoponix, Inc (USA) and NASA, conducted experiments on zeoponic materials as soil amendments for plant growth, studying seed germination, root development, fruit development, fruit quality and yield. Work was carried out on horticultural species and cereal crops. According to results claimed virtually all plants tested had improved yield or quality of growth when growing in soil-less media containing 5-30%

by volume zeoponic fertilizer-amendments [49]. With rising world populations, potential improvements in crop yield indicated with zeoponic systems are of particular importance in meeting rising food demands.

### **Zeolite nutrient interactions in soil**

Studies by Pickering et al. [50] showed that zeolites improve the use of  $\text{NH}_4^+$ -N and  $\text{NO}_3^-$ -N, reducing leaching losses of exchangeable cations along with nutrient use efficiency by increasing P availability from phosphate rocks. To improve plant yield, zeolites which have a high affinity for nutrients, may be used as growth media [38]. Zeolite attracts many trace elements and macro and secondary nutrients i.e. ammonium, potassium, calcium and magnesium. In order to balance the negative charge, zeolite attracts calcium from phosphorus mineral apatite such as rock phosphate or locked up phosphorus in soil. Zeolites have affinity for majority of cation in particular ammonium and potassium. It is known that plants extract ammonium, potassium and other nutrients by active uptake from roots. When cation such as ammonium or potassium source become available either naturally or applied, zeolites get recharged and prevent free nutrients from leaching.

Zeolites are being investigated for use as fertilizer enhancers and other agricultural amendments because of their unique cation-exchange properties. Zeolites serve as a slow release source of nutrients upon precharged with ammonium-nitrogen ( $\text{NH}_4$ -N), potassium ( $\text{K}^+$ ), or iron ( $\text{Fe}^{+2}$ ) when such precharged zeolites are used as a component of soilless media [51]. Studies on controlled release fertilizers have shown zeolite materials to have  $\text{NH}_4$  adsorption kinetics, natural zeolites have also shown  $\text{NH}_4$  adsorbent properties and controlled-release fertilizer properties [52]. Another use of zeolites is in P fertilization [43], ion exchanged zeolites i.e. ammonium exchanged zeolites increases solubility of soil P, enhances its uptake by the plants, and decreases its precipitation and fixation by exchanging  $\text{NH}_4^+$  with other cations [43-45]. In addition, zeolitic tuff can be used in saline soils or when saline water is used for irrigation. In a study on absorption of four  $\text{Na}^+$  salts [sodium chloride ( $\text{NaCl}$ ), sodium bicarbonate ( $\text{NaHCO}_3$ ), sodium carbonate ( $\text{Na}_2\text{CO}_3$ ), and sodium sulfate ( $\text{Na}_2\text{SO}_4$ )] by a natural zeolite (clinoptilolite) from China, it was shown that the zeolite strongly absorbed  $\text{Na}^+$  and anions and that the rate of absorption increased with time as the salt concentration increased [53]. It was concluded in that study that zeolites might be used as amendments for sodic saline alkaline soils. Urea-N losses could be minimized using zeolites as additives in the fertilizers to control the retention and release of  $\text{NH}_4^+$  and convert it as an enhanced efficiency fertilizer (EEF). The fact that zeolite can relatively strongly fix the ammonia nitrogen leads to the reduction of nitrogen leaching.

Among the various zeolites assessed for high cation exchange capacity, clinoptilolite allows for entrapping or releasing various cations [54]. This zeolite can be used to control N loss from urea because of the small molecular size of the open-ringed structure which physically protects  $\text{NH}_4^+$  ions against microbial nitrification [55]. Excessive use of fertilizers results in environmental pollution, studies by McGilloway et al. [3] have shown that adoption of zeolite i.e. clinoptilolite decreases the excessive and unbalanced use of N fertilizers in agriculture by maximizing N use efficiency and water use efficiency. Studies have shown that N use efficiency can be improved by the use of clinoptilolite zeolite and N fertilizers in conjunction with each other [2,56-58]. The increased efficiency of N utilization when urea is used together with clinoptilolite zeolite has been demonstrated by a number of researchers [59-61]. Addition of zeolite to soil reduced soil urease activity through urease adsorption on the zeolite [62] and the prediction equation followed exponential function  $Y = 40.975e^{-3.6848x}$ ,

where Y is the soil urease activity and x is the soil: zeolite ratio,  $R^2=0.95$  [63]. Studies by He et al. [7] showed that natural zeolite clinoptilolite amended soil increases soil surface area and cation exchange capacity which aids in lowering nitrogen concentration in the leachate and increasing moisture and nutrients in the soil.

### **Impact of zeolites on soil properties, nutrient and water use efficiency and crop yield**

Zeolite assists water infiltration and retention in the soil and acts as a natural wetting agent due to its very porous properties and the capillary suction it exerts. In order to assist water distribution through soils, zeolites act as excellent amendment for non-wetting sands [64]. In order to achieve less fertilizer for the same yield or the same amount of fertilizer lasting longer and producing higher yields, zeolites can be used which hold nutrients in the root zone for plants to use when required which leads to more efficient use of N and K fertilizers [65,66]. Zeolites are effective carriers of herbicides, fungicides, and pesticides due to their high adsorption capacities in the dehydrated state and the high ion-exchange capacity. Zeolites can be produced from various sources and they help in absorption and retention of plant nutrients and water and supplemented micronutrients, the synthetic zeolite produced from coal ash is a beneficial soil amendment which aids in the above properties [23].

Zeolites sorption capability and its ability towards uniform release of nutrients into the soil along with its ion-exchange ability prevents against their quick elution [67]. The controlled release unique properties of zeolites allow a gradual and controlled introduction of necessary nutrients i.e. potassium, ammonium or phosphates into the soil. Moreover, zeolites modified with Cu (II) ions can be a good source for Cu deficient soils. Zeolites have the ability to be retained in the soil for several years, therefore the stabilization of ionic exchange balance in soil for several minutes or hours can be considered as fast. The redundant ammonium ions would bind to zeolite in soil and would be gradually release into the soil solution; this will help the liquid nitrogen-based plant nutrition mechanisms.

In semi-arid India, on tropical soils in order to increase crop yield and quality, liming and balanced nutrient supply are essential. Best management practices involve balanced nutrition whereby fertilizers need to be applied as per the need of crop, excessive use of fertilizers (urea in particular) results in nitrate leaching and losses by  $\text{NH}_3$  volatilization. In order to minimize the nitrate leaching and  $\text{NH}_3$  volatilization slow/controlled release fertilizers in the form of zeolites are need of the hour to support the increasing N demand and avoid the low N use efficiency (NUE). Use of aluminosilicates (zeolites) with nitrogenous fertilizer materials can reduce the nitrate leaching [68]. Zeolites used with urea mineral fertilizer can enhance the efficiency of this source by improving the nitrogen use through the control of retention of ammonium ion, contributing to increased N uptake and crop yields. In order to decrease the environmental impact of agriculture, zeolite utilization which maximizes the N use efficiency along with water use efficiency are key as zeolite aids in increasing water holding capacity [7], which helps in reduction of soil irrigation, because water is well retained within zeolite's structure. The other properties of zeolites are reduction of soil acidity [69,70] and reduction of temperature oscillations. It has been reported that natural zeolites not only augment water holding capacity and soil drainage [71,72] but also increase crop water use efficiency [73].

Zeolites not only improve soil conditions but also help in increasing crop yields, in the past studies carried out on various soil types and various crops i.e., potato, maize, rice, tomato, eggplant, carrot showed crop yield increase and in some



cases yield increase reached up to 60% [74]. Due to the zeolites exceptionally high ion-exchange capacity along with vegetable crops, zeolites can also be successfully used in cultivating different cereals, forage crops, vine and fruit crops [75]. In order to improve the horizontal spread of water after irrigation zeolites play an important role as their porous structure assures a permanent water reservoir in the root zone [76,77]. Zeolites make available nutrients exactly when needed thereby preventing unnecessary losses of nutrients [78]. Zeolites being source of nutrients act as excellent carriers, stabilizers and regulators of mineral fertilizers [79]. Zeolites decrease application rate of N and K fertilizers, as they themselves are carriers of N and K fertilizers thereby increasing efficacy [77].

## Conclusion

Zeolites (a separate group of minerals found on earth in abundance and also could be synthesized) find a large number of potential applications in agriculture, particularly in soil management as well as wastewater treatment and heavy metal pollution removal. Nitrogen released as  $\text{NH}_4$  after dissolution of applied chemical fertilizers or decomposition of manures and cover crops is adsorbed by zeolite. They can be used either as carriers of nutrients and/or medium to free nutrients to promote nutrient use efficiency. Zeolites are effectively used as soil ameliorants for treating salinity and have a positive effect on soil fertility, sorption properties of soil and subsequently the formation of phytomass. Also zeolites improve the efficiency of water and nutrient use of plants and decrease runoff and sediments amount by increasing the soil water holding capacity. Acting as slowly soluble fertilizers, they improve water balance and sorption characteristics of light sandy soils in particular, which is reflected in higher yield and better quality. Studies have shown that zeolites significantly adsorb  $\text{NH}_4$  and P and minimize reactive  $\text{NO}_3$  formation and also could be effectively used for wastewater treatment as well as cleaning of heavy metal polluted sites. Usage of zeolites increased plant growth in many research studies across the globe, and also use of zeolite is expected to reduce a third to half Urea-N fertilization. Soil fertilized or amended with organo-zeolite mixtures regulate N release and minimize reactive N ( $\text{NO}_3^-$ ) formation. Research on efficient utilization of zeolites for rainfed dryland agriculture is critical in the current climate change scenario. It is hoped that continued research in enhanced ammonia supply by zeolite as carrier will result in better nitrogen fertilization management through strategies best-suited to support vegetative development of tested crops in drylands. As rainfed agriculture experiences drought year after year, use of zeolites which increases the water holding capacity of various soil types could be alternatives to conventional fertilizers. Finally, zeolites can serve as future environmentally friendly materials for both increasing crop yield and reducing agriculture input costs.

## REFERENCES

1. Nguyen ML, Tanner CC. Ammonium removal from wastewater using natural New Zealand zeolites. *New Zeal J Agric Res.* 1998;41:427-46.
2. Gruener JE, Ming DW, Henderson KE, et al. Common ion effects in zeoponic substrates: Wheat plant growth experiment. *Micropor Mesopor Mat.* 2003;61:223-30.
3. McGilloway R, Weaver R, Ming D, et al. Nitrification in a zeoponic substrate. *Plant Soil.* 2003;256:371-8.

4. Fronczyk J, Garbulewski K. Evaluation of zeolite-sand mixtures as reactive materials protecting groundwater at waste disposal sites. *J Environ Sci.* 2013;25:1764-72.
5. He ZL, Calvert DV, Li AK, et al. Clinoptilolite zeolite and cellulose amendments to reduce ammonia volatilization in a calcareous sandy soil. *Plant Soil.* 2002;247:253-60.
6. Lewis MD, Moore FD, Goldsberry KL. Ammonium exchanged Clinoptilolite and granulated Clinoptilolite with urea as nitrogen fertilizers. *Zeo-Agriculture.* Boulder, Colorado. Westview Press. 2003;105-11.
7. Hatfield JL, Prueger JH. Nitrogen over-use, under-use, and efficiency. "New directions for a diverse planet." Proceedings of the 4<sup>th</sup> International Crop Science Congress. Brisbane, Australia. 2004.
8. Ciambelli P, Corbo P, Liberti L, et al. Ammonium recovery from urban sewage by natural zeolites. *Akademiai Kiado, Budapest, Hungary.* 1988;501-10.
9. Colella C, Aiello R. Ammonium removal from tannery sewages by selective ion exchange using natural Phillipsite. *Akademiai Kiado, Budapest, Hungary.* 1988;491-500.
10. Kovanda F, Kolousek D, Seidl V, et al. Utilization of synthetic Phillipsite as a means for decreasing the concentration of ammonium ions in waste waters. *Ceram Silikaty.* 1994;38:75-9.
11. Santiago O, Walsh K, Kele B, et al. Novel pre-treatment of zeolite materials for the removal of sodium ions: potential materials for coal seam gas co-produced wastewater. *Springer Plus.* 2016;5:1-16.
12. Howe RF. Zeolite catalysts for dehalogenation process. *Appl Catal A Gen.* 2004;271:3-11.
13. Gworek B. Inactivation of cadmium in contaminated soils using synthetic zeolites. *Environ. Pollut.* 1992;75:269-71.
14. Rebedea I, Lepp NW. The use of synthetic zeolites to reduce plant metal uptake and phytotoxicity in two polluted soils. *EnvironGeochem. Hlth.* 1995;16:81-8.
15. Vangronsveld J, Sterckx J, Van Assche F, et al. Rehabilitation studies on an old non-ferrous waste dumping ground: effects of vegetation and metal immobilisation by beringite. *J Geochem Explor.* 1995;52:221-9.
16. Minato H. Characteristics and uses of natural zeolites. *Koatsugasu.* 1968;5:536-47.
17. Allen ER, Hossner LR, Ming DW, et al. Release rates of phosphorus, ammonium, and potassium in clinoptilolite-phosphate rock systems. *Soil Sci Soc Am J.* 1996;60:1467-72.
18. Allen ER, Ming DW, Hossner LR, et al. Modeling transport kinetics in clinoptilolite-phosphate rock systems. *Soil Sci Soc Am J.* 1995;59:248-55.
19. Allen ER, Ming DW, Hossner LR, et al. Growth and nutrient uptake of wheat in clinoptilolite-phosphate rock substrates. *Agron J.* 1995;87:1052-9.

20. Ming DW, Allen ER. Zeoponic substrates for space applications: Advances in the use of natural zeolites for plant growth. *Natural Microporous Materials in Environmental Technology*. Misaelides, P. (Ed). Kluwer Academic Publishers, Netherlands. 1999;157-76.
21. Jakkula VS, Williams CD, Hocking TJ, et al. High selectivity and affinity of Linde type F towards  $\text{NH}_4^+$  on application as a soil amendment for maize growth. *Micropor Mesopor Mat*. 2006;88:101-04.
22. Puschenreiter M, Horak O. Slow-release zeolite-bound zinc and copper fertilizers affect cadmium concentration in wheat and spinach. *Commun Soil Sci Plan*. 2003;34:31-40.
23. Burriesci N, Valente S, Ottana R, et al. Utilization of zeolites in spinach growing. *Zeolites*. 1983;4:5-8.
24. Burriesci N, Valente S, Zipelli C, et al. Studies on zeolites in agriculture. Effect of crop growth of *Prunus persica* and *Vitis vinifera*. *Zeolites*. 1984;4:373-6.
25. Ibrahim KM, Ghir AM, Khoury HN. Mineralogy and chemistry of Jordanian Chabazite-Phillipsite tuff. *Clay Miner*. 1996;39:47-55.
26. Kralova M, Hrozinkova A, Ruzek P, et al. Synthetic and natural zeolites affecting the physical-chemical soil properties. *Rostlinna Vyroba*. 1992;42:131-41.
27. Filcheva EG, Tsadilas CD. Influence of Clinoptilolite and compost on soil properties. *Commun Soil Sci Plan*. 2002;33:95-607.
28. Hershey DR, Paul JL, Carlson RM. Evaluation of potassium enriched Clinoptilolite as a potassium source of potting media. *HortScience*. 1980;15:87-9.
29. Huang ZT, Petrovic AM. Clinoptilolite zeolite influence on nitrate leaching and nitrogen use-efficiency in simulated sand-based golf greens. *J Environ Qual*. 1994;23:1190-4.
30. Karczewska A. Heavy metals in soils polluted by emissions from copper smelters-forms and solubility. *Z Nauk. AR. Wroclaw*. 2002;432:1-159.
31. Fugii S. Heavy Metal Adsorption by Pulverized Zeolites. Japan, Kokai. 1974.
32. Shia W, Shaoa H, Li H, et al. Progress in the remediation of hazardous heavy metal-polluted soils by natural zeolite. *J Hazard Mater*. 2009;170:1-6.
33. Chen ZS, Lee GJ, Liu JC. The effects of chemical remediation treatments on the extractability and speciation of cadmium and lead in contaminated soils. *Chemosphere*. 2000;41:235-42
34. Isildar AA. Effect of addition of zeolite to the soil on nitrification. *Turk J Agric For*. 1994;23:363-8.
35. Michalik J, Missik J. Sorption properties of zeolites from Nizny Hrabovec. *Acta Fytotechnica*. 1995;50:131-41.

36. Petkova D, Popandova S, Popova T, et al. Testing of zeolite-glaucanite-phosphorite mixtures for decrease of lead uptake by plants. *Bulg J Agric Sci.* 2000;6:321-6.
37. Andronikashvili TG, Kadava MA, Gamisonia MK. Effect of natural zeolites on microbe landscape of some soils in Georgia. *Sofia Int Zeolite Meet.* 1995;4:111-2.
38. Leggo PJ. An investigation of plant growth in an organo-zeolitic substrate and its ecological significance. *Plant Soil.* 2000;219:135-46.
39. Pirela HJ, Westfall DJ, Barbarick KA. Use of Clinoptilolite in combination with nitrogen fertilization to increase plant growth. Boulder, Colorado, Westview Press. 1984;113-22.
40. Mumpton FA. La roca magica: Uses of natural zeolites in agriculture and industry. *P Natl Acad Sci USA.* 1999;96:3463-70.
41. Perrin TS, Drost DT, Boettinger JL, et al. Ammonium loaded Phillipsite: A slow release Nitrogen fertilizer for sweet corn. *J Plant Nutr.* 1998;21:515-30.
42. Chesworth W, Van Straaten P, Sadura S. Solubility of apatite in clay and zeolite-bearing systems: applications to agriculture. *Appl Clay Sci.* 1987;2:291-7.
43. Barbarick KA, Lai TM, Eberl DD. Exchange fertilizer (phosphate rock plus ammonium-zeolite) effects on sorghum-Sudan grass. *Soil Sci Soc Am J.* 1990;54:911-6.
44. Dwairi IM. Evaluation of Jordanian zeolite tuff as a controlled slow-release fertilizer for  $\text{NH}_4^+$ . *Environ Geol.* 1998;34:1-4.
45. Notario del Pino IJ. Phosphorus and potassium release from phillipsite-based slow-release fertilizers. *J Control Release.* 1995;34:25-9.
46. Jakkula VS, Williams CD, Hocking TJ, et al. High selectivity and affinity of synthetic Phillipsite compared with natural Phillipsite towards ammonium ( $\text{NH}_4^+$ ) and its potential as a slow release fertilizer. *Arch Agro Soil Sci.* 2011;57:47-60.
47. Jakkula VS. Synthesis of zeolites and their application as soil amendments to increase crop yield and potentially act as controlled release fertilizers. PhD thesis. University of Wolverhampton. 2006;273.
48. Parham WE. Future perspectives for natural zeolites in agriculture and aquaculture. Westview Press, Boulder, Colorado. 1984;283-5.
49. Ming DW, Allen ER. Use of Natural Zeolites in Agronomy, Horticulture, and Environmental Soil Remediation. Blacksburg Press. Virginia. 2001;45:619-53.

50. Pickering HW, Menzies NW, Hunter MN. Zeolite/rock phosphate - a novel slow release phosphorus fertilizer for potted plant production. *Sci Hortic.* 2002;94:333-43.
51. Williams KA, Nelson PV. Using precharged zeolite as a source of potassium and phosphate in a soilless container medium during potted chrysanthemum production. *J Am Soc Hort Sci.* 1997;122:703-08.
52. Kithome M, Paul JW, Lavkulich LM, et al. Kinetics of ammonium adsorption and desorption by the natural zeolite clinoptilolite. *Soil Sci Soc Am J.* 1998;62:622-9.
53. Zhong XZ, Wu J. Studies of the absorbing mechanism of natural zeolite in the improvement of sodic saline-alkali soil. 1. The absorption of the main salts in the sodic saline-alkali soil by natural zeolite. *J Jilin Agric Univ.* 1990;12:40-5.
54. Kamarudin KS, Hamdan H, Mat H. Methane adsorption characteristic dependency on zeolite structures and properties. *Copthorne Orchid Hotel. Penang.* 2003;29-30.
55. Ferguson GA, Pepper IL. Ammonium retention in sand amended with clinoptilolite. *Soil Sci Soc Am J.* 1987;51:231-4.
56. Bouzo L, Lopez M, Villegas R. Use of natural zeolites to increase yields in sugarcane crop minimizing environmental pollution. *World congress of soil science.* 1994;15:695-701.
57. Rehakova M, Cuvanova S, Dzivak M, et al. Agricultural and agrochemical uses of natural zeolite of the clinoptilolite type. *Current Opinion Soil State Materias Sci.* 2004;8:397-404.
58. Ahmed OH, Aminuddin H, Husni MHA. Reducing ammonia loss from urea and improving soil-exchangeable ammonium retention through mixing triple superphosphate, humic acid and zeolite. *Soil Use Manage.* 2002;22:315-9.
59. MacKown C, Tucker T. Ammonium nitrogen movement in a coarse-textured soil amended with zeolite. *Soil Sci Soc Am J.* 1985;49:235-8.
60. Latifah O, Ahmed OH, Nik Muhamad AM. Reducing ammonia loss from urea and improving soil exchangeable ammonium and available nitrate in non-waterlogged soils through mixing zeolite and sago (*Metroxylon sagu*) wastewater. *Int J Phys Sci.* 2011;6:866-70.
61. Rabai KA, Haruna AO, Kasim S. Use of formulated Nitrogen, phosphorus and potassium compound fertilizer using clinoptilolite zeolite in Maize (*Zea mays L.*) cultivation. *Emirates J Food Agric.* 2013;25:713-22.
62. Ramesh K, Reddy KS, Reshmi I, et al. Nanoporous zeolites as soil urease adsorbers: An opportunity for increasing nitrogen use efficiency. 2010.

63. Ramesh K, Biswas AK, Somasundaram J, et al. Nanoporous zeolites in farming: Current status and issues ahead. *Curr Sci India*. 2010a;99:760-5.
64. Szerment J, Ambrozewich-Nita A, Kedziora K, et al. Use of zeolite in agriculture and environmental protection. A Short review. *UDC*. 2014;96:691.
65. Gamze Turan N. The effects of natural zeolite on salinity level of poultry litter compost. *Bioresource Technol*. 2007;99:2097-101.
66. Khodaei-Joghan A, Asilan KS, Zeolite influences on nitrate leaching, nitrogen-use efficiency, yield and yield components of canola in sandy soil. *Arch Agro Soil Sci*. 2012;58:1149-69.
67. Perry CC, Keeling-Tucker T. Model studies of the precipitation of silica in the presence of aluminum; implications for biology and industry. *J Inorg Biochem*. 2000;78:331-9.
68. Alberto CCB, Polidoro JC, Marisa BMM, et al. Enhancing nutrient use efficiency using zeolites minerals-A Review. *Adv Chem Eng Sci*. 2016;6:295-304.
69. Comin JJ, Barloy J, Bourrie G, et al. Differential effects of monomeric and polymeric aluminum on root growth and on the biomass production of root and shoot of corn in solution culture. *Eur J Agron*. 1999;11:115-22.
70. Badora A. Aluminum and manganese mobility in the soil. *Polish J Soil Sci*. 2001;34:1-8.
71. Bigelow CA, Bowman DC, Cassel DK. Physical properties of three sand size classes amended with inorganic materials or sphagnum peat moss for putting green root-zones. *Crop Sci*. 2004;44:900-07.
72. Colombani N, Giambastiani BMS, Mastrocicco M. Predicting salinization trends in a lowland coastal aquifer: Comacchio (Italy). *Water Resour Manag*. 2014;29:603-18.
73. Gholamhoseini M, Ghalavand A, Khodaei-Joghan A, et al. Zeolite-amended cattle manure effects on sunflower yield, seed quality, water use efficiency and nutrient leaching. *Soil Tillage Res*. 2013;126:193-202.
74. Torii K. Utilization of natural zeolites in Japan. Pergamon Press. Elmsford. New York, USA. 1978;441-50.
75. Butorac A, Filipan T, Basic F, et al. Crop response to the application of special natural amendments based on zeolite tuff. *Rostlinná Výroba*. 2002;48:118-24.
76. Treacy MJ, Higgins JB. Collection of Simulated XRD Powder Patterns for Zeolites. (4<sup>th</sup> Edtn). Elsevier. Amsterdam. 2001.
77. Polat E, Karaca M, Demir H, et al. Use of natural zeolite (clinoptilolite) in agriculture. *J Fruit Orn Plant Res*. 2004;12:183-9.
78. Podlesakova E, Kremer J, Bicoovsky K. Die Umtauschreaktion des Kaliums und Natriums am Bentonit von Braňany. *Zeitschrift für Pflanzenernährung Bodenkunde*. 1967;116:1-10.

79. Bagdasarov VR, Kazachenko AA, Rustambekov MK, et al. Prolonged-activity nitrogen-zeolite fertilizer, Russia. 2004.