



ASSESSMENT OF PHYTOCHEMICAL AND ANTI-MICROBIAL ACTIVITY OF *MORINGA OLEIFERA* LAM AGAINST PATHOGENIC BACTERIA

Y. S. BANGINWAR*, N. S. BHAJIPALE and S. T. INGLE^a

Department of Pharmaceutics (Microbiology), S. G. S. P. S, Institute of Pharmacy,
Kaulkhed, AKOLA (M.S.) INDIA

^aDepartment of Plant Pathology, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, AKOLA (M.S.) INDIA

ABSTRACT

Antibacterial activities of aqueous, ethanol, petroleum ether and acetone extract of pod of *Moringa oleifera* (Shevga) were studied. The aqueous leaf extract of *Moringa oleifera* showed prominent antibacterial activities against *Pseudomonas aeruginosa*, *Salmonella typhimurium* and *E. coli*., moderate activity against *Staphylococcus aureus*, *Salmonella typhi* and least activity against *Enterobacter aerogens* and *Proteus vulgaris*. The aqueous extract showed maximum the acetone and petroleum ether extract moderate and ethanol extract least antimicrobial activities. Phytochemical screening revealed the presence of glucosinolates, phenolic (flavonoids, anthocyanin, cinnamate, proanthocyanidin) leaves contain 4-(α -L-rhamnopyranosyloxy) benzylglucosinolate, kaempferol-3-glycoside sugar, D-Mannose and tannin. These antibacterial properties support its traditional use of leaves and pod in the treatment of enteric infection.

Key words: *Moringa oleifera*, *Pseudomonas aeruginosa*, *Salmonella typhimurium*, *E. coli*, *Staphylococcus aureus*, *Salmonella typhi*, *Enterobacter aerogens*, *Proteus vulgaris*, Infection.

INTRODUCTION

From time immemorial, men depend on plants as medicine. The plant kingdom represents a rich store house of organic compounds and many of which have been used for medicinal purposes and could serve as lead for the development of novel agent having good efficacy in various disorders.

In India, use of different parts of several vogue from ancient time and inherited traditionally. The root and bark of *Moringa oleifera* (Moringaceae) have been used in the

* Author for correspondence; E-mail: drbyogesh@gmail.com; Ph.: 0724-2452640; Mo.: +919822701865

indigenous system of medicine in India for the treatment of heart complaint, eye diseases¹. The leave and seed extracts of *Moringa oleifera* show antimicrobial activity against *pseudomonas* and *Staphylococcus aureus*². The presence of active principle flavonoids phenolic compound, glucosinotes, L-rhamnoside anisaldehyde were demonstarated in leaves³. The seed shows presence of phenylacetone⁴.

The enteric or diarrheal infection by water are major public health problems in developing countries and contribute to the death of 3 to 6.1 million children annually. Enteric bacteria comprised of *Salmonella spp.*, *Shigella spp.*, *Proteus spp.*, *Klebsiella spp.*, *E. coli*, *Pseudomonas spp.*, *Vibrio cholerae* and *Staphylococcus aureus*, which are major etiologic agents of sporadic and epidemic diarrhea both in children and adult⁵. W.H.O 1993 reported that 85% population rely mainly on traditional therapies involving the use of plant extracts or their active constituents⁶. The use of medicinal plant in India contributes significantly in primary health and it is interesting to determine, whether actual pharmacological uses are merely based on folkore.

EXPERIMENTAL

Materials and method

The plants of *Moringa oleifera* were collected from Akola city and authenticated by Pharmacognosy Department, SGSPS, Institute of Pharmacy, Akola.

The fresh leaves/pod of *Moringa oleifera* were washed with water and the 0.5% HgCl₂, sterile water and grinded into mortar with pestle. The 50 g fresh leaves/pod were suspended in the 200 mL water and various popular organic solvents and extracted in Soxhlet apparatus and vacuum dried.

Amount of dry recovered powder extracts are recorded (50 g of original leaves and pod). This dry mass of various extract *Moringa oleifera* served for further experimentation.

Bacterial culture

The standard pathogenic bacterial cultures were procured from P.G. Department of Microbiology, Sant Gadge Baba Amravati University, Amravati, and used in the present study (Table 1). The bacteria rejuvenated in Muller-Hinton broth (Hi-media) at 37°C for 18 hrs and then stocked at 40°C in Muller-Hinton Agar subculture prepared from the stock for bioassay. A loopful culture was inoculated in 10 mL of sterile nutrient broth at 37°C for 3-4

hos. Turbidity of culture was standardized to 10^5 CFU with the help of SPC and turbidometry.

Table 1: Bacterial culture used in study

Bacterial pathogen	MTC number
<i>Escherichia coli</i> (<i>E. coli</i>)	452
<i>Staphylococcus aureus</i> (<i>S. aureus</i>)	87
<i>Salmonella typhi</i> (<i>S. typhi</i>)	733
<i>Salmonella typhimurium</i> (<i>S. typhimurium</i>)	98
<i>Proteus vulgaris</i> (<i>P. vulgaris</i>)	426
<i>Pseudomonas aeruginosa</i> (<i>P. aeruginosa</i>)	424
<i>Enterobacter aerogens</i> (<i>E. aerogens</i>)	111

Agar disc diffusion antibiotic activities

For antibacterial properties, 0.1 ml bacterial suspension of 10^5 CFU mL⁻¹ was uniformly spread on Muller-Hinton plate to form lawn cultures. The aqueous, petroleum ether, acetone and ethanol extracts were prepared in their respective solvents in such a manner that ultimate amount (in dry form) in each disc came to 10 mg, 8 mg, 6 mg, 4 mg, and 2 mg. The blotting paper discs (8 mm diameter) were soaked in various diluted extracts, dried in oven at 50°C to remove the excess of solvent and tested for their antibacterial activity against pathogenic bacteria by disc diffusion technique. After incubation of 24 hours at 37°C, zone of inhibition of growth was measured in mm. Tetracycline 10 µg (Hi-Media disc) was used as positive control while disc soaked in various organic solvents and dried were placed on the lawns as negative control.

RESULTS AND DISCUSSION

During the past decades, traditional systems of medicine have become increasingly important in view of their safety. Several antibiotics are currently been used to treat a variety of infectious human diseases have limited antimicrobial spectrum and development of drug resistance in pathogen lead to serious ill effects. Plant extracts have broad spectrum antimicrobial activity, easily available, effective against human infection and also non-toxic to the human beings.

The present study was conducted to investigate antibacterial properties of pod of *Moringa oleifera*. Herbal medicine plays a fundamental role in tradition medicine in rural areas of India.

The amount of dry extracts recovered from various solvents are shown in Table 2. Antibacterial activities of aqueous, petroleum ether, acetone and ethanol extracts of leaves of *Moringa oleifera* were studied. The aqueous extract showed prominent antibacterial activities against *Pseudomonas aeruginosa*, *Salmonella typhimurium* and *Escherichia coli*, where as it was moderate against *Staphylococcus aureus*, *Salmonella typhi* and least active against *Enterobacter aerogens* and *Proteus vulgaris*.

Table 2: Average recovery of dry extract from 50 g of Pod

Type of extract	Amount (g)
Aqueous	34.67
Petroleum ether	28.12
Acetone	29.64
Ethanol	30.16

The petroleum ether extract showed prominent antibacterial activities against *Pseudomonas aeruginosa*, *Salmonella typhimurium* and *Staphylococcus aureus* where as it was moderate against *Salmonella typhi* and *Proteus vulgaris* and least against *Enterobacter aerogens*, *Pseudomonas aeruginosa* and *Escherichia coli*.

The acetone extract showed maximum activity against *Escherichia coli*, moderate against *Pseudomonas aeruginosa*, *Salmonella typhi* and *Proteus vulgaris* and least against *Salmonella typhimurium*, *Enterobacter aerogens* and *Staphylococcus aureus*.

The ethanol extract showed maximum activity against *Salmonella typhimurium*, *Salmonella typhi* and *Pseudomonas aeruginosa*, moderate against *Escherichia coli* and *Proteus vulgaris* and least against *Enterobacter aerogens* and *Staphylococcus aureus* (Table 3).

The aqueous extract showed maximum antibacterial activities against the tested bacterial pathogens due to higher solubility of antibacterial principle in aqueous as compared to organic solvents. Moreover, the pods are highly fibrous and it is difficult to ground it, and it may be difficult to extract the antibacterial principle by organic solvents.

The pods of *Moringa oleifera* contain various phytochemical components such as flavonoids, glycoside, phenolic compound, polysaccharide etc. (Table 4).

Table 4: Phytochemical analysis of *Moringa oleifera* pod

Phytochemical	Result
Alkaloid	Absent
Flavonoid	Present
Carbohydrate	Present
Glycoside	Present
Phenolic compound	Present
Nitriles isothiocyanate	Present
Saponine	Absent
Volatile oil	Absent

REFERENCES

1. G. V. Satyavati and K. Gupta, Medicinal Plants of India ICMR, **2**, 272-278 (1987).
2. A. Caceres, O. Cebrea, O. Morales, P. Miollinedo and P. Mendia, Pharmacological Properties of *M. Oleifera* J. Ethnopharmacol., **33(3)**, 213-216 (1991).
3. M. Leuck and H. Kunz, Synthesis of Active Principle from the Leaves of *M.oleifera* using s-Pent-4 enyl thioglycosides Carbohydr. Res., **312(1-2)**, 33-44 (1998).
4. N. A. Kumar and L. Pan', Antioxidant Action of *M. Oleifera* Against Antitubercular Drug Induced Lipid Peroxidation in Rat J. Med. Food., **6(3)**, 255-259 (2003).
5. Y. Absar, M. R. Uddin, M. A. Malik and K. Ahmed, Studies on Green Leafy Veg. of Bangladesh-2-Biological Availability of Carotene.
6. Y. S. Banginwar, Effect of Various Plant Extract on the Growth of *Vibrio Cholerae*. J. Micro World, **5(1)**, 1-3 (2003).
7. E. G. C. Chopra et al., Glossary of Indian Medicinal Plant CSIR, New Delhi 7st Edi., **26(5)**, (1984) pp. 131-133.

8. I. M. Dayrit, A. D. Akanta and I. M. Villasen, Studies on *M. Oleifera* Seed Part-I. The Antibiotic Compound and its Deactivation in Aqueous Solution, *Philip. J. Sci.*, **119**(1), 23-32 (1990).
9. World Health Organization, Summary of WHO Guidelines for the Assessment of Herbal Medicines, *Herbal Gram.*, **28**, 13-14 (1993).

Revised : 01.10.2012

Accepted : 02.10.2012