

## Black Holes

Pastushenko Vladimir Alexandrovich \*

Department of Space Exploration, Megatrend University, Beograd, Serbia

\* **Corresponding author:** Pastushenko Vladimir Alexandrovich, Department of Space Exploration, Megatrend University, Beograd, Serbia, E-mail: past.vlad2@gmail.com

**Received:** May 28, 2024, Manuscript No. TSSE-24-137504; **Editor assigned:** May 31, 2024, PreQC No. TSSE-24-137504 (PQ); **Reviewed:** June 17, 2024, QC No. TSSE-24-137504; **Revised:** June 03, 2025, Manuscript No. TSSE-24-137504 (R); **Published:** June 10, 2025, DOI. 10.37532/2320-6756.2025.14(3).388

### Abstract

We will talk about the properties of “black spheres” called “black holes”, within the framework of the properties of dynamic space-matter, which are subject to experimental testing. First of all, the presence of new quanta in the cores of planets, in the cores of stars, in the cores of galaxies, in the cores of quasars and in the cores of quasar galaxies. And first of all, stable quanta of the new substance.

**Keywords:** Dynamic space; Euclidean axiomatic; Parallelism; Length

### Introduction

It is generally accepted (in 2020) that there is a “supermassive compact object in the center of the galaxy.” And there is the fact of the presence of dynamic space-matter within the always dynamic ( $\varphi \neq \text{const}$ ) angle of parallelism [1].




FIG. 1. Dynamic space of a bunch of parallel straight lines.

### Literature Review

There is no matter outside space, and there is no space without matter, therefore space, as a form of matter, is one whole. Infinity ( $AC \rightarrow \infty$ ) cannot be stopped, therefore such dynamic space-matter always exists. The limiting case ( $(\varphi=0)=\text{const}$ ) of ( $(\varphi \neq 0) = \text{const}$ ) dynamic space-matter is the Euclidean axiomatics and Riemannian space in particular [2].

- A point is something of which nothing is a part”) (“Principles” by Euclid). and is a point something that has no parts,
- Line-length without width.
- and 5th postulate about parallel straight lines that do not intersect. If a straight line intersecting two straight lines forms

**Citation:** Alexandrovich PV. Black holes. J Space Explor. 2025;14(3):388.

© 2025 Trade Science Inc.

interior one-sided angles less than two right angles, then, extended indefinitely, these two straight lines will meet on the side where the angles are less than two right angles.




FIG. 2. Euclidean axiomatics.

Within the framework of the Euclidean ( $\varphi=0$ ) axes grid, we do not see dynamic ( $X+=Y-$ ), ( $X-=Y+$ ) space-matter and we will not be able to imagine it. Therefore, the axioms of dynamic space-matter are introduced as facts that do not require proof. Already in these axioms the problem of the Euclidean axiomatics of a point, as a set of indivisible sphere-points, is solved in one indivisible sphere-point, but already on (n) convergence, dynamic space-matter [3].




FIG. 3. Dynamic space-matter.




FIG. 4. Dynamic space-matter.

Any fixation (in experiments) of a non-zero ( $\varphi \neq 0$ ) angle of parallelism gives a multi-leaf Riemannian space. Now within the framework of the axioms of dynamic space-matter in the form:

- A non-zero, dynamic angle of parallelism ( $\varphi \neq 0$ )  $\neq$  const, a beam of parallel straight lines, determines the orthogonal fields ( $X-$ )  $\perp$  ( $Y-$ ) of parallel lines - trajectories, as isotropic properties, of space-matter.
- Zero angle of parallelism ( $\varphi=0$ ), gives a “length without width” with a zero or non-zero  $Y_0$  radius of a sphere-point “having no parts” in the Euclidean axiomatics.
- A bundle of parallel lines with zero angle of parallelism ( $\varphi=0$ ), “equally located to all its points,” produces many straight lines in one “widthless” Euclidean straight line [4].
- Internal ( $X-$ ), ( $Y-$ ) and external ( $X+$ ), ( $Y+$ ) fields of line-trajectories are non-zero  $X_0 \neq 0$  or  $Y_0 \neq 0$  material sphere-point, form an Indivisible Area of Localization НОЛ ( $X \pm$ ) or НОЛ ( $Y \pm$ ) dynamic space-matter.
- In unified fields ( $X-=Y+$ ), ( $Y-=X+$ ) orthogonal lines-trajectories, ( $X-$ )  $\perp$  ( $Y-$ ) there are no two identical sphere-points and lines-trajectories.

- A sequence of indivisible areas of localization, (X+), (Y+), (X+), ... along a radius  $X_0 \neq 0$  or  $Y_0 \neq 0$
- $X_0 \neq 0$  or a sphere-point on one line-trajectory gives n convergence, and on different trajectories m convergence.
- Each indivisible area of localization of space-matter corresponds to a unit of all its evolution criteria - CE, in a single (X=+Y+), space (Y=-X+) -matter at m-n convergence

$$HOJ = K\partial(X- = Y+)K\partial(Y- = X+) = 1, \quad HOJ = K\partial(m)K\partial(n) = 1,$$

in a system of numbers of units equal by analogy.

- Fixation of an angle ( $\varphi \neq 0$ )  $\neq$  const or ( $\varphi=0$ ) a bundle of straight parallel lines, space-matter, gives Euclid's 5<sup>th</sup> postulate and the axiom of parallelism [5].

Any point of fixed line-trajectories is represented by local basis vectors of Riemannian space:

$$e_i = \frac{\partial X}{\partial x^i} i + \frac{\partial Y}{\partial x^j} j + \frac{\partial Z}{\partial x^k} k, \quad e^i = \frac{\partial x^i}{\partial X} i + \frac{\partial x^j}{\partial Y} j + \frac{\partial x^k}{\partial Z} k,$$

with fundamental tensor  $e_i(x^n) * e_k(x^n) = g_{ik}(x^n)$  and topology  $(x^n = X, Y, Z)$  in Euclidean space. That is, Riemannian space is a fixed ( $\varphi \neq 0$ )  $\neq$  const state of dynamic ( $\varphi \neq$  const) space-matter. Local basis vectors correspond to the velocity space  $W^N = K^{+N} T^{-N}$ , in multidimensional space-time. Space-time is a special case of a fixed state of dynamic space-matter. At the same time, all Criteria for the Evolution of Matter are formed in multidimensional space-time.

They are presented in the “Unified Theory 2”, in the form of: (P=W<sup>2</sup>) potential, (F=P<sup>2</sup>) force, energy: charge PK=q (Y+ =X -) in electro (Y+=X-) magnetic fields, or mass PK=m (X+=Y-) in gravit (X+=Y-), mass fields, then density  $\rho = \frac{m}{V} = \frac{PK}{K^3} = \frac{1}{T^2} = v^2$  is the square of frequency, energy E=P<sup>2</sup> K, impulse (p=P<sup>2</sup>T), action ( $\hbar=P^2KT$ )..., a single space-matter [6].




FIG. 5. Unified criteria for the evolution of space-matter.

Let us immediately note the “point that has no parts” in the Euclidean axiomatics, and the non-zero radius  $X_0 \neq 0$  or  $Y_0 \neq 0$  material sphere-point of the axioms of dynamic space-matter. In addition, there is a minimum Planck length ( $\lambda=10^{-33}$ cm). These are questions of singularity that are not here, plus the mathematical prohibitions of division by zero.

As part of a dynamic ( $\varphi \neq$  const) space-matter, we have a non-stationary Euclidean space-time (X, Y, Z, cT), or a geodesic variable ( $x^s \neq$  const), fundamental tensor  $g_{ik}(x^s)$  Riemannian space. For example, the no stationary space of Lobachevski geometry, with variable asymptotes of hyperbolas. There is no such mathematics yet.

In other words, we will consider the issues of “black holes” in the axioms of Euclidean space-time, as a special case ( $\varphi=0$ ) or ( $(\varphi \neq 0)=$ const) dynamic ( $\varphi \neq$  const) space-matter.

**Discussion**

**Assumptions:** Within the framework of classical physics, even 100-200 years ago, and in the laws of conservation of energy

$$E_k = \frac{mv^2}{2} \quad \text{and} \quad E_n = mgh, \quad \text{for} \quad g = \frac{GM}{R^2} \quad \text{and} \quad h = R$$

Earth, the maximum speed was determined:  $\frac{v^2}{2} = \frac{GM}{R}$ ,  $v^2 = \frac{2GM}{R}$ , in which the body may not return to Earth ( $M$ ). And even then, the hypothesis of super massive ( $M \neq 0$ ) "black stars", from which light does not come out, arose. The sphere of such "black stars"  $R_0 = \frac{2GM}{c^2}$  was called the Schwarzschild sphere. The reason was considered to be Newton's gravitational force,  $F = \frac{2GMm}{R^2}$ . Here

$R$  is the distance between the centers of massive ( $M \neq 0$ ) and ( $m \neq 0$ ) massive spheres, the Earth and the Moon, for example. But if a small ball is lowered into the diametrical hole of a large massive sphere ( $R \rightarrow 0$ ), then the force does not increase indefinitely.




FIG. 6. Newton's law.

Newton's law doesn't apply here. Newton introduced the very concept of force from the springy collision of two balls, with inverse proportionality to their accelerations of expansion.

$$\frac{m_1}{m_2} = \frac{a_2}{a_1}, m_1 a_1 = m_2 a_2 = F.$$

Newton called this invariant of variable parameters force and said to measure it in newtons without any exchange interaction. Let us immediately note that in dynamic ( $\varphi \neq const$ ) space-matter, all the Evolution Criteria of the space of velocities and in Riemannian space too:  $e_i(x^n) = v_i$ ,  $e_k(x^n) = v_k$ ,  $g_{ik}(x^n) \equiv v^2$ , as a potential in the coordinate -time space of velocities  $W^N = K^+ N T^- N$ , in multidimensional space-time. For charges  $PC=q$  ( $Y+=X-$ ) in electric ( $Y+=X-$ ) magnetic fields, and masses  $PC=m$  ( $X+=Y$ ) in gravitational ( $X+=Y-$ ) mass fields, Maxwell and gravitational equations are derived fields.

$$c * rot_Y B(X-) = rot_Y H(X-) = \epsilon_1 \frac{\partial E(Y+)}{\partial T} + \lambda E(Y+)$$

$$rot_X E(Y+) = -\mu_1 \frac{\partial H(X-)}{\partial T} = -\frac{\partial B(X-)}{\partial T}; c * rot_Y M(Y-) = rot_Y N(Y-) = \epsilon_2 * \frac{\partial G(X+)}{\partial T} + \lambda * G(X+)$$

$$M(Y-) = \mu_2 * N(Y-); rot_Y G(X+) = -\mu_2 * \frac{\partial N(Y-)}{\partial T} = -\frac{\partial M(Y-)}{\partial T}$$

As well as transformations of the relativistic dynamics of the Special Theory of Relativity and quantum relativistic dynamics within the limits of parallelism angles.

$$\bar{X} = \frac{X - WT}{\sqrt{1 - W^2/c^2}}, \quad \bar{T} = \frac{T - \frac{W}{c^2} X}{\sqrt{1 - W^2/c^2}}, \quad \bar{W} = \frac{V + W}{1 + VW/c^2}$$

$$\bar{K}_Y = \frac{a_{11} K_Y + cT}{\sqrt{1 - W^2/c^2}}, \quad \bar{T} = \frac{K_Y/c + a_{22} T}{\sqrt{1 - W^2/c^2}}, \quad \bar{W}_Y = \frac{a_{11} W_Y + c}{a_{22} + W_Y/c}, \text{ in conditions, } (a_{22} \neq a_{11}) \neq 1,$$

For zero angles of parallelism in the Euclidean axiomatics, with velocities lower than the speed of light  $W_Y < c$ , there are limiting cases of transition of quantum relativistic dynamics of vector components,

$$a_{22} = (\cos(\alpha^0 = 0) = 1) = a_{11}, \quad a_{22} = 1, \quad a_{11} = 1, \quad Y = WT,$$

$$(\bar{K}_Y = \bar{Y}) = \frac{(a_{11} = 1)(K_Y = Y) \pm WT}{\sqrt{1 - W^2(X-)/c^2}}, \quad \bar{Y} = \frac{Y \pm WT}{\sqrt{1 - W^2/c^2}}, \quad \bar{T} = \frac{K_Y/c + (a_{22} = 1)T}{\sqrt{1 - W^2(X-)/c^2}}$$

In other words, in Euclidean axiomatics it is impossible in principle to create the Quantum Theory of Relativity. Both theories: Special Theory of Relativity and Quantum Theory of Relativity, allow superluminal ( $v_i=N*c$ ) velocity space:

$$\overline{W}_Y = \frac{c+Nc}{1+c*Nc/c^2} = c, \overline{W}_Y = \frac{a_{11}Nc+c}{a_{22}+Nc/c} = c, \text{ For } a_{11} = a_{22} = 1.$$

Already within the framework of such ideas, we will consider “black holes”. In classical physics with the Euclidean axiomatic of space-time, for super massive “black stars” ( $M \neq 0$ ), with a gravitational radius  $R_0 = \frac{2GM}{c^2}$ , of any mass in theory. And for the masses ( $M \neq 0$ ) = const, inside ( $R < R_0$ ) such a sphere, there must be a superluminal space of velocities ( $v_i > c$ ) or ( $v_i = N * c$ ), ( $N > 1$ ). This does not contradict either the Special Theory of Relativity or the Quantum Theory of Relativity. In the quantum coordinate system of the dynamic ( $\varphi \neq const$ ) space-matter, we are talking about superluminal space of velocities  $v_i = \alpha^{-N} * c$ , where  $\alpha = 1/137,036$  the constant.

But let's return to the laws of classical physics, in which Newton's law of gravity has limits of application, and did not answer the question WHY do masses attract? Studying Maxwell's equations, like electromagnetic fields with Lorentz transformations in two  $(x_0, y_0, z_0, ct_0)$  And  $(x_1, y_1, z_1, ct_1)$  coordinate systems, and from the laws of conservation of energy, back in 1905, Einstein derived a formula, which we will dwell on in more detail.

Body with non-zero( $m \neq 0$ ) mass, emits light with energy ( $L$ ) in system $(x_0, y_0, z_0, ct_0)$  coordinates, with the law of conservation of energy: ( $E_0 = E_1 + L$ ), before and after radiation. For the same mass in a different  $(x_1, y_1, z_1, ct_1)$ coordinate system, the law of conservation of energy with ( $\gamma = \sqrt{1 - \frac{v^2}{c^2}}$ ) Lorentz transformations, Einstein wrote in the form ( $H_0 = H_1 + L/\gamma$ ). Subtracting their difference, Einstein got:

$$(H_0 - E_0) = (H_1 - E_1) + L(\frac{1}{\gamma} - 1), \text{ or } (H_0 - E_0) - (H_1 - E_1) = L(\frac{1}{\gamma} - 1),$$

With separation of the radiation energy difference. Both inertial coordinate systems are moving, but  $(x_1, y_1, z_1, ct_1)$ moving at a speed ( $v$ ) relatively  $(x_0, y_0, z_0, ct_0)$ . And it is clear that blue and red light have an energy difference, which Einstein wrote down in the equation. Einstein wrote down the equation itself as the difference in kinetic energies in the first expansion.

$$(K_0 - K_1) = \frac{L}{2}(\frac{v^2}{c^2} ..) \text{ or } \Delta K = (\frac{\Delta L}{c^2}) \frac{v^2}{2}$$

Here ( $\frac{\Delta L}{c^2} = \Delta m$ ) the multiplier has the properties of the “radiant energy” mass, or:  $\Delta L = \Delta mc^2$ . This formula has been interpreted in different ways. The annihilation energy of  $E = m_0c^2$ the rest mass, or:  $m_0^2 = \frac{E^2}{c^4} - p^2c^2$ , in relativistic dynamics. Here is the mass with zero momentum( $p = 0$ ), has energy:  $E = m_0c^2$ , and the zero mass of the photon: ( $m_0 = 0$ ), has momentum and energy

$E = p * c$ . But Einstein derived another law of “radiant energy” ( $\Delta L = \Delta mc^2$ ), with mass properties. This is not the energy of a photon, and this is not the energy of ( $\Delta E = \Delta mc^2$ )a defect in the mass of nucleons in the nucleus of an atom. Einstein saw something that no one else saw. Like a moving charge, with the magnetic field induction of Maxwell's equations, a moving mass induces mass energy ( $\Delta L = \Delta mc^2$ ), which is what Einstein found. For example, a charged sphere inside a moving carriage has no magnetic field. But the compass on the platform will show the magnetic field of the sphere in the moving carriage. It was precisely this kind of inductive magnetic field, from the moving electrons of a conductor current, that Oersted discovered. Then there were Faraday's experiments, the induction of vortex electric fields in an alternating magnetic field, the laws of induction and self-induction, and Maxwell's equations. By analogy with the inductive energy of a magnetic field from a moving charge, Einstein derived the formula for the inductive, “radiant” energy of mass fields from moving non-zero masses, including stars in galaxies. And here Einstein went beyond the Euclidean ( $\varphi = 0$ )axiomatics of space-time. In the axioms of dynamic space-matter ( $\varphi \neq const$ ), we are talking about inductive  $m(Y -)$ mass fields, in complete analogy with Maxwell’s equations. This is what Einstein saw, and no one else. Already from the Equivalence Principle, the potential of the inductive mass field

$$v^2(Y - X +) = v * \cos\varphi_x(X+) * v * \cos\varphi_x(X+) = Gv^2(X +)$$

in a gravitational field, a constant follows ( $G = \cos^2\varphi_x$ )as a mathematical truth. And already writing the equation of the General

Theory of Relativity, Einstein took the gravitational potential of zero mass:  $\frac{E^2}{p^2} = c^2$ , in the form of  $\frac{L^2(Y-)}{p^2} = Gv^2(X+) = \frac{8\pi G}{c^4} T_{ik}$  an energy-momentum tensor.

A misconception about Einstein's General Theory of Relativity is that it is believed that non-zero mass is represented in the equation as the source of space-time curvature, as the source of gravity. In the equation of Einstein's General Theory of Relativity, as a mathematical truth in dynamic space-matter in its entirety:

$$R_{ik} - \frac{1}{2}Rg_{ik} - \frac{1}{2}\lambda g_{ik} = \frac{8\pi G}{c^4} T_{ik}.$$

There is no mass: ( $M = 0$ ), in its classical sense. In mathematical truth, this is the difference in relativistic dynamics at two fixed points in Riemannian space, one of which is reduced to the Euclidean sphere, in the external, non-stationary ( $\lambda \neq 0$ ) Euclidean space-time. In physical truth, in the equation of the General Theory of Relativity, Einstein, in the unified Criteria of Evolution, Newton's formula (law) is "hardwired":

$$E = c^4K, P = c^4T, (c_i^2 - c_k^2 = \Delta c_{ik}^2) = \frac{E^2}{p^2} = \left(\frac{K^2}{T^2} = c^2\right), \Delta c_{ik}^2 = Gv^2(X+) \neq 0$$

$$\Delta c_{ik}^2 = \frac{c^4 c^4 K^2}{c^4 c^4 T^2} = \frac{G(c^2 K_Y = m_1)(c^2 K_Y = m_2)}{c^2 (c^2 T^2 = K^2)} = \frac{Gm_1 m_2}{c^2 K^2}, \Delta c_{ik}^2 = \frac{Gm_1 m_2}{c^2 K^2}, \Delta c_{ik}^2 c^2 = F$$

As we see, in the equation of Einstein's General Theory of Relativity, the force of gravity acts in fields with zero mass. It reads: the difference in mass flows  $\Delta c_{ik}^2(Y-)$  in the external field of gravity  $c^2(X+)$ , with their Principle of Equivalence, gives strength.

And only now, we will consider the properties of "super massive" ( $M \neq 0$ ) compact ( $R \rightarrow 0$ ) objects discovered in the galactic core as a fact of rarity.

Under the conditions of:  $c^2 = \left(\frac{2G(M=0)}{(R=0)} = 0\right) \neq 0$ , under the conditions of the Planck limit length ( $10^{-33}$  cm), of the quantum field in space-time, under the conditions of the uncertainty principle, as well as the always dynamic one, of the quantum itself, under the conditions of a non-zero difference

$$R_{ik} - \frac{1}{2}Rg_{ik} \neq 0$$

energy-momentum tensor, *i.e.* ( $T_{ik} \neq 0$ ) energy, the presence of:  $c^2 = \left(\frac{2G(M=0)}{(R=0)} = 0\right) \neq 0$  gravitational potential as the reason for the curvature of the "black hole" space itself. In such listed conditions as arguments of mathematical truths, talk about a singularity in the center ( $R = 0$ ) "black hole", this is a conversation about nothing. There is no singularity at the center of the "black holes". The question is closed.

But there is a fact of the presence of "super massive compact objects" discovered in the core of galaxies. And there is another representation of the properties of such objects:

$$(R < R_0) = \frac{2GM}{(v_i > c)^2}$$

with the presence of superluminal space: ( $v_i > c$ ) inside ( $R < R_0$ ) such "black spheres" called "black holes". There are no "holes". The mass of such "black spheres" ( $M \neq 0$ ) is not zero. Next we will talk about the properties of "black spheres" called "black holes", within the framework of the properties of dynamic space-matter which are subject to experimental testing. First of all, the presence of new quanta in the cores of planets, in the cores of stars, in the cores of galaxies, in the cores of quasars and in the cores of quasar galaxies. And first of all, stable quanta of the new substance.

On colliding beams of positrons ( $e^+$ ), which are accelerated in a stream of quanta ( $Y- = \gamma$ ), photons of a "white" laser in the form of:

$$\text{НОЛ}(X \pm = p_1^+) = (Y- = e^+)(X+ = \nu_{\mu}^-)(Y- = e^+) = \frac{2m_e}{G} = 15,3 \text{ TeV},$$

On colliding beams of antiprotons ( $p^-$ ), occurs:

$$\text{НОЛ}(Y \pm = e_2^-) = (X- = p^-)(Y+ = e^+)(X- = p^-) = \frac{2m_p}{\alpha^2} = 35,24 \text{ TeV}.$$

indivisible and stable quanta of matter, similar to the substance of electron quanta.

We are talking about a quantum coordinate system  $OL_{ji}(m-n)$  in the space-matter of the Universe, in each  $OL_j$  level  $OL_i$  there are three ( $X=Y+$ ) charge and two ( $Y=X+$ ) mass isopotentials. And in this quantum coordinate system, "heavy" ( $p_j/e_j$ ) quanta, each of which has its own "depth" of energy levels ( $v_1/\gamma_i$ ) quanta of physical vacuum. Let's imagine them in the form of models such  $R_{ji}(m)$  Indivisible Regions of space - the matter of the Universe.




FIG. 7. Spectrum of indivisible quanta.

This is a certain sphere in space-matter, in the center of which are "heavy" ( $p_j/e_j$ ) quanta that determine "down" and "up" along the radius, up to the level ( $v_i/\gamma_i$ ) quanta of the physical vacuum of space-matter of the Universe, for any similar object within this sphere.

In the axioms of dynamic space-matter,  $HOЛ = KЭ(m)KЭ(n) = 1$ , we obtain for the masses ( $M$ ) of indivisible quanta in ( $OL_{ji}$ ) levels:

$$\begin{aligned}
 HOЛ &= M(e_1 = 1,15 E4)(k = 3.13)M(\gamma_0 = 3.13.E - 5) = 1 \\
 HOЛ &= M(e_2 = 3,524 E7)(k = 3.13)M(\gamma = 9,07 E - 9) = 1 \\
 HOЛ &= M(e_3 = 5,755 E11)(k = 3.86)M(\gamma_1 = 4.5.E - 13) = 1 \\
 HOЛ &= M(e_4 = 1,15 E16)(k = 3.13)M(\gamma_2 = 2,78 E - 17) = 1 \\
 HOЛ &= M(e_5 = 3,97 E19)(k = 3.13)M(\gamma_3 = 8.05.E - 21) = 1 \\
 HOЛ &= M(e_6 = 6,48 E23)(k = 3.83)M(\gamma_4 = 4,03 E - 25) = 1 \\
 HOЛ &= M(e_8 = 4,47 E31)(k = 3.14)M(\gamma_6 = 7,13 E - 33) = 1 \\
 HOЛ &= M(e_{26} = 9,1 E103)(k = 3.14)M(\gamma_{24} = 3,5 E - 105) = 1
 \end{aligned}$$

Obviously, we are talking about vortex mass ( $Y-$ )trajectories:  $c * rot_X M(Y- = \gamma_i) = \epsilon_2 * \frac{\partial G(X+)}{\partial T} + \lambda * G(X+)$  equations of dynamics in a circle ( $k = 3.14 = \pi = \frac{2\pi R=l}{2R}$ ) in each ( $OL_i$ ) level of the physical vacuum. These are spheres around a planet, star, galaxy, quasar.... Using quanta as an example:

$$\begin{aligned}
 HOЛ(X \pm = p_1^+) &= (Y- = e^+)(X+ = \nu_\mu^-)(Y- = e^+) = \frac{2m_e}{G} = 15,3 TeV, \\
 HOЛ(Y \pm = e_2^-) &= (X- = p^-)(Y+ = e^+)(X- = p^-) = \frac{2m_p}{\alpha^2} = 35,24 TeV,
 \end{aligned}$$

we are talking about the synthesis of matter ( $X \pm = p_1^+$ ) using colliding beams ( $e^+ e^+ \rightarrow p_1^+$ ) positrons with virtual quanta ( $\nu_\mu^-$ ), and ( $Y \pm = e_2^-$ ) on counter beams ( $p^- p^- \rightarrow e_2^-$ ) antiprotons and positrons with virtual quanta ( $e^+$ ) similar to electrons ( $e^- = \nu_e^- \gamma^+ \nu_e^-$ ). We can also talk about the consistent synthesis of "heavy" ( $p_j/e_j$ ) quanta, namely substances ( $X \pm = p_j^+$ ), for ( $Y-$ )<sub>A</sub> the ( $X-$ )<sub>A</sub> apparatus, in individual processes. ( $\dots \leftarrow p_6^+ \leftarrow e_5^+ \leftarrow p_3^+ \leftarrow e_2^+ \leftarrow p^+$ ) And ( $\dots \leftarrow p_7^+ \leftarrow e_6^+ \leftarrow p_4^+ \leftarrow e_3^+ \leftarrow p_1^+ \leftarrow e^+$ ) synthesis. The important thing is that the electron ( $e^-$ ) emits and absorbs a photon ( $\gamma^+$ ), but it cannot emit and absorb a "dark" photon ( $\gamma_0$ ). This "dark" photon is emitted and absorbed by a "heavy" electron ( $e_1$ )  $\rightarrow (\gamma_0)$ . In exactly the same way, a "heavy" proton ( $p_1$ )  $\rightarrow (\nu_\mu)$  emits and absorbs a muon neutrino. These are invisible quanta that do not interact and are non-contact with quanta ( $p^+/e^-$ ) atoms of the periodic table. We can neither see nor record them. But these invisible quanta (blue color in the indicated sequences) have charge is potentials and can form Structural Forms that are invisible to us, similar to ordinary ( $p^+/e^-$ ) atoms. These are: structures ( $\nu_\mu/\gamma_0$ ), ( $p_1/e_1$ )... This is how we consistently master the potentials of the core of planets, the core of stars, the core of galaxies and the core of quasars. But for ( $Y-$ )<sub>A</sub> apparatus, we can only form contact

quanta ( $p_4^+$ ) galactic nuclei and quanta ( $p_6^+$ ) substances of the core of quasars. And the device itself  $(Y-)_A$ , is consistently “immersed” in a physical vacuum, such as:  $HOI = (e_4)(k)(\gamma_2) = 1$ ,  $HOI = (e_6)(k)(\gamma_4) = 1$ , superluminal ( $\gamma_2 = 137 * c$ ), and ( $\gamma_4 = 137^2 * c$ ) velocity space.

This is completely acceptable in Special  $\overline{W}_Y = \frac{c+Nc}{1+c*Nc/c^2} = c$ , and Quantum  $\overline{W}_Y = \frac{a_{11}Nc+c}{a_{22}+Nc/c} = c$ , theories of relativity in Euclidean

$a_{ii} = \cos(\varphi = 0)$ ,  $a_{11} = a_{22} = 1$ , angles of parallelism. The apparatus itself  $(Y-)_A$  moves in the indicated sphere of space-matter of the Universe, in various levels of physical vacuum. It is worth noting that the volume of space-matter of a star is “immersed” in velocity space ( $\gamma = c$ ), the volume of galaxies is “immersed” in velocity space ( $\gamma_2 = 137 * c$ ), the volume of quasars is “immersed” in space ( $\gamma_4 = 137^2 * c$ ) already superluminal speeds.

### Admissible objects of the Universe

We will call the objects of the Universe “spheres-points”  $OJ_{ji}(n)$  convergence, in each fixed “point  $OJ_{ji}(m = const)$ ” quantum coordinate system. For example, objects:

$$HOI = M(e_2 = 3,524 E7)(k = 3.13)M(\gamma = 9,07 E - 9) = 1$$

similar to the kernel ( $p/e$ ) ordinary atoms, we are talking about quanta ( $p_2/e_2$ ) star cores. Stars with such a core have the maximum energy level of a physical vacuum, at the level ( $\gamma$ ) photon. Below the photon energy, the star does not manifest itself in a physical vacuum. Similar to proton radiation ( $p^+ \rightarrow \nu_e^-$ ) antineutrino, we are talking about radiation from antimatter matter and vice versa. That is: ( $p_8^+ \rightarrow p_6^-$ ), ( $p_6^- \rightarrow p_4^+$ ), ( $p_4^+ \rightarrow p_2^-$ ), ( $p_2^- \rightarrow p^+$ ), with the corresponding atomic nucleus: ( $p^+/e^-$ ) substances of an ordinary atom, ( $p_2^-/e_2^+$ ) antimatter core of the “stellar atom”, ( $p_4^+/e_4^-$ ) matter of the core of the galaxy, ( $p_6^-/e_6^+$ ) antimatter of the core of the quasar and “ ( $p_8^+/e_8^-$ ) matter of the core of the “quasar galaxy”. Further, we proceed from the fact that quantum ( $e_{*1}^-$ )

substances  $(Y- = p_1^-/n_1^- = e_{*1}^-)$  planetary cores emits a quantum

$$(e_{*1}^+ = 2 * \alpha * (p_1^- = 1,532E7 MeV)) = 223591 MeV, \quad \text{or: } \frac{223591}{p=938,28} = e_{*1}^+ = 238,3 * p$$

mass of the uranium nucleus, quantum of “antimatter”  $M(e_{*1}^+) = M(238,3 * p) = \frac{238}{92}U$ , uranium nuclei. There is such an “antimatter” ( $e_{*1}^+ = \frac{238}{92}U = Y-$ ) unstable, and decays exothermically into a spectrum of atoms in the core of planets.

At superluminal level  $w_i(\alpha^{-N}(\gamma = c))$  physical vacuum, such stars do not manifest themselves. Next, we are talking about the substance of ( $p_3^+ \rightarrow p_1^-$ ) the nucleus ( $Y- = p_3^+/n_3^0 = e_{*3}^+$ ) “black spheres” around which, in their gravitational field, globular clusters of stars form. Similarly, below, we are talking about radiation from antimatter matter and vice versa:

( $p_6^+ \rightarrow p_5^-$ ), ( $p_5^- \rightarrow p_3^+$ ), ( $p_3^+ \rightarrow p_1^-$ ), ( $p_1^- \rightarrow \nu_\mu^+$ ). The general sequence looks like:

$$p_8^+, p_7^+, p_6^-, p_5^-, p_4^+, p_3^+, p_2^-, p_1^-, p^+, \nu_\mu^+, \nu_e^- \dots$$

Further:  $HOI = M(e_4 = 1,15 E16)(k = 3.13)M(\gamma_2 = 2,78 E - 17) = 1$ . These quanta ( $p_4/e_4$ ) galactic nuclei are surrounded by individually emitted quanta ( $p_2/e_2$ ) the cores of stars are the reason for their formation. Such galactic nuclei, in the equations of quantum gravity, have spiral arms of mass trajectories, already:  $w_i(\gamma_2 = \alpha^{-1}c) = 137 * c$ , in superluminal speed space. Below the energy of light photons ( $w_i = 137 * c$ ) in a physical vacuum, galaxies do not manifest themselves. Outside galaxies, we are talking about quanta from the core of mega stars ( $Y- = p_5^-/n_5^- = e_{*5}^-$ ). They generate many quanta ( $e_{*5}^- = 2 * \alpha * p_5^- = e_{*4}^+ = 290p_4^+$ ) galactic nuclei. Likewise, further.

$$HOI = M(e_6 = 6,48 E23)(k = 3.83)M(\gamma_4 = 4,03 E - 25) = 1$$

We're talking about quanta ( $Y- = p_6^-/n_6^- = e_{*6}^-$ ) quasar nuclei, which also individually emit ( $p_4/e_4$ ) quanta of the core of galaxies. In other words, the quasar core is surrounded by quanta from the galactic core. They say that the quasar is in the center of the galaxy. Such quasars plunge into the level of physical vacuum to superluminal speeds  $w_i(\gamma_4 = \alpha^{-2}c) = 137^2 * c$ . This is deeper than the level of the physical vacuum of the galaxy. These are completely different objects. In other words, quasars bend space-matter at the level ( $\gamma_4$ ) quanta Next we talk about quanta of matter kernels ( $Y- = p_7^+/n_7^0 = e_{*7}^+$ ) “black spheres” around which, in their gravitational field, clusters of galaxies are formed, and further:

$$HOI = M(e_8 = 4,47 E31)(k = 3.14)M(\gamma_6 = 7,13 E - 33) = 1$$

We're talking about quanta( $p_8/e_8$ ) the nuclei of quasar galaxies, which also individually emit quanta( $p_6^-/n_6^- = e_{*6}^-$ ) quasar cores. Such quasar galaxies plunge into the level of physical vacuum to superluminal speeds  $w_i(\gamma_6 = \alpha^{-3}c) = 137^3 * c$ . Similarly, further.

In the axioms  $HO\mathbb{I} = K\mathbb{E}(m)K\mathbb{E}(n) = 1$ , or  $M_j(X+) * M_i(Y-) = 1$ , of dynamic space-matter, we are talking about the source of gravity gravitational  $M_j(X+)$  mass in  $O\mathbb{L}_j$  levels and inert  $M_i(Y-)$  mass in  $O\mathbb{L}_i$  levels of physical vacuum, with their Einstein equivalence principle in a single gravitational ( $X+ = Y-$ ) mass field. These masses:  $M_j * M_i = (M = \Pi K)^2 = 1$ , in the form of a quadratic form, are presented in the quantum fields of their interaction:

$$\hbar = Gm_0 \frac{\alpha}{c} Gm_0(1 - 2\alpha)^2 = GM_j \frac{\alpha}{c} GM_i(1 - 2\alpha)^2 = \frac{(6,674 * 10^{-8})^2 * (1 - 2/(137.036))^2}{137.036 * 2.993 * 10^{10}} = 1.054508 * 10^{-27}$$

in quantum:  $G(X+) \left[ \frac{K}{T^2} \right] = \psi \frac{\hbar}{\Pi^2 \lambda} G \frac{\partial}{\partial t} grad_n Rg_{ik}(X+) \left[ \frac{K}{T^2} \right]$ , gravity ( $X+ = Y-$ ) mass fields. This equation of quantum gravity follows directly from the equation of Einstein's General Theory of Relativity. Thus, the maximum mass  $M_j(X+)$  source of gravity is determined by  $M_i(Y-)$  the inertial mass of mass ( $Y- = \gamma_i$ ) fields in  $O\mathbb{L}_i$  levels of physical vacuum, like an object  $O\mathbb{L}_{ji}(n)$  convergence or:  $HO\mathbb{I} = O\mathbb{L}_{ji}(n) = M_j(X+) * M_i(Y- = \gamma_i) = 1$ . Thus, we obtain the maximum masses in the Universe: for example, for a star  $M_j(X+) = M_2(p_2^-/n_2^0) = 1/(\gamma)$  in conditions  $(e_2^+(k)\gamma) = 1$ . Likewise:

Limit mass of planets, for  $1MeV = 1.78 * 10^{-27}g$ :

$$\frac{1}{\gamma_0} = \frac{1}{3.13 * 10^{-5} MeV * 1.78 * 10^{-27}g} = M_1(p_1^-/n_1^-) \approx 1.8 * 10^{31}g \approx \frac{M_s}{100}, \text{ where } (M_s = 2 * 10^{33}g) \text{ is the mass of the Sun. Next is the maximum mass of stars with a core of antimatter:}$$

$$\frac{1}{\gamma} = \frac{1}{9.07 * 10^{-9} MeV * 1.78 * 10^{-27}g} = M_2(p_2^-/n_2^-) \approx 6.2 * 10^{34}g \approx 31M_s, \text{ or ranging from } \frac{M_s}{100} \text{ before } 31M_s \text{ macc.}$$

Similarly, the maximum mass ( $p_3^+/n_3^0 = e_{*3}^+$ ) "black spheres", with a core of matter:

$$\frac{1}{\gamma_1} = \frac{1}{4.5 * 10^{-13} MeV * 1.78 * 10^{-27}g} = M_3(p_3^+/n_3^0) \approx 1.25 * 10^{39}g \approx 625220M_s$$

maximum mass of the galaxy, ( $p_4^+/n_4^0 = e_{*4}^+$ ) with a core of matter:

$$\frac{1}{\gamma_2} = \frac{1}{2.78 * 10^{-17} MeV * 1.78 * 10^{-27}g} = M_4(p_4^+/n_4^0) \approx 2 * 10^{43}g \approx 10^{10}M_s$$

the maximum mass of an extragalactic megastar, ( $p_5^-/n_5^- = e_{*5}^-$ ) with an antimatter core:

$$\frac{1}{\gamma_3} = \frac{1}{8.05 * 10^{-21} MeV * 1.78 * 10^{-27}g} = M_5(p_5^-/n_5^-) \approx 7 * 10^{46}g \approx 3.5 * 10^{13}M_s,$$

the maximum mass quasar, ( $p_6^-/n_6^- = e_{*6}^-$ ) with an antimatter core:

$$\frac{1}{\gamma_4} = \frac{1}{4.03 * 10^{-25} MeV * 1.78 * 10^{-27}g} = M_6(p_6^-/n_6^-) \approx 1.4 * 10^{51}g \approx 7 * 10^{17}M_s,$$

Each core of such objects  $O\mathbb{L}_{ji}(n)$  convergence, generates a set of corresponding quanta ( $2 * \alpha * p_j^\pm = e_{*j}^\mp = N p_{j-1}^\mp$ ) indicated in the table, and emits ( $p_j^\pm \rightarrow p_{j-2}^\mp$ ). This is a lot ( $N$ ) quanta of the core of planets, stars, galaxies, quasars....

### Conclusion

For example, the core of the Sun, like a star, emits hydrogen nuclei ( $p_2^- \rightarrow p^+ \rightarrow v_e^-$ ) and electron antineutrino, but generates ( $2 * \alpha * p_2^- = e_{*2}^+ = N p_1^+$ ) quanta of, let's say, "stellar matter" ( $p_1^+/e_1^-$ ) in the solid surface of the star. This is "star stuff" ( $p_1^+/e_1^-$ ) cannot interact with hydrogen ( $p^+/e^-$ ), but can emit muon antineutrino ( $p_1^+ \rightarrow v_\mu^-$ ), which in the Earth's atmosphere forms muons, which in decays give: ( $e^+$ ) positrons: ( $Y_\pm = \mu$ ) = ( $X- = v_\mu^-$ ) ( $Y+ = e^+$ ) ( $X- = v_e^-$ ). Or, quanta core of a mega star with ( $p_5^-/n_5^- = e_{*5}^-$ ) emit quanta ( $p_5^- \rightarrow p_3^+$ ) of matter, but generate quanta from the nuclei of galaxies ( $2 * \alpha * p_5^- = e_{*5}^+ = N p_4^+$ ). We see, as it were, the "surface" of the galaxy, but the core of such an object  $O\mathbb{L}_{ji}(n)$  convergence, has a mass ranging from ( $10^{10}M_s$ ) before ( $3.5 * 10^{13}M_s$ ) mass of the Sun.

We are talking about valid objects  $O\mathbb{L}_{ji}(n)$  convergence, in the dynamic space-matter of the Universe. In this case, the calculated cause-and-effect relationships are indicated.

## References

1. Skenderis K, Taylor M. The fuzzball proposal for black holes. *Phys Reports*. 2008;467(4-5):117-171.
2. Alexander DM, Hickox RC. What drives the growth of black holes? *New Astron Rev*. 2012;56(4):93-121.
3. Taylor EF, Wheeler JA. *Exploring black holes*. San Francisco: Addison Wesley Longman, 2000.
4. Novikov ID, Frolov VALERYP. *Physics of black holes*. Springer Science & Business Media, 2013.
5. Hawking SW. Black holes and thermodynamics. *Phys Rev D*. 1976;13(2):191.
6. Bekenstein JD. Black holes and the second law. In *JACOB BEKENSTEIN: The Conservative Revolutionary*, 2020:303-306.