



Trade Science Inc.

# Natural Products

*An Indian Journal*


---

**Full Paper**

NPAIJ, 3(1), 2007 [54-57]

## Chemical Composition Of The Essential Oil Of *Heracleum Persicum* Seeds Of Iran



Corresponding Author

Avat(Arman)Taherpour  
Chemistry Department, Graduate School, Islamic  
Azad University, P.O.Box 38135-567, Arak Branch,  
Arak, (IRAN)

†Unusual Molecule and Reactive Intermediate  
Group, Chemistry Building, School of Microbial  
& Molecular Sciences, The University of  
Queensland-QLD 4072, Brisbane, (AUSTRALIA)  
E-mail: a.taherpour@uq.edu.au

Received: 6<sup>th</sup> February, 2007Accepted: 21<sup>st</sup> February, 2007Web Publication Date : 25<sup>th</sup> February, 2007

Co-Authors

Ramin Karimzadeh

Chemistry Department, Graduate School, Islamic Azad University,  
P.O.Box 38135-567, Arak Branch, Arak, (IRAN)

### ABSTRACT

The volatile chemical constituents of the essential oil of *Heracleum Persicum* seeds, that is growing wild in Ardebile-Iran were investigated by GC and GC/MS technique. Twenty-nine compounds, representing 75% of the total oil were identified and 12 unknown compounds detected. The main components were: hexylbutyrate(37.7%), hexylbutanoate(36.7%), octylacetate(16.3%), hexyl-2-methylbutanoate(5.7%), hexylisobutyrate (4.7%), hexylhexanoate(4.3%), heptyl-2-methyl-butyrates(2.3%), n-butylbutanoate(2.25%), hexylvalerate(1.9), octyl-butanoate(1.7%) and linalole (1.5%). © 2007 Trade Science Inc. - INDIA

### KEYWORDS

*Heracleum persicum*;  
Octylacetate;  
Hexyl-2-methylbutanoate.

### INTRODUCTION

*Heracleum* genus has 10 species in Iran. *H.persicum* desf. ex fischer (syn. *H.pubescens* Rech., *H.glabrescens* boiss. and hohen.) (Apiaceae) is an annual herb, indigenous to the alborz region, the northern part of Iran, where it grows at an altitude ranging from 2000

to 3000m<sup>[1-4]</sup>. The genus *heracleum* are noted for their rich furanocoumarin content. Furanocoumarins are effective dermal photosensitizing agents, and are widely used in the treatment of leucoderma and in various 'suntan' lotions. *Heracleum Persicum* desf (umbellifereae) is an annual plant which grows in the northern parts of Iran. The seeds of this plant is

used as spices and the young stems are used for making pickles<sup>[5]</sup>. The *Heracleum persicum* materials of this study were collected from ardebil-Iran in aug. of this year. A voucher specimen has been deposited in herbarium of research institute of forest and rangelands, ardebil, Iran. Its fruits are used commonly in Iran as spices, while the fruits and the young shoots are used in the preparation of pickles. From the aerial parts of this herb, especially the fruits were administered because of their carminative activity in local medicine by folks<sup>[3]</sup>.

### Analytical methods

Dried aerial parts of *Heracleum persicum* were subjected to hydrodistillation for 5 hours using clevenger-type apparatus to produce yellow oil in 0.28%(w/w) yield. The oil of the aerial parts of *Heracleum persicum* was examined by GC/MS(GC:HP 6890, MS:HP 5973), column(HP5-MS, 30m×0.25mm fused silica capillary column, film thickness 0.32µm) by temperature program 60°C(3 min)-210°C(2min) at the rate of 6°C/min(injection temperature 250°C, carrier gas: helium(with purity 99.999%), detector temperature 150°C, ionization energy in mass was 70eV, mass range 10-300 amu, and scan time was 1 sec.

The list of identified components is presented in TABLE-1. The constituents were identified by comparing their MS spectra with those in computer library or with authentic compounds. The identifications were confirmed by comparison of their retention indices either with those of authentic compounds or with data in the literature<sup>[6-10]</sup>.

In the aerial parts of *Heracleum persicum* the major identified components and the relative amounts based on peak area were: hexylbutyrate(37.7%), hexylbutanoate(36.7%), octylacetate(16.3%), hexyl-2-methylbutanoate(5.7%), hexylisobutyrate(4.7%), hexylhexanoate(4.3%), heptyl-2-methylbutyrate(2.3%), n-butylbutanoate(2.25%), hexylvalerate(1.9), octylbutanoate(1.7%) and linalole(1.5%).

### DISCUSSION

Plants from genus *Heracleum* have been previously studied and reported in the literature. There are some nice studies of the chemical composition of essen-

tial oil of *Heracleum* species<sup>[11-16]</sup>. A report in literature shows that roots of *Heracleum persicum* have been investigated because of their furanocoumarins; five of which compounds were isolated and identified<sup>[11]</sup>. The literature survey presence of such compounds in leaves and seeds of this species<sup>[12]</sup>. From the diethyl ether extracts of the fruits of *Heracleum persicum*, an aglycone was identified which was demonstrated to be quercetin<sup>[13]</sup>. In this investigation, the volatile constituents of the essential oil of *Heracleum persicum* growing wild in ardebile-Iran were investigated by GC and GC/MS technique. As could see in TABLE 1, two component i.e. hexylbutyrate(37.7%) and hexylbutanoate(36.7%) have the highest percentage and more than 35% among the known compounds in the essential oil of this herb. Some other components like octylacetate(16.3%) and hexyl-2-methylbutanoate(5.7%), are located in the second level of the concentration in the essential oil. Although, in accordance with the data in TABLE 1, some components i.e. hexylisobutyrate(4.7%), hexylhexanoate(4.3%), heptyl-2-methylbutyrate(2.3%), n-butylbutanoate(2.25%), hexylvalerate(1.9), octylbutanoate(1.7%) and linalole(1.5%) have the medium up to low relative percentages, could see some important compounds with effects like mold and mildew preventive, microscopy, preservative and antioxidant with percentages under 1.5%. Biological and aroma effects of the main and minor compounds of the essential oil of *Heracleum persicum* are discussable in terms of their possible use in medicine, cosmetics and foods. Perhaps, the high densities of the main compounds give some biological activities to the essential oil or to this herb. Although no records of toxicity have been found for this plant, it belongs to a family that includes many poisonous plants so some caution is advised<sup>[9,10]</sup>.

### CONCLUSION

*Heracleum persicum* is one of the *Heracleum* genus of that was collected from west area in Iran. It is utilized as the medicinal herb for the various purposes in local and traditional medicine by folks in Iran. Twenty-nine compounds, representing 75% of the total oil were identified and 12 unknown compounds detected.

## Full Paper

TABLE 1: Essential oil constituents of *Heracleum persicum*

No.	Name of Compounds	Scan	KI*	RT**	%
1	Unknown	15	846.478	4.131	0.938
2	Unknown	62	846.788	4.260	2.368
3	Butyl-2-methylpropanoate	496	940.258	5.444	2.368
4	Isobutylbutanoate	515	942.5	5.496	0.308
5	Unknown	592	951.551	5.706	0.505
6	n-Butylbutanoate	846	981.422	6.399	2.25
7	n-Octanal	909	988.836	6.571	1.017
8	Isobutyl-3-methylbutanoate	932	991.551	6.634	0.490
9	Hexylethanoate	994	998.836	6.803	1.501
10	Para-Cymene	1102	1010.120	7.098	1.057
11	cis- $\beta$ -Ocimene	1208	1020.783	7.387	0.320
12	n-Butyl-2-methylbutanoate	1238	1023.843	7.469	0.402
13	Butylpentanoate	1278	1029.910	7.578	1.057
14	$\gamma$ -Terpinene	1398	1040.111	0.905	0.784
15	1-Octanol	1504	1050.932	8.195	0.929
16	Linalol	1759	1076.865	8.890	1.529
17	Hexylpropinoate	1796	1080.634	8.991	0.714
18	Unknown	1945	1095.820	9.398	0.343
19	Hexylisobutyrate	2180	1119.029	10.039	4.703
20	Hexylbutanoate	2628	1162.050	11.235	32.71
21	Hexylbutyrate	2628	1163.021	11.262	37.76
22	Unknown	2659	1166.043	11.346	5.454
23	Unknown	2670	1167.122	11.376	0.393
24	Decanal	2695	1169.604	11.445	0.200
25	Octylacetate	2772	1177.517	11.655	16.297
26	Hexyl-2-methylbutanoate	2958	1195.396	12.162	5.745
27	Hexylvalerate	2987	1198.237	12.241	1.902
28	Unknown	3731	1273.137	14.272	0.188
29	Octylisobutyrate	3823	1282.399	14.523	1.383
30	Unknown	4078	1308.423	15.219	0.447
31	Hexylhexanoate	4162	1317.231	15.448	4.307
32	Octylbutanoate	4182	1319.308	15.502	1.676
33	Unknown	4237	1325.077	15.652	0.247
34	Unknown	4436	1345.962	16.195	0.121
35	Heptyl-2-methylbutyrate	4524	1355.231	16.436	2.286
36	Octylvalerate	4554	1358.346	16.517	0.714
37	Unknown	5618	1475.875	19.421	0.220
38	Heptyloctanoate	6940	1627.387	23.028	0.200
39	Unknown	2840	1183.813	11.840	0.178
40	n-Octylpropinoate	3486	1248.450	13.603	0.141
41	Undeca-6-ol	3636	1263.542	14.012	0.093

\* Kovats index; \*\* Retention time

These components were identified by GC and GC/MS technique. In this herb, hexylbutyrate(37.7%) and hexylbutanoate(36.7%) have the most percentages among compounds of the essential oil. Some other components like octylacetate(16.3%) and hexyl-2-methylbutanoate(5.7%) are located in the second level of the concentration in the essential oil. The folks in common medicinal have been used from the aerial parts of this herb because of their carminative activity.

## REFERENCES

- [1] I.Mandenova, K.H.Rechinger; 'Heracleum, Flora Iranica Umbelliferae', In:(Ed), 492-502 (1987).
- [2] A.Parsa; 'Flore de l'Iran', Danech, Tehran, 2, 481 (1948).
- [3] G. Amin; 'Iranian Traditional Medicinal Plants'. Research Deputy of Health Ministry, Tehran, 130 (1991).
- [4] F.Mojab, B.Nickavar; Iranian Journal of Pharmaceutical Research, 245-247 (2003).
- [5] Y.Aynehchi, Z.Aliabadi, M.H.Salehi Sormaghi; ISHS Acta Horticulturae, I International Symposium on Spices and Medicinal plants, No. of Article: 43, 73, 1 (1978).
- [6] R.P.Adams; 'Identification of Essential oil Components by Gas Chromatography/Mass Spectroscopy', Allured Publishing Corporation, Illinois, 78-330 (1995).
- [7] 'Eight Pick Index of Mass Spectra.', 2<sup>nd</sup> Ed., Mass Spectroscopy Data Center, Reading, U.K., (1974).
- [8] W.Jennings, J.Shibamoto; 'Quantitative Analysis of Flavour and Fragrance Volatiles by Capillary Gas Chromatography', Academic Press, New York, (1980).
- [9] Dictionary of Natural Product, Chapman & Hall, U.K. (1992).
- [10] G.G.Hawley; 'Condensed Chemical Dictionary', 13<sup>th</sup> Ed., Van Nostrand Reinhold Company Inc., New York, USA, (1997).
- [11] Y.Aynehchi, Z.Aliabadi, M.H.S.Surmaghi; Acta Hort., 73, 103-107 (1978).
- [12] A.Merijanlian, T.Colasurdo, P.Samtak, J.Ullrich, J.Spagnuolo; Rev.Latinoam.Quim., 11, 51-53 (1980).
- [13] B.Ghods; Bull.Trav.Soc.Pharm.Lyon., 20, 3-8 (1976).
- [14] J.J.C.Scheffer, R.Hiltunen, Y.Aynehchi, M.von Schantz, A.Baerheim-Svendsen; Planta Med., 50, 56-60 (1984).
- [15] F.Mojab, A.Rustaiyan, A.R.Jasbi; DARU (and the literature cited there in.), 10, 6-8 (2002).
- [16] Y.K.Sarin, Kapoor; Perfumery Essent.Oil Record, 54, 437-443 (1963).