



Trade Science Inc.

ISSN : 0974 - 7532

Volume 6 Issue 9

Research & Reviews in

BioSciences

Regular Paper

RRBS, 6(9), 2012 [258-263]

## Cytotoxic and genotoxic effects of *Bacopa monnieri* in the cultured human peripheral blood lymphocytes

Srinivas Rao Bammidi<sup>1</sup>, Sharan Suresh Volluri<sup>2</sup>, Nandam Sree Satya<sup>1</sup>, Meena Vangalapati<sup>1\*</sup>

<sup>1</sup>Centre of Biotechnology, Department of Chemical Engineering, AUCE (A),  
Andhra University, Visakhapatnam - 530 003, (INDIA)

<sup>2</sup>Department of Biotechnology, DVR & HS MIC College of Technology, Kanchikacherla, (INDIA)

E-mail: bsrinivas3441@gmail.com; sharansuresh.volluri@yahoo.co.uk;  
nandam.sree@gmail.com; meena\_sekhar09@yahoo.co.in

Received: 28<sup>th</sup> July, 2012 ; Accepted: 15<sup>th</sup> September, 2012

### ABSTRACT

*Bacopa monnieri* is a medicinal herb which has been valued in ayurvedic medicine for its different activities. In the present studies, Bacopa methanolic extract was prepared by Soxhlet extraction and then evaluated for the Cytotoxicity and Genotoxicity in cultured human peripheral blood lymphocytes. Mitotic Index (MI) Cell Proliferation Kinetics (CPK) and Sister-chromatid exchanges (SCE) were scored to measure the cytotoxic and genotoxic effects of the Bacopa extract in cultures set up from the three different healthy donors. The treatment of the cell culture was done employing two different Bacopa methanolic extract concentrations (500 & 1000 µg/ml) and the control (did not receive any additive). Our present studies revealed that the MI, CPK and SCES for control are 4.28, 0.472 and 5.813 respectively where as the Bacopa extracts has the MI, CPK and SCES for 500µg/ml are 4.09, 0.46 and 5.386 respectively and for 1000 µg/ml are 3.62, 0.468 and 5.35 respectively. From the results, we can conclude that the MI and frequency of SCES of Bacopa methanolic extracts are almost similar to that of control which indicates that the Bacopa plant extracts has no significant cytotoxicity and genotoxicity effects in cultured human peripheral blood lymphocytes. © 2012 Trade Science Inc. - INDIA

### INTRODUCTION

In recent years that has been a surge in the demand for the plant based drugs and food additives. Medicinal plants, herbs and spices as herbal remedies are known to Ayurveda in India since long times. The value of medicinal plants, herbs, spices as herbal remedies is being lost due to lack of awareness and deforestation. The result is many valuable medicinal herbs are becoming rare and precious information is lost. Medicinal herbs have been used in folk medicine for millennia. Simply, in recent times, scientific study of their effects has flourished. Nevertheless, some of them

can cause adverse effects or have the potential to interact with other medications<sup>[1]</sup>. Moreover, there is little information on the potential risk to health of such herbs. Based on their long-term use by humans, one might expect herbs used in traditional medicine to have low toxicity. It is known that green plants in general are a primary source of antimutagens as well as natural toxic agents<sup>[2]</sup>, and many plants contain cytotoxic and genotoxic substances. Recent investigations have revealed that many plants used as food or in traditional medicine have mutagenic, cytotoxic and genotoxic effects in vitro and in vivo assays<sup>[3]</sup>.

*Bacopa monnieri* is a perennial creeper, faintly aro-

matic and a valuable medicinal herb of which is distributed throughout tropical and subtropical regions of World such as India, China, Nepal, Srilanka, and Indonesia etc. Traditionally, *Bacopa monnieri* has been valued for centuries in ayurvedic medicine for the treatment of leprosy, ulcer, asthma, bronchitis, elephantiasis, anxiety, urethritis, eye troubles, diarrhoea among children, skin diseases, wound healing etc.,. Phytochemical analysis of *Bacopa monnieri* plant extracts revealed the presence of various biochemical compounds such as alkaloids, flavonoids, glycosides, triterpenoids and Saponins etc. Several pharmacological studies on *Bacopa monnieri* reported that it has anti-inflammatory<sup>[4]</sup>, antibacterial, antimicrobial<sup>[4]</sup>, antispasmodic, antioxidant<sup>[5]</sup>, antitubercular<sup>[6]</sup>, immunomodulatory, antiarthritic<sup>[7]</sup>, anticancer<sup>[8]</sup> and antiulcer activities. In this study, Cytotoxic and genotoxic effects of *Bacopa monnieri* in cultured human peripheral blood lymphocytes have been evaluated.

### Mitotic index (MI) as a measure of cytotoxicity

Cell population growth occurs as cells pass through interphase and mitosis to complete the cell cycle. Many cells lose the capacity to divide as they mature or divide only rarely. Other cells are capable of rapid cell division. Cytotoxicity is a quality of being toxic to cells. Treating cells with a cytotoxic compound can result in a variety of cell fates. Examples of toxic agents are a chemical substance, plant extract, an immune cell or some types of venom. By quantifying aspects of a dividing cell population; we can examine how cells differ in their capability to divide. Experimentally, we can change properties of the cell's environment and quantify the effects on cell division "as discussed by Kannan T.P et al.<sup>[9]</sup>, The way to quantify cell division is by using the mitotic index. If you administer colchicine or other colchicine-derivative medications (i.e. colcemid) you can arrest the cell cycle at this point leaving the chromosomes in their visible form. Colchicine disrupts the microtubule formation which is necessary for the spindle fibers to separate the chromosomes during anaphase. Mitotic index is a measure for the proliferation status of a cell population. It is defined as the ratio between the number of cells in mitosis and the total number of cells. The mitotic index can be worked out from a slide, even with light microscopy. It is the number of cells containing visible chromosomes divided by the total

number of cells in the field of view.

### Sister chromatid exchange test as a measure of genotoxicity

Genotoxicity is a quality of being toxic to cell's genetic material affecting its integrity. Genotoxic substances are known to be potentially mutagenic or carcinogenic, specifically those capable of causing genetic mutation and of contributing to the development of tumors. This includes certain chemical compounds, plant extracts and certain types of radiation. The Sister Chromatid Exchange (SCE) assay is a short-term test for the detection of reciprocal exchanges of DNA between two sister chromatids of a duplicating chromosome. The SCE test is usually performed on human peripheral blood lymphocytes. As peripheral lymphocytes are in the resting G<sub>0</sub> stage of the cell cycle, they have to be stimulated to divide by an aspecific antigen, like phytohaemagglutinin. To collect a sufficient number of mitotic cells, a spindle inhibitor like colcemid may be added shortly before fixation (at 72 hours), to block cells in (pro) metaphase of the second mitosis. To allow for a differential staining that enables the researcher to distinguish both chromatids, BrdU (bromo-deoxy-uridine) is added to the culture medium for the duration of two complete cell cycles. The Giemsa staining was able to stain due to the presence of bromo-deoxy-uridine (BrdU) analogous base which was introduced to the desired chromatid.

Chromatids in which only one strand of DNA incorporated BrdU show a normal dark Giemsa staining, whereas those with two substituted strands stain less darkly. If an exchange occurred, this can be seen as the dark part changes to the other arm: "harlequin chromosomes". The technique can also be used to estimate the division rate, as first metaphases stain uniformly dark "as discussed by Kannan T P et al.<sup>[9]</sup> and "Maria P et al.,<sup>[10]</sup>

## MATERIALS AND METHODS

### Extraction and processing of the plant material

The fresh whole plant of *Bacopa monnieri* was collected from Araku Valley, Near Visakhapatnam, and Andhra Pradesh, India. The plant material was garbled and air dried under the shade up to 48 days. By using the kitchen blender the plant material was grounded to a fine powder form. Powdered plant material (5g) was mixed with 500 ml of methanol. Using the Soxhlet ap-

## Regular Paper

paratus continuous extraction was done for 24 hrs. The extract was filtered and methanol was evaporated on a rotary evaporator under vacuum at a lower temperature of 50°C to dryness. These plant extracts were weighed and preserved at 4°C in a desiccator.

### Blood sample collection

Three fresh blood samples from the volunteers were collected, generally from the arm by venipuncture, and placed into a heparinised tube.

### Peripheral blood culture initiation

Few drops of whole blood (0.5ml) was cultured in medium RPMI 1640 and a mitogen phytohemagglutinin (PHA) 100µL was added. *Bacopa monnieri* plant extracts of 2 different doses (500µg/ml and 1000 µg/ml) were added. And for mutagen cultures we were added Mitomycin-c 0.1 µg. After 24<sup>th</sup> hour of the cultures BrdU was added. Peripheral blood cultures were placed in a 37°C incubator for 72 hrs.

### Harvesting of the blood cultures

By administering colchicine or other colchicine-derivative medications (i.e. colcemid) you can arrest the cell cycle at this point leaving the chromosomes in their visible form. Colchicine disrupts the microtubule formation which is necessary for the spindle fibers to separate the chromosomes during anaphase. To arrest the cell division at metaphase, 0.1 ml (1 µg) of colchicine was added to the cultures just 90 minutes before harvesting. After addition of colchicine the cultures were incubated for 40 minutes and centrifuged at 1200 RPM for 10 minutes. To the pellet pre-warmed hypotonic solution (0.75M KCl) of 6ml was added. Incubated for 10 minutes and centrifugation was done at 1200RPM for 10 minutes.

### Fixation of blood cultures

To the pellet 5 ml of fixative solution (3:1 ratio of methanol and acetic acid) was added and centrifuged. 4-5 times of fixative washes were given. The fine pellet was dropped onto microscopic sterile and chilled slides.

### Preparation of metaphase chromosome spreads

Chromosome spreads were prepared by gently dropping the cell suspension from a height of 1 foot on to a clean grease free chilled slide and air-dried.

### Mitotic index (MI) assay

The slides were stained with 10 per cent Leishman's stain in phosphate buffer saline (pH 6.8) for 5 minutes. The slides were dried, mounted and the chromosomes were observed under the microscope<sup>[9]</sup>. Slides were prepared from each blood culture and a total of 1000 lymphocytes were counted per culture to determine the mitotic index. The mitotic index was used to determine if the *Bacopa monnieri* plant extracts produced any Cytotoxicity. It is a measure for the proliferation status of a cell population. It can be defined as the ratio between the number of cells in mitosis and the total number of cells or the number of cells containing visible chromosomes divided by the total number of cells in the field of view. It can be calculated as follows:

$$\text{Mitotic Index (MI)} = \frac{\text{Number of dividing cells}}{\text{Total number of cells}} \times 100$$

### Cell proliferation kinetics (CPK)

Cell-cycle specific patterns were determined by M<sub>1</sub>, M<sub>2</sub>, and M<sub>3</sub> metaphases. These were defined by the number of cell cycles completed. M<sub>1</sub> metaphases substituted BrdU unifarly in both chromatids without differential staining. Chromosomes of M<sub>2</sub> metaphases contained a bifilar substitution of BrdU in one chromatid, which, consequently, was lightly stained, thus displaying the typical sister chromatid differentiation pattern. Bifilar incorporation of BrdU in the third cell cycle yielded light staining of both chromatids. Such metaphases were classified as M<sub>3</sub> metaphases. The cell proliferation kinetic index is calculated using

$$\text{Cell Proliferation Kinetics (CPK)} = \frac{M_1 + M_2 + M_3}{100}$$

Where M<sub>1</sub> are Dark stained metaphases, M<sub>2</sub> are Differential stained metaphases and M<sub>3</sub> are Light stained metaphases.

### Sister chromatid exchange assay

After the third day of the preparation of the slides, then they were used for Sister Chromatid Exchange assay. For this assay we kept a drop of Hoechst 3325 solution onto the slide and mounted with cover slip for 25 minutes. Washed the slides with water and kept for drying in the dark chamber and then placed the slides in petridish which contains 2XSSC, for 25 minutes and exposed to UV for 40 minutes. Washed the slides with water and dried for few minutes. Incubate the slides in

2XSSC at 56°C for 2hrs. Stain the slide with 2% Giemsa's stain with Sorenson's buffer.<sup>[9,10]</sup> The sister chromatid exchanges were used to determine if the *Bacopa monnieri* plant extracts produced any genotoxicity. Slides were prepared from each blood culture and No. of sister chromatid exchanges in the centromere was counted in 1000 M2 Metaphases in the chromosome spreads per culture was assessed to score for any chromosomal aberrations that may be induced due to the *Bacopa monnieri* plant extracts. SCEs can be calculated using the formula as follows:

$$\text{SCE} = \frac{\text{Total no. of sister chromatid exchanges observed}}{\text{Total no. of metaphases observed with sister chromatid exchanges}} \times 100$$

## RESULTS AND DISCUSSIONS

### Determination of mitotic index (MI)

For the calculation of mitotic index, number of dividing cells has to be counted for every 1000 cells counted. From the slides, we have observed the 2 types of dividing cells

Prophase-stage of mitosis in which the chromatin condenses (it becomes shorter and fatter) into a highly ordered structure called a chromosome in which the chromatin becomes visible. This process, called chromatin condensation

Metaphase - stage of mitosis in the eukaryotic cell cycle in which condensed & highly coiled chromosomes, carrying genetic information, align in the middle of the cell before being separated into each of the two daughter cells.


From the observed slides, dividing and non dividing cells were counted to determine the mitotic index for the assessment of the cytotoxicity in the control and *Bacopa monnieri* plant extracts of concentration 500 & 1000 µg/ml using three blood samples for the accuracy. The Mean Mitotic Index (MI) values of the cultures which are exposed to two doses of plant extracts and control are obtained as shown in the TABLE 1. The mean value of MI of the control is 4.39 where as the *Bacopa monnieri* extracts have the mean value of MI of the 500µg/ml is 4.09 and the mean value of MI of the 1000 µg/ml is 3.62. MI of *Bacopa monnieri* plant extracts was not much different from the MI of

control.

From the Figure 1 of MI Vs Type of culture, obtained equation was  $y = -0.385x + 4.803$  and  $R^2 = 0.984$ , which indicates that MI of *Bacopa monnieri* plant extracts were not significantly different from the MI of control.

**TABLE 1 : Mitotic index of *Bacopa monnieri* plant extracts (500 & 1000 µg/ml) and control using three blood samples.**

S. No	Sample	Control	Bacopa 500 µg/ml	Bacopa 1000 µg/ml
1.	Sample1	4.44	4.36	3.67
2.	Sample2	4.23	3.92	4.21
3.	Sample3	4.5	3.98	2.97
4.	Mean	4.39	4.09	3.62



**Figure 1 : Graphical representation of the overall mean mitotic index (MI) of *Bacopa monnieri* extracts (500 & 1000 µg/ml) and control**

### Determination of the frequency of sister chromatid exchanges (SCES)

Slides were prepared from each blood culture and 1000 metaphase chromosome spreads were counted per culture was assessed to score for any chromosomal aberrations (sister chromatid exchanges) that may be induced due to the *Bacopa monnieri* plant extract of concentration 500 & 1000µg/ml using three blood samples for accuracy.

The Mean Sister chromatid exchanges (SCES) values of the cultures which are exposed to two doses of plant extracts and control are obtained as shown in TABLE 2. The mean value of SCES of the control is 5.813 where as the *Bacopa monnieri* extracts have the mean value of SCES of the 500µg/ml is 5.386 and the mean value of SCES of the 1000 µg/ml is 5.35. SCES of *Bacopa monnieri* plant extracts were not


## Regular Paper

much different from the SCES of control.

From the Figure 2 of SCES Vs Type of culture, obtained equation was  $y = -0.239x + 5.989$  and  $R^2 = 0.829$ , which indicates that the frequency of SCES in *Bacopa monnieri* plant extracts were not significantly different from the SCES of control.

**TABLE 2 : Sister Chromatid exchanges (SCES) of *Bacopa monnieri* plant extracts (500 & 1000 µg/ml) and control using three blood samples.**

	Control	Bacopa 500 µg	Bacopa 1000 µg
Sample - 1	5.81	5.6	5.45
Sample - 2	5.75	5.16	5.1
Sample - 3	5.88	5.4	5.5
Mean	5.813	5.386	5.35



**Figure 2 : Graphical representation of the overall mean Sister chromatid exchanges (SCES) of *Bacopa monnieri* extracts (500 & 1000 µg/ml) and control.**


### Determination of cell proliferation kinetics (CPK)

Slides which were prepared for genotoxicity studies are also used to assess cell proliferation kinetics (CPK) and 25 metaphase chromosome spreads per culture were counted to determine the cell proliferation kinetics (CPK) in the control and *Bacopa monnieri* plant extracts of concentration 500 & 1000 µg/ml using three blood samples for the accuracy. The Mean Cell proliferation kinetics (CPK) values of the cultures which are exposed to two doses of plant extracts and control are obtained as shown in TABLE 3. The mean value of CPK of the control is 0.472 where as the *Bacopa monnieri* extracts have the mean value of CPK of the 500µg/ml is 0.46 and the mean value of CPK of the 1000 µg/ml is 0.468. CPK of *Bacopa monnieri* plant extracts were not much different from the CPK of control.

From the Figure 3 of CPK Vs Type of culture, obtained equation was  $y = -0.0055x + 0.478$  and  $R^2 = 0.9758$ , which indicates that CPK of *Bacopa monnieri* plant extracts were not significantly different from the CPK of control.

**TABLE 3 : Cell proliferation kinetics (CPK) of *Bacopa monnieri* plant extracts (500 & 1000 µg/ml) and control using three blood samples.**

Sr. No	Sample ID	Control	CPK (500 µg/ml)	CPK (1000 µg/ml)
1	Sample-1	0.47	0.49	0.49
2	Sample-2	0.49	0.45	0.45
3	Sample-3	0.48	0.41	0.49
	Mean	0.472	0.46	0.468



**Figure 3 : Graphical representation of the overall means Cell Proliferation Kinetics (CPK) of *Bacopa monnieri* extracts (500 & 1000 µg/ml) and control.**

## CONCLUSION

In the present studies, *Bacopa monnieri* methanolic crude extract was prepared by soxhlet extraction and then evaluated for the cytotoxicity and genotoxicity in human peripheral blood lymphocyte cultures. However, working with crude extracts also means working with complex mixtures of biologically active compounds. Some of these compounds can be Cytotoxic and/or Genotoxic; others can be Cytoprotective and/or antigenotoxic.

Mitotic Index (MI) Cell Proliferation Kinetics (CPK) and Sister-chromatid exchanges (SCE) were scored to measure the cytotoxic and genotoxic effects of the *Bacopa monnieri* extract in cultures set up from the three different healthy donors. From the results, the MI and frequency of SCES of *Bacopa*

*monnieri* methanolic extracts are almost similar to that of control which indicates that the *Bacopa monnieri* plant extracts has no significant cytotoxicity and genotoxicity effects in cultured human peripheral blood lymphocytes.

### REFERENCES

- [1] Abhishek Mathur, Satish K. Verma, Reena Purohit et al.; 'Dual pharmacological investigation of *Bacopa monnieri* on the basis of antioxidant, antimicrobial and anti-inflammatory properties', Journal of Chemical and Pharmaceutical Research, **2(6)**, 191-198.
- [2] V. Elangovan, S. Govindasamy, N. Ramamoorthy, K. Balasubramanian; 'In vitro studies on the anti-cancer activity of *Bacopa monniera*', Fititerapia, **66**, 211-215 (1995).
- [3] T.P. Kannan et al.; 'Mitotic index and chromosomal analyses for hydroxyapatite implantation in rabbits', Archives of Orofacial Sciences, **1**, 15-20 (2006).
- [4] P. Maria et al.; 'Genotoxicity of commercial petrol samples in cultured human lymphocytes', Rev. Int. Contam. Ambient, **13(1)**, 15-21 (1997).
- [5] Meena Vangalapati, Srinivas Rao Bmmidi, Sharan Suresh Volluri, Seema Chaitanya Chippada, Sumanjali Avani gadda; 'A review on pharmacological studies of *Bacopa Monnieri*', Journal of Chemical, Biological and Physical Sciences, **1(2)**, 250-259 (2011).
- [6] M.J. Plewa, E.D. Wagner; 'Activation of promutagens by green plants', Annual Review of Genetics, **27**, 93-113 (1993).
- [7] O. Schimmer et al.; 'An evaluation of the 55 commercial plant extracts in the ames mutagenicity test', Pharmazie, **49(6)**, 448-451 (1994).
- [8] Sharan Suresh Volluri, Srinivas Rao Bammidi, Seema Chaitanya Chippada, Meena Vangalapati; 'In-vitro anti-arthritis activity of methanolic extract of *Bacopa Monniera*', International Journal of Chemical, Environmental and Pharmaceutical Research, **2(2,3)**, 156-159 (2011).
- [9] Tirtha Ghosh, Tapan Kumar Maity, Mrinmay Das et al.; 'In vitro antioxidant and hepatoprotective activity of ethanolic extract of *Bacopa Monnieri* linn. aerial parts', Iranian Journal of Pharmacology and Therapeutics, **6**, 77-85 (2007).
- [10] T. Zink, J. Chaffin; 'Herbal health products: what family physicians need to know', American Family Physician, **58(5)**, 1133-1140 (1998).