

Extraction of beta-carotene as natural dyes from *Cucumis melo* via soxhlet extraction method and its applications for coloring and decorating sweets

Hassan Abdalla Almahy*

Department of Chemistry, Faculty of Science, University of Sudan.

*Corresponding author: Hassan Abdalla Almahy, Department of Chemistry, Faculty of Science, University of Sudan, E-mail: abuamna4@hotmail.com.

Received: October 24, 2021; Accepted: November 8, 2021; Published: November 15, 2021

Abstract

Natural dyes are pronounced as an interesting alternative to fabricated dyes. Accordingly, the attractive demand to develop the extraction techniques of the best ingredients of solid and liquid for natural dyes colorants from fruits materials and their application in human life. Extraction is processed using solvents or using modern techniques like ultrasonication and microwave. In this respects, the extraction of natural colorant from *Cucumis melo* has been investigated as a model. Determination of the structure of active ingredients was performed conducted by UV spectroscopy and gravimetric analysis. Through the analysis of the results, it will be confirmed that the dye extracted from *Cucumis melo* of intense color is beta-carotene, which is considered as one of the most colored dyes. Therefore, we recommend using such dyes in the industrial fields especially for coloring and decorating sweets.

Keywords: *Cucumis melo*; extraction; ultra-sonication; β -carotene; colorants; sweets.

Introduction

Recently, the consumer attracted for essential materials instead of fabricated analogous which need an extensive role in the food industry. Particularly, due to dietary supplementation and healthy equities associated there from. Such as *Cucumis melo* as a member of Cucurbitaceae family, the fruit is recognized locally in Kingdom of Saudi Arabia as "Shamam" and it is widely established as the consequence of its calyces. Usually, they are used in large scale to construct cold beverages in several worlds' countries, additionally in phytotherapeutic applications [4, 13]. Several pharmacological activities have been effectively cited, such as antihypertensive [21] and cardio protective agents (Napier et al ., 201 ; Paris et al . , 2012), hepatoprotective (

Citation: Hassan Abdalla Almahy. Extraction of beta-carotene as natural dyes from *Cucumis melo* via soxhlet extraction method and its applications for coloring and decorating sweets. Int J Chem Sci. 2021;19(10):413

© 2021 Trade Science Inc

Sebastian et al., 2010), an inhibitor against porcine pancreaticamylase (Vouldoukis et al., 2008; Lin et al., 2007), sedative (Ali et al., 2005) and antioxidant agents (Hansawasdi et al., 2000; Chen et al., 2004). Moreover, *Cucumis melo* has attained an essential position in the soft drinks market; its commercial extract preparations are currently advertised as supplements laying for their attractive health benefits and as a replacement colorant of the fabricated dyes.

The intensive red colorant in the yellow calyces of *Cucumis melo* is beta-carotene and it was usually analyzed and detected using mass spectroscopy (Zhang et al., 2004; Chirinos et al., 2006). The distinctions for beta-carotene are correlated by how many the -OH and -COOH groups, the nature, location, and the amount of sugars (e.g. D-glucose, D-galactose, etc.) incorporated to the molecule as well as aromatic acids coupled to sugars in the molecule (Bhuyan and Saikia 2004). Distinguished glycones are of eighteen naturally existing most plants are pronounced as delphinidin (Dp), pelargonidin (Pg), peonidin (Pn), cyaniding (Cy), malvidin (Mv) and petunidin (Pt) (Bednar et al., 2003; Ichianagi et al., 2000; Calvo et al., 2008). The most prevalent glycosides in nature are the non-methylated three anthocyanidins (Cy, Dp and Pg) (Arráez-Román et al., 2009) and their structures were elucidated using spectroscopic and chemical transformation methods (Carrasco et al., 2006; Bednar et al., 2005).

Additionally, organic dyes, whether natural or synthetic are compounds of molecular structure that contain π -electrons. It can be affected by light beams that caused the electron oscillation between lower and higher energy levels. The color appearance depends on the following (Hou et al., 2007; Gradinaru et al., 2002; Prior et al., 1998).

- The chromophoric groups found in the molecular structure.
- The conjugated double bonds which help the molecule polarization.
- The auxochromophoric groups.

Interestingly, beta-carotene investigation by botanists and plant physiologists was concerned in respect to their importance as pollination attractants and photoprotective agents. Kong *et. al.*, introduced beta-carotene as a very useful colorant in taxonomic studies, realizing their accessible priority in affecting the color performance of the fresh and treated fruits and vegetables (Kong et al., 2003). It was reported that first started studying these compounds to overcome the color degeneration of strawberry preserves fruit juice concentrates. Subsequent works were constructed to define their utility as natural foodstuff colorants and to authorize fruit juice, which may prevent disease (Almahy 2011; Almahy et al., 2015).

The main target of the present study is to apply a modern selective extraction method, separation and identification using spectroscopic methods in respect to determining a qualitative estimation of beta-carotene in *Cucumis melo*. The above findings encouraged us to extract the β -carotene as natural dyes from *Cucumis melo*, investigate its chemical composition and exploring its applications for coloring and decorating sweets in Kingdom of Saudi Arabia.

Experimental Methods

Plant materials

The fresh fruit *Cucumis melo* were collected, the fruit collected was further shelled and dried organs were ground to a reduced fine powder using Thomas-Willey Milling Machine.

Extraction process

24.0 g of homogenized *Cucumis melo* powder dry parts were immersed in one liter of methanol and stirred at room temperature for 48 hours. The reaction mixture was filtered off and the filtrate was mixed with 40.0 g Amberlite XAD-2 (pore size 9 nm, particles size 0.3-1.2 mm) and stirred for 40 minutes at room temperature. Then packing the particles in a glass column (64cm x 42cm) and the column was washed with deionized water until a colorless solution was obtained. Elution of the carotenoids remained absorbed on the column using methanol (1L) and glacial acetic acid (1%). The intense red solution obtained was subjected to dryness under reduced pressure.

Dyeing process

The method of dyeing procedure in each case depends on the type of the foodstuff and the dyeing conditions that were performed by varying: temperature, pH, the addition of other additives and duration of treatments. Each time 20.0g of samples are used adding ingredients and controlling the required conditions.

Procedures

- Start with cool foodstuff e.g. cakes; protect the surface, not to decorate by covering with a sheet of paper.
- Prepare the coloring dye.
- Place the coloring dye on the places to be decorated.
- Keep in liquid for up to 5 minutes; leave it longer for a darker hue.

Carefully remove the sheet of paper to get the decoration.

Results and Discussions

UV-VIS spectroscopic analysis of the natural dye derived from *Cucumis melo* was obtained at the wavelength of 480 nm. Regarding the absorbance of natural dye extract derived from both magnetic stirring and ultrasound. Are marked 13% enhancement in percent of yield for the extract accompanying the use of ultrasound in comparable to the control process was recorded.

The color reflectance of dyed silk fabrics using *Cucumis melo* extracts belongs to boiling and solvent extraction techniques. Indeed, the same *Cucumis melo* extracted dye was blended with various mordant to give many colors. Instantly, fabrics dyed using iron as mordant produced the darkest color in comparison with the other mordants for both dyes from *Cucumis melo*.

β -carotene is a red-orange organic compound and is chemically allocated as a distinctive terpenoid, derived from isoprene moiety; it is biosynthesized from geranylpyrophosphate 27. β -carotene is characterized for the general class of carotenes by the β -rings at both terminals of the molecule (Hou et al., 2007; Gradinaru et al., 2002). β -carotene is considered as a non-polar dye, so it can be isolated via the non-polar solvation (Prior et al., 1998). It is characterized by high conjugation, deeply colored, and as a hydrocarbon which lacking functional groups, it is aliphilic (Tsuda et al., 2002; Kong et al., 2003).




FIG.1. Extracted Dye from *Cucumis melo*

The dye extracted from *Cucumis melo* of intense color is beta-carotene (FIG.1.), which is considered as one of the most colored dyes. Our results were revealed that, an attractive modification in the yield percent for colorant isolated has been observed in case of using ultrasound. It can be interpreted owing to the extent degree of binding of coloring matter related to plant cell membranes. On the other hand, the chemical constituents of plant species is another important factor which responsible for the chromophoric moiety and its solubility origin. Regarding the model selection of the plant, the basic chromophore group is carotenoid possess the OH group. According to Almahy (Almahy 2011), "The hydroxyl group is expected to be extracted better applying aqueous solvation using organic solvents". Conversely, in the case of Green wattle, the main chromophore investigated is poly phenolic (kaempferol), which is preferred to be performed better using water. Hence, better yields are observed for

some plants as compared to Greenwattle using water as solvent (Almahy et al., 2015).

Conclusions

β-carotene is providing an eco-friendly and safe facility for the coloring of food stuff and other materials. Application of ultrasound was proved to perform the extraction process for a larger extent owing to the better leaching of natural dyes of the plant cell membranes; consequently a mass transfer to solvent was assisted by acoustic cavitation's assisted by ultrasound. Additional process parameters such as solvent system, temperature, ultrasound power, amount of dye material etc., are attracting parameters for our future investigation work.

REFERNCES

1. Ali BH, Wabel AL, Blunden N. Ultrasound-assisted enhancement in natural dye extraction from beetroot for industrial applications and natural dyeing of leather. *Phytother. Res.* 2005;19:369-375
2. Almahy HA. Anthocyanin (Natural dye) Sensitizing TiO₂ in Photodegradation Processes. The Fourth International Chemistry Conference. Riyadh. Saudi Arabia. 2011;November:19-21
3. Almahy HA, Abdel-Razik HH, Elbadry YA, et al. Ultrasonic extraction of anthocyanin as natural dyes from Hibiscus Sabdariffa(karkade) and its application on dyeing foodstuff and beverages. *American J Biological and Pharmaceutical Research (AJBPR)*. 2015;125-131
4. Amaro AL, Beaulieu JC, Grimm CC. Effect of oxygen on aroma volatiles and quality of fresh-cut cantaloupe and honeydew melons. *Food Chem.* 2012;130(1):49-57
5. Amos S, Binda L, Chindo BA, et al. Potential applications of the oxidoreductive enzymes in the decolorization and detoxification of textile and other synthetic dyes from polluted water. *Pharm. Biol.* 2003;41:325-329
6. Arráez-Román D, Cortacero-Ramírez S, Segura-Carretero A, et al. Natural dye-yielding plants and indigenous knowledge on dye preparation in Arunachal Pradesh Northeast India. *Electrophoresis.* 2009; 47:2197-2207
7. Bednar P, Papoušková B, Müller L, et al. Synthetic organic food coloring agents and their degraded products effects on human and rat cholinesterases. *J. Sep. Sci.* 2005;28:1291-1299
8. Bednar P, Tomassi A, Presutti V, et al. Applications of natural dyes extracted from plants for coloring and decorate foods. *Chromatographia.* 2003;58:283-287
9. Bhuyan R, Saikia CN. Isolation of color components from native dye-bearing plants in Northeastern India, *Bioresour. Technol.* 2004;95:363-372
10. Brat P, George S, Bellamy A. Daily Polyphenol Intake in France from Fruit and Vegetables. *J. Nut.* 2006; 136(9):2368-73
11. Calvo D, Saenz-Lopez R, Fernandez-Zurbano P, et al. Extraction of natural dyes from cantaloupe species. *Anal. Chim. Acta.* 2008;524:207-213
12. Carrasco A, Arráez-Román D, Segura-Carretero A, et al. Uses of natural dyes extracted from carrot. *Electrophoresis.* 2006;27:2182-2196
13. Chen CC, Chou FP, Holy YC, et al. Natural dye-yielding plants and indigenous knowledge on dye preparation in Arunachal Pradesh Northeast India. *J. Sci. Food. Agric.* 2004;84:1989-1996

14. Chirinos R, Campos D, Betalleluz I, et al. Toxicological effect of indole and its azo dye derivatives on some microorganisms under aerobic conditions. *J. Agric. Food Chem.* 2006;54:7089-7097
15. Daniel CR, Prabhakaran D, Kapur KA. Cross-sectional investigation of regional patterns of diet and cardio-metabolic risk in India. *Nutr. J.* 2011;10(12). doi: 10.1186/1475-2891-10-12
16. Gonda I, Bar E, Portnoy V. Branched-chain and aromatic amino acid catabolism into aroma volatiles in Cucumis melo L. fruit. *J. Exp. Bot.* 2010;61(4):20-24
17. Gradinaru G, Biliaderis CG, Kallithraka S, et al. Ultrasound-assisted enhancement in wattle bark (*Acacia mollissima*) vegetable tannin extraction for leather processing. *Food Chem.* 2002;82:422-434
18. Hansawasdi C, Kawabata J, Kasai T. Extraction of natural dye from African marigold flower (*Tagetes erecta*) for textile coloration. *Biosci. Biotechnol. Biochem.* 2000;64:1041-1043
19. Hou DX, Tong X, Terahara N, et al. Aromatic amines from azo dye reduction: status review with emphasis on direct UV spectrophotometric detection in textile industry wastewaters. *Arch. Biochem. Biophys.* 2007;448:244-248
20. Ichihara T, Kashiwada Y, Ikeshiro Y, et al. Comparison between natural and synthetic dyes. *Chem. Pharm. Bull.* 2000;52:226-229
21. Ismail HI, Chan KW, Mariod AA. Phenolic content and antioxidant activity of cantaloupe (*Cucumis melo*) methanolic extracts. *Food Chem.* 2011;119(2):643-647
22. Ismail M, Mariod A, Bagalkotkar G. Fatty acid composition and antioxidant activity of oils from two cultivars of Cantaloupe extracted by supercritical fluid extraction. *J. Grasas Aceites.* 2009;61(1):37-44
23. Kong JM, Chia LS, Goh NK, et al. Application of anthocyanin's extracted from malevaceae. *Phytochemistry.* 2003;64(5):923-933
24. Kourkoutas D, Elmore JS, Mottram DS. Comparison of the volatile compositions and flavour properties of cantaloupe. *Food Chem.* 2006;97(1):95-102
25. Lin HH, Chen JH, Kuo WH, et al. Mutagenic and carcinogenic potential of a textile azo dye processing plant effluent that impacts a drinking water source. *Chem. Biol. Interact.* 2007;165:59-75
26. Melo ML, Narain N, Bora PS. Characterisation of some nutritional constituents of melon (*Cucumis melo* hybrid seeds). *Food Chem.* 2008;68(4):411-414
27. Naito Y, Akagiri S, Uchiyama K. Reduction of diabetes-induced renal oxidative stress by a cantaloupe melon extract biopolymers, oxykine, in mice. *Biofactors.* 2005;23(2):85-95
28. Napier AB, Crosby KM, Park SO. Identifying Molecular Markers Correlating with High Beta-carotene Content in Muskmelon. *HortScience.* 2014;41:1049
29. Paris HS, Amar Z, Lev E. Medieval emergence of sweet melons, *Cucumis melo* (Cucurbitaceae). *Ann. Bot.* 2012;110(1):23-33
30. Prior R, Cao G, Martin A. Antioxidant capacity as influenced by total phenolic and anthocyanin's content, maturity, and variety of *Vaccinium* species. *J. Agric. Food Chem.* 1998;46(7):2686-2693
31. Sebastian P, Schaefer H, Telford IR. Cucumber (*Cucumis sativus*) and melon (*Cucumis melo*) have numerous wild relatives in Asia and Australia and the sister species of melon is from Australia. *Proc. Natl.*

Acad. Sci. USA. 2010;107(32):14269-73

32. Tsuda T, Horio F, Osawa T. Applications and decolorization of dyes from plants. J. Nutr.Sci. Vitaminol. 2002;48(4):305-310
33. Vouldoukis I, Lacan D, Kamate C. Antioxidant and anti- inflammatory properties of a Cucumis melo extract rich in superoxide dismutase activity. J. Ethnopharmacol. 2008;94(1):67-75
34. Zhang Z, Kou X, Fugal K, et al. Determination of banned azo dyes in consumer goods. J. Agric. Food. Chem. 2004;52:688-691