

## Research on Plant Hormones: A Review

**Taagore Anandakumar B\***

<sup>1</sup>Department of Biotechnology, Vydehi Institute of Biotech Sciences, Karnataka, India

\* **Corresponding author:** Taagore Anandakumar B, Department of Biotechnology, Vydehi Institute of Biotech Sciences, Karnataka, India E-mail: tagore4890@yahoo.co.in

**Received:** February 02, 2017; **Accepted:** February 20, 2017; **Published:** February 22, 2017

### Abstract

I am certain you've heard the familiar adage about how one rotten one riches the bundle? Indeed, it's actual, and the cause is a hormonal unevenness. In nature, the primary ready apple of the season drops to the ground and starts to deteriorate. Amid the disintegration procedure, the apple discharges a gas called ethylene. Ethylene is a plant development hormone, or phytohormone, that triggers close-by apples to age and tumble to the ground. The sweet possess a scent reminiscent of every one of those decaying apples pulls in searching creatures, who eat the apples and spread the seeds far and wide

**keywords:** *Hormones; Phytohormone; Auxins; Cytokinins*

### Introduction

Ethylene and other plant development hormones are crucial to all parts of plant development and advancement, and a little learning and comprehension about them and their uses can enhance your yields.

At the point when a plant sets blooms, the bearing it extends, the measure of its natural products, when it drops them to the ground and practically every other part of plant improvement is controlled by hormones. Natural signs trigger creation of these intense mixes. The hormones are created either in the leaves, blossoms, shoots, roots or natural products, contingent upon which hormone is underway. They are made on the smooth endoplasmic reticulum inside plant cells, then transported through the cell dividers and circled all through the plant. Regardless of whether some portion of the typical life cycle or in light of stress, the interfacing plant hormones are in charge of all development changes.

### Types of Hormones

There are five general classes of hormones: auxins, cytokinins, gibberellins, ethylene, and abscisic corrosive [1-3].

An auxin, indole-3-acetic corrosive (IAA), was the principal plant hormone distinguished. It is made basically in the shoot tips (in leaf primordia and youthful leaves), in incipient organisms, and in parts of creating blossoms and seeds. Its vehicle from cell to cell through the parenchyma encompassing the vascular tissues requires the use of ATP vitality. IAA moves in

one bearing just—that is, the development is polar and, for this situation, descending. Such descending development in shoots is said to be basipetal development, and in roots it is acropetal.

Auxins alone or in mix with different hormones are in charge of numerous parts of plant development. IAA specifically [3-10]:

- Activates the separation of vascular tissue in the shoot peak and in calluses; starts division of the vascular cambium in the spring; advances development of vascular tissue in recuperating of wounds.
- Activates cell prolongation by expanding the versatility of the cell divider.
- Maintains apical strength in a roundabout way by fortifying the generation of ethylene, which specifically restrains sidelong bud development.
- Activates a quality required for making a protein important for development and different qualities for the blend of divider materials made and discharged by dictyosomes.
- Promotes start and development of unusual roots in cuttings.
- Promotes the development of many organic products (from auxin delivered by the creating seeds).
- Suppresses the abscission (partition from the plant) of foods grown from the ground (brought down creation of auxin in the leaf is related with arrangement of the abscission layer).
- Inhibits most blooming (however advances blossoming of pineapples).
- Activates tropic reactions.
- Controls maturing and senescence, lethargy of seeds.

Manufactured auxins are broadly utilized as herbicides, the most generally known being 2,4-D and the infamous 2,4,5-T, which were utilized as a part of a 1:1 blend as Agent Orange amid the Vietnam War and showered over the Vietnam backwoods as a defoliant [11-20].

International Conference on “Plant Science” held on September 21-23, 2015 San Antonio, USA. Some of the prominent speakers are Grace Chen, US Department of Agricultural sciences, Joao Paulo Rodrigues Marques University of Sao Paulo Brazil Grace Chen. They spoke about Plant hormones in the Conference.

The Arab Society for Plant Protection (ASPP) is a non-profit skilled and scholarly organization that works with the promotion of plant hormones and its applications in today’s world. This society works vigorously for the promotion and welfare of Plant development and protections. Right quantity of Plant hormones are essential for the plant development.

VEGETOS journal has been working laboriously towards the development of plant research since 29 years and has successfully published article related to plant research and development. To analyse various substances present in the plant sample the assistance of computer technology is essential. The Journal of Applied Bioinformatics & Computational Biology publishes articles relevant to computational analysis.

The current article combines the study of plant tissue culture performed by different scientists worldwide. Plant hormones have an everlasting impact on the growth and development of plants. Plant hormones have to be well regulated and maintained in right proportion to ensure smooth functioning of the plants [21-30].

### **Cytokinins**

Named in light of their found part in cell division (cytokinesis), the cytokinins have an atomic structure like adenine. Actually happening zeatin, separated first from corn (Zea mays), is the most dynamic of the cytokinins. Cytokinins are found in locales of dynamic cell division in plants—for instance, in root tips, seeds, organic products, and clears out. They are

transported in the xylem and work within the sight of auxin to advance cell division. Contrasting cytokinin:auxin proportions change the way of organogenesis. On the off chance that kinetin is high and auxin low, shoots are framed; if kinetin is low and auxin high, roots are shaped. Parallel bud improvement, which is hindered by auxin, is advanced by cytokinins. Cytokinins additionally defer the senescence of leaves and advance the development of cotyledons.

#### Gibberellins

The gibberellins are boundless all through the plant kingdom, and more than 75 have been disengaged, to date. As opposed to giving each a particular name, the mixes are numbered—for instance, GA1, GA2, et cetera. Gibberellic corrosive three (GA3) is the most across the board and most altogether considered. The gibberellins are particularly copious in seeds and youthful shoots where they control stem extension by animating both cell division and lengthening (auxin fortifies just cell prolongation). The gibberellins are conveyed by the xylem and phloem. Various impacts have been recorded that include around 15 or less of the gibberellic acids. The more noteworthy number with no known impacts evidently are antecedents to the dynamic ones.

Experimentation with GA3 showered on hereditarily predominate plants invigorates prolongation of the smaller person plants to ordinary statures. Normal-height plants splashed with GA3 get to be monsters [31-43].

### **Making Plant Hormones Work for You**

These five classes of hormones work synergistically to trigger all the important physiological procedures plants experience to finish their life cycles and guarantee another era. Cytokinins and auxins offset the change from vegetative to blossoming development, gibberellins and abscisic corrosive cooperate to advance heavier fruiting, and ethylene and auxins organize to bring about the dropping of takes off. Low measurements of gibberellic corrosive advances development, while abnormal states repress it. A mixed drink of chemicals is continually moving through a developing plant, the formula of which is constantly fluctuating.

Talking about formulas, what about the ones that are in some of your most loved develop items? Frequently the genuine dynamic fixing is not recorded on the name, which is the reason that container of potassium sulfate appears to get enchantment going. Plant development hormones are now and then recorded on the name; different circumstances, they are definitely not. In the event that a hormone is made artificially, it is known as a plant development controller, or PGR. Two basic PGRs seen on item names are the auxins indole-3-butyric corrosive (IBA) and naphthaleneacetic corrosive (NAA), which you have likely utilized as a part of establishing hormone items. These two PGRs imitate the normal hormone IAA (indole-3-acidic corrosive), and start the arrangement of a callus and after that root advancement. Utilizing an item with IBA or NAA will guarantee that time spent taking and establishing cuttings is not squandered [44-61].

PGRs can be utilized like an apparatuses in a tool stash, modifying a garden's development however a cultivator sees fit. A typical issue indoor and nursery producers face is coming up short on room, which can be cured by utilizing PGRs that repress stem prolongation. You may have known about items that utilization paclobutrazol, flurprimidol or trinexapac-ethyl, which prevent originates from extending by repressing gibberellin biosynthesis. Contingent upon the planning of their utilization, you can keep vegetative plants shorter for more, or make a blossoming plant create short, tight internodes with expanded sidelong stretching. PGRs can be valuable apparatuses if utilized appropriately, however because of wellbeing and security reasons, some PGRs are intended to be utilized on elaborate products as it were. On the off chance that you choose to utilize them, ensure they are sheltered to use on your planned yield, utilize the suggested security gear, focus on reentry periods, and do a post-reap wash of all deliver.

On the off chance that you need the advantages of a PGR, yet would prefer not to utilize engineered chemicals, Mother Nature has everything made sense of. Intense phytohormones are delivered in plants, growths and green growth. They can be straightforwardly connected to the garden, or removed and amassed into simple to-utilize fluids and powders. A few are in items you may as of now be utilizing.

Willow bark powder is an awesome common attaching hormone because of the high measures of salicylic corrosive present in the bark, which advances root start. This normal wonder specialist additionally has an influence in instigating efficient obtained resistance, bringing about a plant to build up its whole protection framework, and decreasing odds of future ailment or irritation issues. A foliar splash of willow bark water will toughen up your plants and keep them more grounded for more [62-87].

Another common plant development hormone source is yeast, which creates the auxin indole-3-acidic corrosive. Yeast concentrates are likely on the rundown of unlisted fixings in items that make your garden take off. Grown seed teas (SSTs) are beginning to pick up fame among probiotic ranchers. A growing seed is stuffed with abscisic and gibberellic acids, and in addition a bundle of other bioactive chemicals and valuable proteins. These teas are made by splashing seeds, regularly grain or rye, in water until they grow their radicle (the principal bit of root that rises up out of a developing seed), mixing the growing seeds into a slurry, or simply gathering the water they are absorbed. This organically dynamic fluid can be utilized as a part of a root splash or foliar shower. Plants will react to the hormones with root and shoot improvement, cell lengthening and substantial blooming.

Another normal wellspring of plant hormones is kelp. Kelp items contain auxins, gibberellins and cytokinins, making plants develop more leaves, and also fortifying blooming, expanding parallel stretching, growing more roots and separating more cells. Diverse items will have distinctive focuses and proportions relying upon the extraction and fixation handle and the sort of kelp utilized. High cytokinin levels cause mammoth kelp to grow up to 2 feet for each day, however that sort of development may escape hand even in the biggest of distribution center greenery enclosures [88-103].

## Conclusion

Plant development hormones resemble devices in an instrument trunk. Utilizing the correct item at the perfect time permits producers to tailor their greenery enclosures how they see fit. One can prompt vertical development and leaf advancement to meet vegetative development objectives, and after that stop vertical development and advance sidelong fanning and bloom start to complete enormous, or hold clones a long time longer by stopping development by and large. Savvy utilization of phytohormones can convey your garden to the following level—simply recollect keeping an eye out for the rotten ones.

## REFERENCES

1. Kabara JI, Chen JS. Microdetermination of lipids classes after thin-layer chromatography. *Anal Chem.* 1976;48:814-17.
2. Hagio M, Sakurai I, Sato S, et al. Phosphatidylglycerol is essential for the development of thylakoid membranes in *Arabidopsis thaliana*. *Plant Cel Physiol.* 2002;43:1456-64.
3. Wu HC, Chen HM, Shiau CY. Free amino acids and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*). *Food Res Int.* 2003;36:949-57.
4. Hancock REW, Sahl HG. Antimicrobial and host-defense peptides as new anti-infective therapeutic strategies. *Nat Biotechnol.* 2006;24:1551-57.
5. Korhonen H, Pihlanto A. Bioactive peptides: production and functionality. *Int Dairy J.* 2006;16:945-60.

6. <http://www.ncbi.nlm.nih.gov/protein>
7. <http://us.expasy.org/tools/protparam.html>
8. Combet C, Jambon M, Deleage G, et al. Geno3D: Automatic comparative molecular
9. modelling of protein. *Bioinformatics* . 2002;18:213-14.
10. Arnold K, Bordoli L, Kopp J, et al. The SWISS-MODEL workspace: a web-based
11. environment for protein structure homology modelling. *Bioinformatics*. 2006;22:195-01.
12. Agyei D, Danquah MK, Sarethy IP, et al. Antioxidative peptides derived from food proteins. *Free Radicals in Human Health and Disease*. Springer India. 2015
13. Guruprasad K, Reddy BVP, Pandit MW. Correlation between stability of a protein and its dipeptide composition: a novel approach for predicting in vivo stability of a protein from its primary sequence. *Prot Eng*. 1990;4:155-64.
14. Christian JAS, Cerutti L, Hulo N, et al. Prosite: a documented database using patterns and profiles as motif descriptors. *Brief Bioinform*. 2002;3:265-74.
15. Sarika M, Iquebal A, Rai A. In silico analysis and homology model of legume antioxidant proteins. *Onl J Bioinform*. 2012;13:120-29.
16. Duraimurugan P, Jagadish A (2002) Seasonal incidence and effect of weather parameters on the population dynamics of illi thrips *Scirtothrips dorsalis* Hood (Thysanoptera: Thripidae) on rose, Resources management in plant protection during twenty first century, Hyderabad, India: 180-183.
17. Shukla A. Seasonal activity of thrips, scirtothrips dorsalis on chilli crops. *Indian J of Tropical Biodiversity*. 2006;14:177-74.
18. Panickar BK, Patel JR. Population dynamics of different species of thrips on chilli, cotton and pigeonpea. *Indian J of Entomol*. 2001;63:170-75.
19. Nandini, Giraddi RS, Mantur SM, et al. Population dynamics and extent damage of pests of capsicum under protected cultivation. *Karnataka J Agril Sci*. 2010;25:150-51.
20. Vanisree K, Rajashekar P, Rao RG, et al. Seasonal incidence of thrips and its natural enemies on chilli (*capsicum annum* L.) in andhra pradesh. *The Andhra Agril J*. 2011;58:185-91.
21. Goswami BK, Mital A. Management of root-knot nematode infecting tomato by trichoderma viride and <http://trove.nla.gov.au/work/24055138?selectedversion=NBD13452663> aecilomyces lilacinus, *Indian phytopathol*. 2004; 57: 235-36.
22. Jatala P. Biological control of plant parasitic nematodes. *Ann Rev phytopathol*. 1986;24:453-89.
23. Krishnamoorthi R, Kumar R. Management of meloidogyne incognita by paecilomyces lilacinus – influence of soil PH and soil types. *Ann Plant Protect Sci*. 2008;16:236-65.
24. Ioannon N. Soil solarization as a substitute for methyl bromide fumigation in greenhouse tomato production in Cyprus. *Phytoparasitica*. 2002;28:248-56.
25. Jianhua Wu, Peiyue Li, Hui Qian, et al. Assessment of soil salinization based on a low-cost method and its influencing factors in a semi-arid agricultural area, northwest China. *Environ Earth Sci*. 2014;71:3465-75.
26. Gu L F. The Xinjiang production and construction corps cotton drip irrigation under membrane formation and development. *Water saving irrigation*. 2003;1:27-29.

27. Ibranhim MK, Miyazaki T, Nishimura T, et al. Contribution of shallow groundwater rapid fluctuation to soil salinization under arid and semiarid climate. *Arab J Geosci.* 2014;7:3901-11.
28. Aragues R, Medina ET, Zribi W, et al. Soil salinization as a threat to the sustainability of deficit irrigation under present and expected climate change scenarios. *Irrig Sci.* 2015;33:67-79.
29. Dehghanisanij H, Agassi M, Anyoji H, et al. Improvement of saline water use under drip irrigation system. *Agric Water Manage.* 2006;85:232-42.
30. Colombani N, Mastrocicco M, Sole BM, et al. Predicting salinization trends in a lowland coastal aquifer: Comacchio (Italy). *Water Resour Manage.* 2015;29:603-18.
31. Wang GX, Cheng GD. The spatial pattern and influence caused by water resources in arid desert oasis. *Acta Ecologica Sinica.* 2000;20:363-68.
32. Ling H, Xu H, Shi W, et al. Regional climate change and its effects on the runoff of Manas River, Xinjiang, China. *Environ Earth Sci.* 2011;64:2203-13.
33. Meena RS, Ameta OP, Meena BL. Population dynamics of sucking pests and their correlation with weather parameters in chilli capsicum annum. *Crop An In Quart. Js. of Life Sci: The Bioscan.* 2013;8:177-80.
34. Jayakumar M, Rajavel M. Weather based forewarning models for coffee borer incidence. *J Agrometeorol.* 2014;16:140-43.
35. Adak T, Gundappa, Balaji R. Application of thermal indices in predicting mango hopper dynamics in lucknow region of uttar pradesh. *Proceedings of International symposium on "New-dimensions in Agrometeorology for sustainable Agriculture"*. GBPUAT, Pantnagar, India. 2014.
36. Adak T, Singh VK, Ravishankar H. Radiation dynamics and agroclimatic models as a tool to predict impact of climate change on dynamics of flowering in mango. *CISH, Lucknow, India.* 2014.
37. Vargas-Solorzano JW, Carvalho CWP, Takeiti CY, et al. Physicochemical properties of expanded extrudates from colored sorghum genotypes. *Food Res Int.* 2014;55:37-44.
38. Huibin Yu, Song Y, Xi B, et al. Application of chemometrics to spectroscopic data for indicating humification degree and assessing salinization processes of soils. *J Soils Sediments.* 2012;12:341-53.
39. Kaskavalci G. Effect of soil solarization and organic amendments treatment for controlling *meloidogyne incognita* in tomato cultivars in western anatolia. *Turk J Agric.* 2007;31:159-67.
40. Verma KK, Nandal SN. Comparative efficacy of VAM, *glomus fasciculatum* and *g.mosseae* for the management of phosphorus levels. *Nat J Plant improve.* 2006;8:174-76.
41. Al-Hazmi AS, Dawabah AAM. Effect of urea and certain NPK fertilizers on the cereal cyst nematode (*Heterodera avenae*) on wheat. *Saudi J Biol Sci.* 2014;21:191-96.
42. Mojtahedi H, Lownsbery BF. The effects of ammonia-generating fertilizer on *criconemoides xenoplax* in pot cultures. *J nematol.* 1976;8:306-09.
43. Santana-Gomes SM, Dias-Arieira CR, Roldi M, et al. Mineral nutrition in the control of nematodes. *Afr J Agric Res.* 2013;8:2413-20.
44. Seifi S, Bide AK. Effect of mineral fertilizers on cereal cyst nematode *heterodera filipjevi* population and evaluation of wheat world *Appl. Prog.* 2013;3:137-41.
45. Wahundeniya I. Effect of poultry manure on root knot nematode (*meloidogyne spp.*) in tomato (*lycopersicon esculentum* Mill). *Trop Agric.* 1991;147:143-53.

46. Faruk I, Mustafa MH, Dey TK. Effect of two organic amendments and a nematicide for controlling root-knot nematode (*Meloidogyne incognita*) on country bean. *Amer-Eurasian J Agric & Env Sci*. 2012;12:608-15.
47. Noe JP. Crop-and nematode-management systems. *Plant-parasitic nematode interactions*. Amer Soc Agron, Madison Wisconsin, USA.1998.
48. Oka Y, Pivonia S. Use of ammonia-releasing compounds for control of the root-knot nematode *Meloidogyne javanica*. *Nematology*. 2002;4:65-71
49. Sasser JN, Carter CC, Hartman KM. Standardization of host suitability studies and reporting of resistance to root-knot nematodes. North Carolina State Univ. USA. 1984.
50. Kuiper D, Smid A. Genetic differentiation and phenotypic plasticity in *Plantago major* the effect of differences in level of irradiance on growth, photosynthesis, respiration and chlorophyll content. *Physiol Plant*. 1985;60:520-28.
51. Lambers H, Posthumus F, Stulen I, et al. Energy metabolism of *Plantago major* ssp *major* as dependent on the supply of mineral nutrients. *Physiol Plant*. 1981;51:245-52.
52. Mudrik V, Kosobrukhov A, Knyazeva I, Changes in the photosynthetic characteristics of *Plantago major* plants caused by soil drought stress. *Plant Growth Regul*. 2003;40:1-6.
53. Kuiper PJC, Bos M. *Plantago: a multidisciplinary study*. Springer, Gemany. 1992.
54. Gurib-Fakim A. Medical plants: traditions of yesterday and drugs of tomorrow. *Molecul Aspects Med*. 2006;27: 1-93.
55. Chiang LC, Chiang W, Chang MY, et al. Antiviral activity of *Plantago major* extracts and related compounds in vitro. *Antiviral Res*. 2002;5:53-62.
56. Jamilah J, Sharifa AA, Sharifah NRSA. GC-MS analysis of various extracts from leaf of *Plantago major* used as traditional medicine. *World Appl Sci J*. 2012;17:67-70.
57. Noor H, Juing M, Chee BJ, et al. Medicinal properties of *Plantago major*: Hypoglycaemic and male fertility studies. *Pertanika J Trop Agricul Sci*. 2000;23:29-35.
58. Samuelsen AB. The traditional uses, chemical constituents and biological activities of *Plantago major* L. *J Ethnopharmacol* 2000;71:1-21.
59. Chemical, physicochemical, and nutritional evaluation of *Plantago* (*Plantago ovata* Forsk). *Cereal Chem* 83: 358-362.
60. Weryszko-Chmielewska E, Matysik-Woźniak A, Sulborska A, et al. Commercially important properties of plants of the genus *Plantago*. *Acta Agrobot*. 2012;65:11-20.
61. Haddadian K, Haddadian K, Zahmatkash M. A review of *Plantago* plant. *Indian J Tradit Knowl*. 2014;13 681-685.
62. Braam J, Sistrunk ML, Polisensky DH, Xu W, Purugganan M M, et al. Plant responses to environmental stress: regulation and functions of the *Arabidopsis*. *TCH Genes. Planta*. 1997;203:S35-41.
63. Chinnusamy V, Schumaker K, Zhu J-K. Molecular genetic perspectives on cross-talk and specificity in abiotic stress signalling in plants. *J Experiment Biol*. 2004;55:225-36.
64. Larkindale J, Vierling E. Core genome responses involved in acclimation to high temperature. *Plant Physiol*. 2008;146:748-61.

65. Wood KA, Stillman RA, Clarke R T, et al. Understanding plant community responses to combinations of biotic and s of the plant growth cycle. PLOS ONE 7. 2012;e49824.
66. Xiong L, Schumaker KS, Zhu J-K. Cell signaling during cold, drought, and salt stress. The Plant Cell Supplem.2002;S165-83.
67. Gurr MI, Harwood JL, Frayn KN. Lipid biochemistry. Wiley, USA. 2008.
68. Los' DA, Murata N. Membrane fluidity and its roles in the perception of environmental signals. Biochim Bioph Acta. 2004;166:142-57.
69. Golovko TK, Dalke IV, Zakhochiy IG, et al. Functional plasticity of photosynthetic apparatus and its resistance to photoinhibition in *Plantago media*. Russian J Plant Physiol. 2011;58:549-59.
70. Rozentsvet OA, Golovko TK, Bogdanova ES, et al. Polar lipid pool modification in leaves of hoary plantain (*Plantago media* L) plants during their light adaptation under natural conditions. Proceed Academy Sci, Biol Series. 2013;2:152-60.
71. Grebenkina TM, Nesterov VN, Rozentsvet OA. The dynamics of lipid and pigment composition in *plantago media* (Plantaginaceae) during daylight. Veget Resour. 2012;4:565-78.
72. Rozentsvet OA, Golovko TK, Nesterov VN, et al. Lipid content variation in *plantago media* leaves in response to light conditions. SDRP J Earth Sci Environ Studies. 2015;1:1-6.
73. Anikiev VV, Kutuzov FF. A new method of determining the area of a sheet surface in cereals. Russian J Plant Physiol. 1996;8:375-77.
74. Lukatkin AS, Bashmakov DI, Kipaykina NV. A protective role of thidiazuron treatment for cucumber seedlings affected by heavy metals and chill. Russian J Plant Physiol. 2003;50:305-07.
75. Bligh EG, Dyer WJ. A rapid method of lipid extraction and purification. Canad J Biochem Physiol. 1959;37: 911-17.
76. Vaskovsky VE, Latyshev NA. Modified jungnickel's reagent for detecting phospholipids and other phosphorus compounds on thin-layer chromatograms. J Chromatography. 1975;115:246-49.
77. Kates M. Techniques of lipidology: isolation, analysis and identification of lipids Elsevier, New York, USA. 1975.
78. Piao S, Ciais P, Huang Y. The impacts of climate change on water resources and agriculture in China. Nature. 2010;467:43-51.
79. Xu GQ, Wei WS. Climate change of Xinjiang and its impact on eco-environment. Arid land geography . 2004;27:14-18.
80. Shi-Peng Yu, Jing-Song Yang, Guang-Ming Liu, et al. Multiple time scale characteristics of rainfall and its impact on soil salinization in the typical easily salinized area in Huang-Huai-Hai Plain, China. Stoch Environ Risk Assess. 2012;26:983-92.
81. [\(2\)MOD16A2data">https://glcf.umd.edu/data/landsat](https://glcf.umd.edu/data/landsat)
82. <https://ntsg.umt.edu/project/mod16>
83. Chethan S, Malleshi NG. Finger millet polyphenols: Characterization and their nutraceutical potential. American Journal of Food Technology. 2007;2:582-92.
84. Devi PB, Vijayabharathi R, Sathyabama S, et al. Health benefits of finger millet (*Eleusine coracana* L.) polyphenols and dietary fiber: a review. J Food Sci Technol. 2014;51:1021-40.



85. Shobana S, Sreerama YN, Malleshi NG. Composition and enzyme inhibitory properties of finger millet (*Eleusine coracana* L.) seed coat phenolics: Mode of inhibition of  $\alpha$ -glucosidase and pancreatic amylase. *Food Chem.* 2009;115:1268-73.
86. Angioloni A, Collar C. Effects of pressure treatment of hydrated oat, finger millet and sorghum flours on the quality and nutritional properties of composite wheat breads. *J Cereal Sci.* 2012;56:713-19.
87. Martinez M, Rosell C, Gomez M. Modification of wheat flour functionality and digestibility through different extrusion conditions. *J Food Eng.* 2014;143:74-79.
88. Roman L, Martinez MM, Gomez M. Assessing of the potential of extruded flour paste as fat replacer in O/W emulsion: A rheological and microstructural study. *Food Res Int.* 2015;74:72-79.
89. Martinez M, Sanz T, Gomez M. Influence of wheat flour subjected to different extrusion conditions on the rheological behaviour and thermal properties of batter systems for coating. *J Food Sci Technol.* 2015;64: 1309-14.
90. Ding Q, Ainsworth P, Tucker G, et al. The effect of extrusion conditions on the physicochemical properties and sensory characteristics of rice-based expanded snacks. *J Food Eng.* 2005;66:283-89.
91. Ding QB, Ainsworth P, Plunkett P, et al. Effect of extrusion conditions on the functional and physical properties of wheat-based expanded snacks. *J Food Eng.* 2006;73:142-48.
92. Chang YH, Ng PKW. Effect of extrusion process variables on quality properties of wheat-ginseng extrudates. *International Journal of Food Properties.* 2011;14:914-25.
93. Zeng J, Gao H, Li G, et al. Extruded corn flour changed the functionality behaviour of blends. *Czech J Food Sci.* 2011;29:520-27.
94. Alvarez-Martinez L, Koundry KP, Harper JM. A general model for expansion of extruded products. *J Food Sci.* 1988;53:609-15.
95. Fan J, Mitchell JR, Blanshard JMV. The effect of sugars on the extrusion of maize grits: II. Starch conversion. *Int J Food Sci Technol.* 1996;31:67-76.
96. Anderson RA, Conway HF, Pfeifer VF. Gelatinization of corn grits by roll and extrusion cook. *Cereal Science Today.* 1969;14:4-11.
97. Li E, Hasjim J, Dhital S, et al. Effect of a gibberellin-biosynthesis inhibitor treatment on the physicochemical properties of sorghum starch. *J Cereal Sci.* 2011;53:328-34.
98. Chakraborty SK, Singh DS, Kumbhar BK, et al. Process parameter optimization for textural properties of ready-to-eat extruded snack food from millet and legume pieces. *J Texture Stud.* 2009;40:710-26.
99. Stojceska V, Ainsworth P, Plunkett A, et al. The effect of extrusion cooking using different water feed rate on the quality of ready-to-eat snacks made from food by-products. *Food Chem.* 2009;114:226-32.
100. Gutkoski LC, El-Dash AA. Effect of extrusion process variables on physical and chemical properties of extruded oat products. *Plant Foods Hum Nutr.* 1999;54:315-25.
101. Martinez M, Rosell C, Gomez M. Modification of wheat flour functionality and digestibility through different extrusion conditions. *J Food Eng.* 2014;143:74-79.
102. Zhang S, Rodrigue D, Riedl B. Preparation and morphology of polypropylene/wood flour composite foams via extrusion. *Polym Compos.* 2005;26:731-38.

103. Melton TA, Barker K, Koenning SR, et al. Temporal efficacy of selected nematicides on *Meloidogyne* species on tobacco. *J Nematol.* 1995;27:263-72.